

The Sarin-like Organophosphorus Agent bis (isopropyl methyl)phosphonate Induces Apoptotic Cell Death and COX-2 Expression in SK-N-SH Cells

Yosuke ARIMA¹⁾, Kanji YOSHIMOTO²⁾, Akira NAMERA³⁾, Ryosuke MAKITA⁴⁾,
Kazuhiro MURATA³⁾ and Masataka NAGAO^{3, *)}

- 1) Department of Forensic Medicine, Graduate School of Biomedical and Health Sciences, Hiroshima University, Kasumi, Minami-ku, Hiroshima 734-8551, Japan
- 2) Department of Food Sciences and Biotechnology, Faculty of Life Sciences, Hiroshima Institute of Technology, Miyake, Saeki-ku, Hiroshima 731-5193, Japan
- 3) Department of Forensic Medicine, Institute of Biomedical and Health Sciences, Hiroshima University, Kasumi, Minami-ku, Hiroshima 734-8551, Japan
- 4) Department of Rehabilitation, Faculty of Health Sciences, Hiroshima Cosmopolitan University, Ozuka-Higashi, Asa-Minami-ku, Hiroshima 731-3166, Japan

ABSTRACT

Organophosphorus compounds, such as sarin, are highly toxic nerve agents that inhibit acetylcholinesterase (AChE), but not cholinesterase, via multiple mechanisms. Recent studies have shown that organophosphorus compounds increase cyclooxygenase-2 (COX-2) expression and induce neurotoxicity. In this study, we examined the toxicity of the sarin-like organophosphorus agent bis(isopropyl methyl)phosphonate (BIMP) and the effects of BIMP on COX-2 expression in SK-N-SH human neuroblastoma cells. Exposure to BIMP changed cell morphology and induced caspase-dependent apoptotic cell death accompanied by cleavage of caspase 3, caspase 9, and poly (ADP-ribose) polymerase (PARP). It also increased COX-2 expression, while pretreatment with a COX inhibitor, ibuprofen, decreased BIMP-dependent cell death and COX-2 expression in SK-N-SH cells. Thus, our findings suggest that BIMP induces apoptotic cell death and upregulates COX-2 expression.

Key words: Sarin, Organophosphates, Apoptosis, COX-2

Organophosphorus compounds have been used as pesticides and nerve agents worldwide; their toxicity is due to irreversible inhibition of acetylcholinesterase (AChE)^{6,7,11,15,26,43}. The resulting accumulation of acetylcholine in cholinergic synapses results in overstimulation of cholinergic neurons, an event known as an acute cholinergic crisis. Moreover, a recent study showed that organophosphorus compounds cause multiple toxicities²⁷, for example, DNA damage^{3,39}, mitochondria dysfunction²¹, and an inflammatory response⁴. These toxicities cannot be explained only by toxicity with canonical acetylcholinesterase targeting toxicity. Elucidation of various toxic mechanisms of organophosphorus compounds is very important

for optimal treatment.

Organophosphorus compounds can also elicit toxicity independently of their acetyl cholinesterase inhibitory functions; this toxicity is due to mitochondrial dysfunction^{32,44}. Mitochondria have an important role in the production of adenosine triphosphate and reactive oxygen species (ROS), and are also key players in some types of caspase-dependent cell death. Given their central role in several pathways, dysfunctional mitochondria elicit multisystem disorders. For example, oxidative damage alters mitochondrial membrane potential, thereby inducing cell death. Moreover, mitochondria-dependent cell death pathways, such as apoptosis and necrosis, can lead to changes in mitochondrial function and morpholo-

Abbreviations: AChE, acetylcholinesterase; COX, cyclooxygenase; BIMP, bis(isopropyl methyl)phosphonate; ROS, reactive oxygen species; PGE₂, prostaglandin E₂; NSAID, nonsteroidal anti-inflammatory drug; PARP, poly (ADP-ribose) polymerase; DMSO, dimethyl sulfoxide; FBS, fetal bovine serum; LDH, lactate dehydrogenase; RIPA, radioimmunoprecipitation assay; SDS, sodium dodecyl sulfate; HRP, horseradish peroxidase; SEM, standard error of the mean; BPMP, bis(pinacolyl methyl)phosphonate

*Corresponding author at: Department of Forensic Medicine, Graduate School of Biomedical and Health Sciences, Hiroshima University, Japan
Tel: +81-82-257-5171, Fax: +81-82-257-5174, E-mail: nagao@hiroshima-u.ac.jp

gy^{22,33}). Organophosphorus compounds can induce ROS and mitochondrial dysfunction-dependent cell death^{8,25}, and we previously reported that a soman-like organophosphate agent alters the mitochondrial membrane¹⁹. However, the precise mechanisms through which organophosphate compounds induce mitochondria-dependent cell death are unclear.

Inflammation is an important response of the immune system to injury or infection¹⁶; however, excessive inflammatory responses are associated with the pathology of many degenerative diseases^{5,42}. Cyclooxygenase (COX) is an important inflammatory molecule that catalyzes the first step in the formation of prostaglandins from arachidonic acid. There are two isoforms of COX, COX-1 and COX-2; COX-1 is responsible for basal and constitutive prostaglandin synthesis, while COX-2 is an inducible enzyme whose expression is increased in response to various stimuli³⁵. COX-2 expression and production of inflammatory factors, such as prostaglandin E₂ (PGE₂), can induce dopaminergic neuronal cell disorder¹⁷. Interestingly, organophosphorus compounds can modulate inflammatory responses^{10,20,41}, and soman, a compound that is similar to sarin, increases inducible COX-2 expression and inflammation in rats^{2,41}. Thus, organophosphorus compounds may modulate inflammatory responses, such as COX-2 overexpression, thereby contributing to the onset of neurodegenerative disorders. Moreover, ibuprofen, a nonsteroidal anti-inflammatory drug (NSAID) that inhibits both COX-1 and COX-2^{13,31}, has been widely used to relieve pain and inflammation and may also have neuroprotective effects^{5,18,36}.

Sarin is a highly toxic organophosphorus agent, and its synthesis and usage are strictly regulated in Japan. Therefore, in a previous study, we synthesized a sarin-like organophosphate agent, bis(isopropyl methyl)phosphonate (BIMP), that has a phosphonate group (isopropyl methylphosphonate) that is similar to that of sarin. This imparts sarin-like activity on BIMP, as it acts as an AChE antagonist (Fig. 1). After intravenous injection, the LD₅₀ value of BIMP in mice was 0.8 mg/kg, and AChE activity in murine blood and brain cells reduced in a dose-dependent manner²⁸. Consistent with this mechanism of action, BIMP exhibits a toxicity profile similar to that of sarin²⁸. Importantly, BIMP has almost no volatility at room temperature and is easily hydrolyzed in water, making BIMP suitable for experiments in ordinary laboratories^{30,40}. Therefore, BIMP may be a suitable alternative compound for mechanistic studies of sarin toxicity.

In this study, we used BIMP to elucidate the mechanisms through which organophosphorus compounds mediate their cytotoxic effects. To this end, we examined the effect of BIMP on both COX-2 expression and cytotoxicity in the human neuroblastoma cell line, SK-N-SH.




Fig. 1. Chemical structures of sarin and BIMP. BIMP has similar phosphonate group with sarin.

MATERIALS AND METHODS

Reagents

BIMP was prepared as previously described^{28,30}. Anti-caspase 3, anti-cleaved caspase 9, anti-nuclear poly (ADP-ribose) polymerase (PARP), and anti-COX-2 antibodies were purchased from Cell Signaling Technology (Beverly, MA, USA). Anti- β -actin was purchased from Sigma-Aldrich Co. (St. Louis, MO, USA). Dimethyl sulfoxide (DMSO) and ibuprofen were purchased from Wako (Japan).

Cell culture and BIMP treatment

Human neuroblastoma SK-N-SH cells were obtained from the European Collection of Cell Cultures (ECACC, UK) and cultured in DMEM (Invitrogen, Carlsbad, CA, USA) supplemented with 10% fetal bovine serum (FBS; Invitrogen) and 1% penicillin-streptomycin-neomycin (Invitrogen) at 37°C in a humidified atmosphere with 5% CO₂. Culture media were replaced every 3 days. For treatment, BIMP was dissolved in DMSO, and cells were exposed to various concentrations of BIMP for 0-24 hr in medium containing 0.1% FBS.

Measurement of cell viability

Cell viability was measured using the MTT assay (Cell Proliferation Kit I [MTT]; Roche Applied Science, Germany). The cells were seeded in 96-well plates (2×10^4 cells/well) and exposed to BIMP for 24 hr. MTT was then added to the cells, and the plates were incubated at 37°C for 4 hr in a humidified atmosphere with 5% CO₂. Finally, the cells were lysed in lysis buffer, and the quantity of formazan product was measured by determining the absorbance at 490/650 nm, which is directly proportional to the number of living cells in culture. Results were expressed as a percentage of the control cells.

Detection of lactate dehydrogenase (LDH) activity

Cytotoxicity was measured by determining the total LDH content (Cytotoxicity Detection Kit Plus

[LDH]; Roche Applied Science); LDH release was considered an indicator of cytotoxicity. Cells were seeded in 96-well plates (2×10^4 cells/well) and exposed to BIMP for 24 hr. The cellular extract was mixed with LDH reaction mixture (1:1), and the mixture was then incubated in the dark at room temperature for 20-30 min. The reaction was stopped with 0.2N HCl. The absorbance was measured at 490/630 nm in a microplate reader, and the results were expressed as a percentage of the control cells.

Western blot analysis

Cells were seeded in 6-well plates (4×10^5 cells/well) and exposed to BIMP. Cells were then lysed in radioimmunoprecipitation assay (RIPA) lysis buffer (50 mM Tris-HCl [pH 8.0], 150 mM NaCl, 1% sodium deoxycholate, 1% NP-40, and 0.1% sodium dodecyl sulfate [SDS]) containing 1 mM Protease and Phosphatase Inhibitor Cocktail (Thermo Scientific, USA) on ice. Collected cells were broken by sonication on ice and centrifuged at $20,000 \times g$ for 20 min at 4°C. The protein concentration was measured using a BCA protein assay kit (Thermo Scientific). Next, 20 μ g of protein from each sample was separated by SDS-polyacrylamide gel electrophoresis and then transferred to

PVDF membranes. The membranes were incubated in the presence of different primary antibodies at 4°C overnight, followed by incubation with horseradish peroxidase (HRP)-linked secondary antibodies for 1 hr. Immunoreactivity was visualized using VersaDoc (Bio-Rad, Hercules, CA, USA), and protein bands were analyzed by Quantity one (Bio-Rad).

Statistical analysis

Statistical analysis of the data was performed using one-way ANOVA followed by the Tukey-Kramer method. Differences with p values less than 0.05 were considered significant. Error bars represent the mean \pm standard error of the mean (SEM).

RESULTS

BIMP-dependent cytotoxicity reduces SK-N-SH cell viability

To evaluate whether BIMP could induce SK-N-SH cell death independently of its acetyl cholinesterase inhibitory activity, we measured the effects of BIMP on cell viability and cytotoxicity. Cells were exposed to BIMP (100-750 μ M) and incubated for 24 hr. MTT assays revealed that BIMP significantly reduced cell viability in a concentration-dependent manner (Fig. 2A). Additionally, LDH assays showed that BIMP significantly induced cytotoxicity in a concentration-dependent manner (Fig. 2B). Ob-




Fig. 2. BIMP reduced cell viability and induced cytotoxicity in SK-N-SH cells.

Cells were exposed to various concentrations of BIMP (0-750 μ M) for 24 hr in medium containing 0.1% FBS. (A) Cell viability was measured using an MTT assay after 24 hr (mean \pm SEM, n = 6). (B) Cytotoxicity was measured using an LDH release detection assay after 24 hr (mean \pm SEM, n = 6). **p < 0.01 vs. control, one-way ANOVA followed by the Tukey-Kramer method.




Fig. 3. Light microscopy of SK-N-SH cell morphology. (a) Untreated SK-N-SH cells. (b, d) Cells were exposed to 0.1% DMSO for 12 (b) or 24 hr (d). (c, e) Cells were exposed to 500 μ M BIMP for 12 (c) or 24 hr (e). All incubations were performed in medium containing 0.1% FBS. The bar represents 100 μ m.

ervation of cell morphology by light microscopy revealed that treatment with 500 μM BIMP for 12 or 24 hr caused morphological changes and increased the numbers of floating cells, consistent with the onset of cell death (Fig. 3c and 3e). Taken together, these data suggested that BIMP induced SK-N-SH cell death and morphological changes.

BIMP-dependent viability reduction is associated with markers of apoptosis in SK-N-SH cells

To determine the type of BIMP-induced cell death in SK-N-SH cells, we treated them with various concentrations of BIMP (100-750 μM) for 24 hr and then performed western blot analysis for detection of cleaved caspase 3, cleaved caspase 9, and cleaved PARP in whole-cell lysates (Fig. 4). Cleaved caspase 3 expression was significantly increased in a concentration-dependent manner. The same was also true for caspase 9, with a significant increase in cleavage following exposure to 750 μM BIMP. Levels of cleaved PARP were also

significantly increased following exposure to 500 μM BIMP. Interestingly, cleaved PARP expression was lower in cells treated with 750 μM BIMP when compared with the 500 μM dose. This is likely due to the substantial amount of cell death observed at a high BIMP concentration, which leads to general cellular degradation. Owing to its dramatic and reproducible biological effects, we subsequently used 500 μM BIMP in all other experiments. Further analysis revealed that levels of cleaved caspase 3 increased in a time-dependent manner following treatment with BIMP over a 24-hr period (Fig. 4E). Taken together, these results suggested that BIMP induced caspase-dependent apoptotic cell death.

BIMP induces COX-2 expression in SK-N-SH cells

Next, we examined the effects of BIMP on COX protein expression in SK-N-SH cells exposed to 500 μM BIMP over a 24-hr period. Western blot analysis (Fig. 5) revealed that COX-2 expression




Fig. 4. Expression of apoptosis-related proteins following treatment with BIMP in SK-N-SH cells.

(A) Western blot analysis of cleaved caspase 3, cleaved caspase 9, cleaved PARP, and β -actin. Cells were exposed to various concentrations of BIMP (0-750 μM) for 24 hr in medium containing 0.1% FBS. (B) Densitometry analysis of western blots for cleaved caspase 3, (C) cleaved caspase 9, and (D) cleaved PARP. The results were normalized individually to the level of the loading control (β -actin). Data are shown as the mean \pm SEM ($n = 3$). * $p < 0.05$, ** $p < 0.01$ vs. control, one-way ANOVA followed by the Tukey-Kramer method. (E) Western blot analysis of cleaved caspase 3. Cells were exposed to 500 μM BIMP for 0, 1, 3, 6, 12, or 24 hr in medium containing 0.1% FBS.




Fig. 5. Expression of COX-1 and COX-2 following treatment with BIMP in SK-N-SH cells.

(A) Western blot analysis of COX-1, COX-2, and β-actin. The cells were treated with 500 μM BIMP for 0, 1, 3, 6, 12, or 24 hr in medium containing 0.1% FBS. (B, C) Densitometric analysis of western blots for COX-1 (B) and COX-2 (C). The results were normalized individually to the loading control (β-actin). Data are shown as the mean ± SEM (n = 3). *p < 0.05, **p < 0.01 vs. the control, one-way ANOVA followed by the Tukey-Kramer method.

was significantly increased in a time-dependent manner (Fig. 5B). However, COX-1 expression was not significantly altered. These data showed that stimulation with BIMP induced the expression of COX-2, but did not affect the expression of COX-1.

Ibuprofen protects against BIMP-induced cell death

To investigate the effects of ibuprofen on BIMP-induced cell death, we exposed SK-N-SH cells to




Fig. 6. Effects of ibuprofen on BIMP-treated SK-N-SH cells.

(A) Light microscopy of cell morphology. (a) Untreated SK-N-SH cells. (b) Cells were exposed to 500 μM BIMP. (c) Cells were treated with 1 mM ibuprofen. (d) Cells were pretreated with 1 mM ibuprofen for 30 min prior to treatment with 500 μM BIMP. All incubations were performed in medium containing 0.1% FBS for 24 hr. The bar represents 100 μm. (B) Cells were exposed to 500 μM BIMP with or without 1 mM ibuprofen, and after 24 hr, cell viability was measured using an MTT assay (mean ± SEM, n = 6). (C-F) Western blot analysis of COX-2 (C and D) and cleaved caspase 3 (E and F). Cells were pretreated with 1 mM ibuprofen for 30 min and then exposed to 500 μM BIMP for 24 hr in medium containing 0.1% FBS. The results were normalized individually to the loading control (β-actin). Data are the mean ± SEM (n = 3). *p < 0.05, **p < 0.01, one-way ANOVA followed by the Tukey-Kramer method. D = DMSO (0.1%), B = BIMP (500 μM), I = ibuprofen (1 mM).

500 μ M BIMP for 24 hr after pretreatment for 30 min with ibuprofen, and then observed cell morphology by microscopy. This revealed a clear protective effect of the NSAID. However, cell morphology was still altered compared with that of cells treated with 0.1% DMSO (Fig. 6A). Additionally, as shown using MTT assays, ibuprofen itself elicited no toxicity, while the drug significantly decreased BIMP-induced death (Fig. 6B). These data suggest that ibuprofen had the potential to attenuate BIMP-induced toxicity in SK-N-SH cells.

Ibuprofen inhibited BIMP-induced COX-2 expression and apoptotic cell death

Due to its mechanism of action, we next determined whether the protective effects of ibuprofen were associated with changes in the expression of inflammatory markers. Cells were exposed to 500 μ M BIMP for 24 hr after pretreatment for 30 min with 1 mM ibuprofen, and western blot analysis was performed for detection of the pro-inflammatory enzyme, COX-2, and cleaved caspase 3 expression. Interestingly, BIMP-dependent induction of COX-2 and caspase 3 cleavage was significantly decreased by ibuprofen (Fig. 6C-F). These results showed that ibuprofen effectively blocked BIMP-induced COX-2 expression and apoptotic cell death.

DISCUSSION

In this study, we examined the mechanisms through which the sarin-like organophosphate agent BIMP affects viability in neuroblastoma cells. We found that BIMP induced toxicity in SK-N-SH cells, resulting in altered cell morphology and reduced cell viability. In our previous study, we found that the soman-like organophosphate bis (pinacolyl methyl)phosphonate (BPMP) induced mitochondrial vacuolation and stellation in rat astrocytes¹⁹. Here, we observed that BIMP induced similar phenotypes, as it caused loss of neuronal cell projections, and subsequent cell rounding. Thus, our findings suggest that BIMP could induce neurotoxicity.

Previous studies have shown that organophosphorus compounds induce apoptotic cell death^{8,25}. Similarly, our results showed that BIMP induced apoptotic cell death in SK-N-SH cells, as indicated by the induction of cleaved caspase 3, cleaved caspase 9, and cleaved PARP. Caspase 3 is a critical executioner of apoptosis, as it is either partially or completely responsible for the proteolytic cleavage of many key proteins, including PARP²⁹. Cleavage of both these proteins is a hallmark of apoptosis. Additionally, caspase 9 is an important member of the caspase family that further processes other caspases, including caspase 3, upon cytochrome c release from mitochondria³⁴. Therefore, we conclude that BIMP induced mitochondrial dysfunction and




Fig.7. Schematic model for BIMP induced neurotoxicity in SK-N-SH cells

mediated the activation of caspase-9, leading to mitochondria-dependent apoptosis.

Interestingly, while exposure to BIMP did not affect COX-1 expression, there was a significant time-dependent increase in the level of COX-2. COX-2 induces production of the inflammation product PGE₂, which promotes apoptotic cell death in rat cortical cells³⁷. The mechanism through which COX-2 induces neurotoxicity is still unknown. However, high expression of COX-2 has been shown to affect the nervous system adversely. Moreover, consistent with our results, organophosphorus compounds increase COX-2 expression both *in vivo* and *in vitro*^{9,23}, and sarin vapor has been shown to increase prostaglandin levels in guinea pigs²⁴ and rats¹⁴. Our data also indicate that BIMP induced COX-2 expression and caused inflammation, similar to the effects of other organophosphorus compounds. Thus, BIMP may directly induce inflammation in neurons involving COX-2 expression. Further studies are required to elucidate the details of these mechanisms.

In this study, we found that ibuprofen could attenuate BIMP-induced toxicity. Treatment of the cells with ibuprofen inhibited COX-2 expression and reduced levels of cleaved caspase 3, concurrent with suppression of BIMP-induced cell death. In previous studies, treatment with NSAIDs increased survival rates and attenuated brain edema in mice following exposure to soman¹. Moreover, the organophosphorus compound VX induces COX-2 expression and PGE₂ release in rat neurons, while treatment with NSAIDs suppressed these effects and reduced VX-dependent cell death³⁸. Moreover, the NSAIDs, indomethacin and NS-398 reduced COX-2 expression in the human neuroblastoma cell line, MSN¹². Thus, NSAIDs have an important role in attenuating inflammation induced by organophosphorus compounds, which is consistent with our current results. In addition, our findings indicate that anti-inflammatory drugs, such as ibuprofen, could exert neuroprotec-

tive effects by suppressing the cytotoxic effects of BIMP.

In conclusion, we found that the sarin-like organophosphate BIMP induces apoptotic cell death via mitochondrial dysfunction and activation of inflammation, resulting in increased expression of COX-2. Additionally, COX-2 expression played an important role in BIMP-induced neurotoxicity, and this effect was attenuated by ibuprofen (Fig. 7). These results are consistent with those of other recent studies that show the inflammatory response is intimately involved in the toxicity of organophosphorus compounds, and that treatment of inflammation might prevent some of the toxicity of the organophosphorus compounds. However, further studies are required to determine the precise mechanisms of BIMP-induced neurotoxicity and inflammation.

CONFLICT OF INTEREST

The authors declare that there are no conflicts of interest.

ACKNOWLEDGMENTS

This work was carried out at the Analysis Center of Life Science, Natural Science Center for Basic Research and Development, Hiroshima University.

(Received November 17, 2015)

(Accepted December 9, 2015)

REFERENCES

- Amitai, G., Adani, R., Fishbein, E., Meshulam, H., Laish, I. and Dachir, S. 2006. Bifunctional compounds eliciting anti-inflammatory and anti-cholinesterase activity as potential treatment of nerve and blister chemical agents poisoning. *J. Appl. Toxicol.* **26**: 81-87.
- Angoa-Pérez, M., Kreipke, C.W., Thomas, D.M., Van Shura, K.E., Lyman, M., McDonough, J.H., et al. 2010. Soman increases neuronal COX-2 levels: possible link between seizures and protracted neuronal damage. *Neurotoxicology* **31**: 738-746.
- Bagchi, D., Bagchi, M., Hassoun, E. A. and Stohs, S.J. 1995. In vitro and in vivo generation of reactive oxygen species, DNA damage and lactate dehydrogenase leakage by selected pesticides. *Toxicology* **104**: 129-140.
- Banks, C.N. and Lein, P.J. 2012. A review of experimental evidence linking neurotoxic organophosphorus compounds and inflammation. *Neurotoxicology* **33**: 575-584.
- Blasko, I., Apochal, A., Boeck, G., Hartmann, T., Grubeck-Loebenstein, B. and Ransmayr, G. 2001. Ibuprofen decreases cytokine-induced amyloid beta production in neuronal cells. *Neurobiol. Dis.* **8**: 1094-1101.
- Brezenoff, H.E. and Giuliano, R. 1982. Cardiovascular control by cholinergic mechanisms in the central nervous system. *Ann. Rev. Pharmacol. Toxicol.* **22**: 341-381.
- Brimblecombe, R.W. 1977. Drugs acting on central cholinergic mechanisms and affecting respiration. *Pharmacol. Ther.* **3**: 65-74.
- Chan, J.Y., Chan, S.H., Dai, K.Y., Cheng, H.L., Chou, J.L. and Chang, A.Y. 2006. Cholinergic receptor-independent dysfunction of mitochondrial respiratory chain enzymes, reduced mitochondrial transmembrane potential and ATP depletion underlie necrotic cell death induced by the organophosphate poison mevinphos. *Neuropharmacology* **51**: 1109-1119.
- Chapman, S., Kadar, T. and Gilat, E. 2006. Seizure duration following sarin exposure affects neuro-inflammatory markers in the rat brain. *Neurotoxicology* **27**: 277-283.
- Damodaran, T.V., Greenfield, S.T., Patel, A.G., Dressman, H.K., Lin, S.K. and Abou-Donia, M.B. 2006. Toxicogenomic studies of the rat brain at an early time point following acute sarin exposure. *Neurochem. Res.* **31**: 367-381.
- De Candole, C.A., Douglas, W.W., Evans, C.L., Holmes, R., Spencer, K.E., Torrance, R.W., et al. 1953. The failure of respiration in death by anticholinesterase poisoning. *Br. J. Pharmacol. Chemother.* **8**: 466-475.
- Ferrera, P. and Arias, C. 2005. Differential effects of COX inhibitors against β -amyloid-induced neurotoxicity in human neuroblastoma cells. *Neurochem. Int.* **47**: 589-596.
- Gierse, J., Koboldt, C., Walker, M., Seibert, K. and Isakson, P. 1999. Kinetic basis for selective inhibition of cyclo-oxygenases. *Biochem. J.* **339**: 607-614.
- Grauer, E., Chapman, S., Rabinovitz, I., Raveh, L., Weissman, B.A., Kadar, T., et al. 2008. Single whole-body exposure to sarin vapor in rats: long-term neuronal and behavioral deficits. *Toxicol. Appl. Pharmacol.* **227**: 265-274.
- Gupta, R.C., Patterson, G.T. and Dettbarn, W.D. 1991. Comparison of cholinergic and neuromuscular toxicity following acute exposure to sarin and VX in rat. *Fundam. Appl. Toxicol.* **16**: 449-458.
- Hanisch, U.K. and Kettenmann, H. 2007. Microglia: active sensor and versatile effector cells in the normal and pathologic brain. *Nat. Neurosci.* **10**: 1387-1394.
- Hunter, R.L., Dragicevic, N., Seifert, K., Choi, D.Y., Liu, M., Kim, H.C., et al. 2007. Inflammation induces mitochondrial dysfunction and dopaminergic neurodegeneration in the nigrostriatal system. *J. Neurochem.* **100**: 1375-1386.
- Hunter, R.L., Choi, D.Y., Ross, S.A. and Bing, G. 2008. Protective properties afforded by pioglitazone against intrastriatal LPS in Sprague-Dawley rats. *Neurosci. Lett.* **432**: 198-201.
- Isobe, I., Maeno, Y., Nagao, M., Iwasa, M., Koyama, H., Seko-Nakamura, Y., et al. 2003. Cytoplasmic vacuolation in cultured rat astrocytes induced by an organophosphorus agent requires extracellular signal-regulated kinase activation. *Toxicol. Appl. Pharmacol.* **193**: 383-392.
- Johnson, E.A., Dao, T.L., Guignet, M.A., Geddes, C.E., Koemeter-Cox, A.I. and Kan, R.K. 2011. Increased expression of the chemokines CXCL1 and MIP-1a by resident brain cells precedes neutrophil

- infiltration in the brain following prolonged soman-induced status epilepticus in rats. *J. Neurochem.* **8**: 41.
21. **Karami-Mohajeri, S. and Abdollahi, M.** 2013. Mitochondrial dysfunction and organophosphorus compounds. *Toxicol. Appl. Pharmacol.* **270**: 39-44.
 22. **Kim, J.S., He, L. and Lemasters, J.J.** 2003. Mitochondrial permeability transition: a common pathway to necrosis and apoptosis. *Biochem. Biophys. Res. Commun.* **304**: 463-470.
 23. **Lee, J.E., Park, J.H., Jang, S.J. and Koh, H.C.** 2014. Rosiglitazone inhibits chlorpyrifos-induced apoptosis via modulation of the oxidative stress and inflammatory response in SH-SY5Y cells. *Toxicol. Appl. Pharmacol.* **278**: 159-171.
 24. **Levy, A., Chapman, S., Cohen, G., Raveh, L., Rabinovitz, I., Manistersky, E., et al.** 2004. Protection and inflammatory markers following exposure of guinea pigs to sarin vapour: comparative efficacy of three oximes. *J. Appl. Toxicol.* **24**: 501-504.
 25. **Liu, H., Liu, J., Xu, L., Zhou, S., Li, L. and Liu, W.** 2010. Enantioselective cytotoxicity of isocarbofos is mediated by oxidative stress-induced JNK activation in human hepatocytes. *Toxicology* **276**: 115-121.
 26. **Lotti, M.** 2000. Organophosphorus compounds, p.897-925. *In* P. Spencer, H. Schaumburg and A. Ludolph (eds.), *Experimental and clinical neurotoxicology*, second edition. New York.
 27. **Mostafalou, S. and Abdollahi, M.** 2013. Pesticides and human chronic diseases: evidences, mechanisms, and perspectives. *Toxicol. Appl. Pharmacol.* **268**: 157-177.
 28. **Nagao, M., Takatori, T., Matsuda, Y., Nakajima, M., Nijima, H., Iwase, H., et al.** 1997. Detection of sarin hydrolysis products from sarin-like organophosphorus agent-exposed human erythrocytes. *J. Chromatogr. B Biomed. Sci. Appl.* **701**: 9-17.
 29. **Nicholson, D.W., Ali, A., Thornberry, N.A., Vaillancourt, J.P., Ding, C.K., Gallant, M., et al.** 1995. Identification and inhibition of the ICE/CED-3 protease necessary for mammalian apoptosis. *Nature* **376**: 37-43.
 30. **Nijima, H., Nagao, M., Nakajima, M., Takatori, T., Matsuda, Y., Iwase, H., et al.** 1999. Sarin-like and soman-like organophosphorous agents activate PLC γ in rat brains. *Toxicol. Appl. Pharmacol.* **156**: 64-69.
 31. **Robertson, F.M., Parrett, M.L., Joarder, F.S., Ross, M., Abou-Issa, H.M., Alshafie, G., et al.** 1998. Ibuprofen-induced inhibition of cyclooxygenase isoform gene expression and regression of rat mammary carcinomas. *Cancer Lett.* **122**: 165-175.
 32. **Saleh, A.M., Vijayasarathy, C., Masoud, L., Kumar, L., Shahin, A. and Kambal, A.** 2003. Paraoxon induces apoptosis in EL4 cells via activation of mitochondrial pathways. *Toxicol. Appl. Pharmacol.* **190**: 47-57.
 33. **Skulachev, V.P.** 2006. Bioenergetic aspects of apoptosis, necrosis and mitoptosis. *Apoptosis* **11**: 473-485.
 34. **Slee, E.A., Harte, M.T., Kluck, R.M., Wolf, B.B., Casiano, C.A., Newmeyer, D.D., et al.** 1999. Ordering the cytochrome c-initiated caspase cascade: hierarchical activation of caspases-2, -3, -6, -7, -8, and -10 in a caspase-9-dependent manner. *J. Cell Biol.* **144**: 281-292.
 35. **Smith, W.L., Garavito, R.M. and DeWitt, D.L.** 1996. Prostaglandin endoperoxide H synthases (cyclooxygenases)-1 and -2. *J. Biol. Chem.* **271**: 33157-33160.
 36. **Świątkiewicz, M., Zaremba, M., Joniec, I., Członkowski, A. and Kurkowska-Jastrzębska, I.** 2013. Potential neuroprotective effect of ibuprofen, insights from the mice model of Parkinson's disease. *Pharmacol. Rep.* **65**: 1227-1236.
 37. **Takadera, T., Yumoto, H., Tozuka, Y. and Ohyashiki, T.** 2002. Prostaglandin E₂ induces caspase-dependent apoptosis in rat cortical cells. *Neurosci. Lett.* **317**: 61-64.
 38. **Tenn, C.C., Weiss, M.T., Beaup, C., Peinnequin, A., Wang, Y. and Dorandeu, F.** 2012. Cyclooxygenase-2 contributes to VX-induced cell death in cultured cortical neurons. *Toxicol. Lett.* **210**: 71-77.
 39. **Ündeğer, Ü. and Başaran, N.** 2005. Effects of pesticides on human peripheral lymphocytes in vitro: induction of DNA damage. *Arch. Toxicol.* **79**: 169-176.
 40. **Watanabe, Y., Itoh, T., Shiraishi, H., Maeno, Y., Arima, Y., Torikoshi, A., et al.** 2013. Acute effects of a sarin-like organophosphorus agent, bis(isopropyl methyl)phosphonate, on cardiovascular parameters in anaesthetized, artificially ventilated rats. *Toxicol. Appl. Pharmacol.* **272**: 61-66.
 41. **Williams, A.J., Berti, R., Yao, C., Price, R.A., Velarde, L.C., Koplovitz, I., et al.** 2003. Central neuro-inflammatory gene response following soman exposure in the rat. *Neurosci. Lett.* **349**: 147-150.
 42. **Whitton, P. S.** 2007. Inflammation as a causative factor in the aetiology of Parkinson's disease. *British J. Pharmacol.* **150**: 963-976.
 43. **Wood, J.R.** 1951. Chemical defense. *J. Am. Med. Assoc.* **145**: 1264-1267.
 44. **Xu, W.N., Liu, W.B. and Liu, Z.P.** 2009. Trichlorfon-induced apoptosis in hepatocyte primary cultures of *Carassius auratus gibelio*. *Chemosphere* **77**: 895-901.