

# Existence of tangential limits for $\alpha$ -harmonic functions on half spaces

Yoshihiro Mizuta

## Abstract

Our aim in this paper is to prove the existence of tangential limits for Poisson integrals of the fractional order of functions in the  $L^p$  Hölder space on half spaces.

## 1 Introduction

For  $0 < \alpha < 2$ , Riesz [6] defined the notion of  $\alpha$ -harmonic functions on a domain  $\Omega$  in the  $n$ -dimensional Euclidean space  $\mathbf{R}^n$ , as solutions of the fractional Laplace operators (see also Itô [3] and the book by Landkof [4]).

In the half space  $H = \{x = (x', x_n) \in \mathbf{R}^{n-1} \times \mathbf{R} : x_n > 0\}$ , consider

$$P_\alpha f(x) = c_\alpha \int_{\mathbf{R}^n \setminus \overline{H}} \left( \frac{x_n}{|y_n|} \right)^{\alpha/2} |x - y|^{-n} f(y) dy$$

for a measurable function  $f$  on  $\mathbf{R}^n$ , where  $c_\alpha = \Gamma(n/2) \pi^{-n/2-1} \sin(\pi\alpha/2)$ . Here  $f$  satisfies

$$\int_{\mathbf{R}^n \setminus \overline{H}} |y_n|^{-\alpha/2} (|y| + 1)^{-n} |f(y)| dy < \infty, \quad (1)$$

which is equivalent to

$$P_\alpha |f| \neq \infty.$$

If this is the case, then we see from Remarks 4 and 5 below that  $P_\alpha f$  is  $\alpha$ -harmonic in  $H$ .

Let  $1 < p < \infty$  and  $\beta > 1/p$ . Recently, Bass and You [1] have shown the existence of nontangential limits for  $P_\alpha f$  with  $f \in \Lambda_\beta^{p,\infty}(\mathbf{R}^n)$ , which is the space of  $L^p$  Hölder continuous functions of order  $\beta$ ; more precisely,  $f \in L^p(\mathbf{R}^n)$  and it

---

2000 Mathematics Subject Classification : Primary 31B25, 31B05

Key words and phrases :  $\alpha$ -harmonic functions, (non)tangential limits,  $L^p$  Hölder space

satisfies  $L^p$  Hölder continuity of order  $\beta$ , that is, there exists a positive constant  $C$  such that

$$\left( \int |f(y+h) - f(y)|^p dy \right)^{1/p} \leq C|h|^\beta \quad \text{for all } h \in \mathbf{R}^n. \quad (2)$$

Our aim in this note is to extend their result on the existence of nontangential limits for  $P_\alpha f$ . For  $\gamma > 0$  and  $\xi \in \partial H$ , our approach region is defined by

$$T_\gamma(\xi) = \{x \in H : |x - \xi|^{1+\gamma/(n-\alpha/2)} < x_n\},$$

which is tangential to the boundary  $\partial H$  at  $\xi$ .

**THEOREM.** *Let  $0 < \alpha < 2(1 - 1/p)$  and  $0 < \gamma < \beta - 1/p$ . If  $f$  is a measurable function in  $\Lambda_\beta^{p,\infty}(\mathbf{R}^n)$  satisfying (1), then there exists  $E \subset \partial H$  of  $(n-1)$ -dimensional measure zero such that  $P_\alpha f$  has a finite limit along  $T_\gamma(\xi)$  for every  $\xi \in \partial H \setminus E$ .*

**REMARK 1.** Note that  $|y_n|^{-\alpha/2}$  is locally  $L^{p'}$  integrable on  $\mathbf{R}^n$  where  $p' = p/(p-1)$  if and only if  $0 < \alpha < 2(1 - 1/p)$ . But, in case  $2(1 - 1/p) \leq \alpha < 2$ , we do not know whether  $P_\alpha f$  has a tangential limit at almost every boundary point or not.

**REMARK 2.** Bass and You [2] have also obtained nontangential limit result for  $\alpha$ -harmonic functions in Lipschitz domains. It will be expected that tangential convergence holds in a way similar to the proof of the theorem given later, because they have shown a good estimate for Poisson kernel near the boundary; but the discussions are left to be open.

Our method of proof is carried out directly, as in the discussions of harmonic case (see Landkof [4], Stein [7] and the author [5]).

## 2 Proof of the Theorem

Throughout this note, let  $C$  denote various constants independent of the variables in question. For a locally integrable function  $g$  on  $\mathbf{R}^n$ , we define its integral mean over the ball  $B(x, r)$  centered at  $\xi$  of radius  $r > 0$  by

$$\fint_{B(x,r)} g(y) dy = \frac{1}{\sigma_n r^n} \int_{B(x,r)} g(y) dy,$$

where  $\sigma_n$  denotes the  $n$ -dimensional Lebesgue measure of the unit ball.

Let  $f$  be a measurable function on  $\mathbf{R}^n$  as in the theorem. First note that

$$\begin{aligned} & \fint_{B(\xi,r)} |y_n|^{-\alpha/2} \left| f(y) - \fint_{B(\xi,r)} f(z) dz \right| dy \\ & \leq C r^{-2n} \int_{B(0,r)} \int_{B(0,r)} |y_n|^{-\alpha/2} |f(\xi + y) - f(\xi + z)| dy dz \end{aligned}$$

for  $\xi \in \partial H$ . Further, for  $E \subset \partial H$ , denote by  $|E|$  the  $(n-1)$ -dimensional measure of  $E$ .

To complete a proof of the theorem, we first prepare two lemmas.

LEMMA 1. Let  $0 \leq \alpha < 2(1 - 1/p)$  and  $0 < \gamma < \beta - 1/p$ . For  $f \in \Lambda_\beta^{p,\infty}(\mathbf{R}^n)$ , set

$$E(\gamma) = \left\{ \xi \in \partial H : \limsup_{r \rightarrow 0^+} r^{\alpha/2 - \gamma - 2n} \times \int_{B(0,r)} \int_{B(0,r)} |y_n|^{-\alpha/2} |f(\xi + y) - f(\xi + z)| dydz > 0 \right\}.$$

Then  $|E(\gamma)| = 0$ .

PROOF. For  $M > 0$  and a positive integer  $j$ , set

$$E(\gamma, M, j) = \left\{ \xi \in \partial H : 2^{j(\gamma + 2n - \alpha/2)} \times \int_{B(0,2^{-j})} \int_{B(0,2^{-j})} |y_n|^{-\alpha/2} |f(\xi + y) - f(\xi + z)| dydz > 1/M \right\}.$$

Let  $a > 0$ . Then we have

$$|E(\gamma, M, j) \cap B(0, a)| \leq M 2^{j(\gamma + 2n - \alpha/2)} \times \int_{B(0,a) \cap \partial H} \left( \int_{B(0,2^{-j})} \int_{B(0,2^{-j})} |y_n|^{-\alpha/2} |f(\xi + y) - f(\xi + z)| dydz \right) d\xi.$$

Letting  $w = \xi + y = \xi + (y', y_n)$ , we see from Hölder's inequality and (2) that

$$\begin{aligned} & \int_{\{y_n: |y_n| < 2^{-j}\}} \left( \int_{B(0,a) \cap \partial H} |y_n|^{-\alpha/2} |f(\xi + y) - f(\xi + y + (0, z_n))| d\xi \right) dy_n \\ & \leq \int_{\{w \in B(0,a+1): |w_n| < 2^{-j}\}} |w_n|^{-\alpha/2} |f(w) - f(w + (0, z_n))| dw \\ & \leq C(2^{-j})^{-\alpha/2 + 1 - 1/p + \beta} \end{aligned}$$

for  $y, z \in B(0, 2^{-j})$ . Similarly, letting  $w = \xi + z = \xi + (z', z_n)$ , we obtain

$$\begin{aligned} & \int_{\{z_n: |z_n| < 2^{-j}\}} \left( \int_{B(0,a) \cap \partial H} |f(\xi + y + (0, z_n)) - f(\xi + z)| d\xi \right) dz_n \\ & \leq \int_{\{w \in B(0,a+1): |w_n| < 2^{-j}\}} |f(w + (y' - z', y_n)) - f(w)| dw \\ & \leq C(2^{-j})^{1 - 1/p + \beta} \end{aligned}$$

for  $y, z \in B(0, 2^{-j})$ . Hence it follows that

$$\begin{aligned} & |E(\gamma, M, j) \cap B(0, a)| \\ & \leq CM 2^{j(\gamma + 2n - \alpha/2)} (2^{-j})^{-\alpha/2 + 1 - 1/p + \beta} 2^{-j(n-1)} 2^{-jn} \\ & \quad + CM 2^{j(\gamma + 2n - \alpha/2)} (2^{-j})^{1 - 1/p + \beta} (2^{-j})^{-\alpha/2 + n} 2^{-j(n-1)} \\ & \leq CM 2^{-j(\beta - 1/p - \gamma)}. \end{aligned}$$

Define  $E(\gamma, M) = \bigcap_{k=1}^{\infty} \bigcup_{j=k}^{\infty} E(\gamma, M, j)$ . Then, since  $\beta - 1/p - \gamma > 0$ , we find

$$|E(\gamma, M) \cap B(0, a)| = 0$$

for all  $a > 0$ , which implies that  $|E(\gamma, M)| = 0$ . Noting that  $E(\gamma) = \bigcup_{M=1}^{\infty} E(\gamma, M)$ , we see that  $|E(\gamma)| = 0$  and

$$\lim_{j \rightarrow \infty} 2^{j(\gamma+2n-\alpha/2)} \int_{B(0,2^{-j})} \int_{B(0,2^{-j})} |y_n|^{-\alpha/2} |f(\xi + y) - f(\xi + z)| dydz = 0$$

for every  $\xi \in \partial H \setminus E(\gamma)$  ■

LEMMA 2. Let  $f$  be a function in  $L^p(\mathbf{R}^n)$  satisfying (2). If  $0 < \gamma < \beta - 1/p$ , then

$$A(\xi) = \lim_{r \rightarrow 0^+} \int_{B(\xi, r)} f(y) dy$$

exists and is finite for almost every  $\xi \in \partial H$ . Further,

$$\lim_{r \rightarrow 0^+} r^{\alpha/2-\gamma-n} \int_{B(\xi, r)} |y_n|^{-\alpha/2} |f(y) - A(\xi)| dy = 0$$

for almost every  $\xi \in \partial H$ .

It is well known that if  $f \in L^p(\mathbf{R}^n)$ , then  $A(x)$  exists and is finite for almost every  $x \in \mathbf{R}^n$ . Since  $\partial H$  is of  $n$ -dimensional measure zero, we need an additional condition like  $L^p$  Hölder continuity for  $f$  in the present lemma.

PROOF OF LEMMA 2. For simplicity, set  $A(\xi, r) = \int_{B(\xi, r)} f(y) dy$ . Then, for  $r \leq t \leq 2r$ , note that

$$|A(\xi, t) - A(\xi, r)| \leq Cr^{-2n} \int_{B(0,2r)} \int_{B(0,2r)} |f(\xi + y) - f(\xi + z)| dydz,$$

so that Lemma 1 (with  $\alpha = 0$ ) yields

$$\lim_{r \rightarrow 0^+} r^{-\gamma} |A(\xi, 2r) - A(\xi, r)| = 0$$

for almost every  $\xi \in \partial H$ . In this case,

$$\lim_{k \rightarrow \infty} 2^{k\gamma} \sum_{j=k}^{\infty} |A(\xi, 2^{-j+1}) - A(\xi, 2^{-j})| = 0,$$

which implies that  $A(\xi)$  exists and

$$\lim_{r \rightarrow 0^+} r^{-\gamma} |A(\xi) - A(\xi, r)| = 0.$$

This proves the first part of the result.

Similarly, we see from Lemma 1 that

$$\lim_{r \rightarrow 0^+} r^{\alpha/2 - \gamma - n} \int_{B(\xi, r)} |y_n|^{-\alpha/2} |f(y) - A(\xi, 2r)| dy = 0$$

for almost every  $\xi \in \partial H$ . Note here that

$$\begin{aligned} & \int_{B(\xi, r)} |y_n|^{-\alpha/2} |f(y) - A(\xi)| dy \\ & \leq \int_{B(\xi, r)} |y_n|^{-\alpha/2} |f(y) - A(\xi, 2r)| dy + Cr^{n-\alpha/2} |A(\xi) - A(\xi, 2r)|. \end{aligned}$$

This yields the second result. ■

PROOF OF THE THEOREM. Let  $f$  be as in the theorem. Then Lemma 2 implies that

$$A(\xi) = \lim_{r \rightarrow 0^+} \int_{B(\xi, r)} f(y) dy$$

exists and is finite for almost every  $\xi \in \partial H$ . Moreover we see from Remark 4 below that

$$P_\alpha f(x) - A(\xi) = c_\alpha \int_{\mathbf{R}^n \setminus \bar{H}} \left( \frac{x_n}{|y_n|} \right)^{\alpha/2} |x - y|^{-n} \{f(y) - A(\xi)\} dy.$$

In view of Lemma 2 we see that

$$\lim_{r \rightarrow 0^+} r^{\alpha/2 - \gamma - n} \int_{B(\xi, r)} |y_n|^{-\alpha/2} |f(y) - A(\xi)| dy = 0$$

holds for almost every  $\xi \in \partial H$ . Hence, for given  $\varepsilon > 0$  and  $\xi \in \partial H$ , we assume that

$$r^{\alpha/2 - \gamma - n} \int_{B(\xi, r)} |y_n|^{-\alpha/2} |f(y) - A(\xi)| dy < \varepsilon \quad (3)$$

whenever  $0 < r < r_0$ . Since  $0 < \alpha/2 < 1 - 1/p$ , we see that

$$\lim_{x \rightarrow \xi} \int_{\mathbf{R}^n \setminus \{\bar{H} \cup B(\xi, r_0)\}} \left( \frac{x_n}{|y_n|} \right)^{\alpha/2} |x - y|^{-n} |f(y) - A(\xi)| dy = 0. \quad (4)$$

Letting  $r = 2|x - \xi| < r_0$ , we obtain by (3)

$$\begin{aligned} & \int_{B(\xi, r) \setminus \bar{H}} \left( \frac{x_n}{|y_n|} \right)^{\alpha/2} |x - y|^{-n} |f(y) - A(\xi)| dy \\ & \leq C x_n^{\alpha/2 - n} \int_{B(\xi, r) \setminus \bar{H}} |y_n|^{-\alpha/2} |f(y) - A(\xi)| dy \\ & \leq C \varepsilon x_n^{\alpha/2 - n} r^{\gamma - \alpha/2 + n} \end{aligned}$$

and

$$\begin{aligned}
& \int_{B(\xi, r_0) \setminus \{\bar{H} \cup B(\xi, r)\}} \left( \frac{x_n}{|y_n|} \right)^{\alpha/2} |x - y|^{-n} |f(y) - A(\xi)| dy \\
& \leq C \int_{B(\xi, r_0) \setminus \{\bar{H} \cup B(\xi, r)\}} \left( \frac{x_n}{|y_n|} \right)^{\alpha/2} |\xi - y|^{-n} |f(y) - A(\xi)| dy \\
& \leq C x_n^{\alpha/2} \int_r^{r_0} \left( \int_{B(\xi, t) \setminus \bar{H}} |y_n|^{-\alpha/2} |f(y) - A(\xi)| dy \right) t^{-n-1} dt.
\end{aligned}$$

If  $\gamma - \alpha/2 < 0$ , then

$$\int_{B(\xi, r_0) \setminus \{\bar{H} \cup B(\xi, r)\}} \left( \frac{x_n}{|y_n|} \right)^{\alpha/2} |x - y|^{-n} |f(y) - A(\xi)| dy \leq C \varepsilon x_n^{\alpha/2} r^{\gamma - \alpha/2},$$

which together with (4) implies that

$$\limsup_{x \rightarrow \xi, x \in T_\gamma(\xi)} |P_\alpha f(x) - A(\xi)| \leq C \varepsilon.$$

If  $\gamma - \alpha/2 \geq 0$ , then

$$\int_{B(\xi, r_0) \setminus \{\bar{H} \cup B(\xi, r)\}} \left( \frac{x_n}{|y_n|} \right)^{\alpha/2} |x - y|^{-n} |f(y) - A(\xi)| dy \leq C \varepsilon x_n^{\alpha/2} \log(r_0/r),$$

so that

$$\limsup_{x \rightarrow \xi, x \in T_\gamma(\xi)} |P_\alpha f(x) - A(\xi)| \leq C \varepsilon.$$

Hence it follows that

$$\lim_{x \rightarrow \xi, x \in T_\gamma(\xi)} |P_\alpha f(x) - A(\xi)| = 0,$$

as required. ■

### 3 Further remarks

REMARK 3. Let us consider the Riesz potential

$$U_\alpha \mu(x) = \int |x - y|^{\alpha - n} d\mu(y)$$

for a nonnegative measure  $\mu$  on  $\mathbf{R}^n$ , where  $0 < \alpha \leq 2$  and  $U_\alpha \mu \not\equiv \infty$ . Then it is known that  $U_\alpha \mu$  is  $\alpha$ -superharmonic in  $\mathbf{R}^n$  and  $\alpha$ -harmonic outside the support of  $\mu$ ; for this, see Riesz [6] and Landkof [4].

REMARK 4. Let

$$P_\alpha(x, y) = c_\alpha \left( \frac{x_n}{|y_n|} \right)^{\alpha/2} |x - y|^{-n}$$

for  $x \in H$  and  $y \in \mathbf{R}^n \setminus \overline{H}$ . For  $x_0 \in H$ , we consider the inversion with respect to the ball  $B_0 = B(x_0, R)$  with  $R = (x_0)_n$ . For  $x \in \mathbf{R}^n$ , write

$$x^* = x_0 + R^2 \frac{x - x_0}{|x - x_0|^2}.$$

Set  $\tilde{x}_0 = (x'_0, R/2)$  and  $\tilde{B}(x_0) = B(\tilde{x}_0, R/2)$ . Then, for  $x \in B(x_0, R)$  and  $y \in \mathbf{R}^n \setminus \overline{H}$ , we have

$$\begin{aligned} P_\alpha(x, y) &= c_\alpha R^{-2n} |x_0 - x^*|^{n-\alpha} \left( \frac{|x^* - \tilde{x}_0|^2 - (R/2)^2}{(R/2)^2 - |y^* - \tilde{x}_0|^2} \right)^{\alpha/2} \\ &\quad \times |x^* - y^*|^{-n} |x_0 - y^*|^{n+\alpha}, \end{aligned}$$

so that (A.1) of [4, Appendix] gives

$$\int_{\mathbf{R}^n \setminus \overline{H}} P_\alpha(x, y) dy = 1.$$

REMARK 5. In the same way as above, we find

$$\int_{\mathbf{R}^n \setminus \overline{H}} P_\alpha(x, y) |x_0 - y|^{\alpha-n} dy = |x - x_0|^{\alpha-n}$$

for  $x_0 \in H$ . This is the key result to discuss the  $\alpha$ -harmonicity of  $P_\alpha f$ ; see Itô [3, Lemma 1] and Landkof [4, Lemma 1.13 in Chap.1].

REMARK 6. Let  $0 < \alpha < 2(1 - 1/p)$  and  $\beta p > 1$ . Suppose  $f$  is a measurable function in  $\Lambda_\beta^{p, \infty}(\mathbf{R}^n)$ . Then we can find a set  $E \subset \partial H$  of measure zero such that  $P_\alpha f$  has a finite limit along  $T_\gamma(\xi)$  for every  $\xi \in \partial H \setminus E$  and  $\gamma$  with  $0 < \gamma < \beta - 1/p$ .

## References

- [1] R. F. Bass and D. You, A Fatou theorem for  $\alpha$ -harmonic functions, *Bull. Sci. Math.* **127** (2003), 635–648.
- [2] R. F. Bass and D. You, A Fatou theorem for  $\alpha$ -harmonic functions in Lipschitz domains, *Prob. Th. rel. Fields* 133 (2005), 391-408.
- [3] M. Itô, On  $\alpha$ -harmonic functions, *Nagoya Math. J.* **26** (1966), 205–221.
- [4] N. S. Landkof, *Foundations of modern potential theory*, Springer-Verlag, 1972.
- [5] Y. Mizuta, *Potential theory in Euclidean spaces*, Gakkōtōsyō, Tokyo, 1996.
- [6] M. Riesz, *Intégrales de Riemann-Liouville et potentiels*, *Acta Szeged* **9** (1938), 1–42.

- [7] E. M. Stein, Singular integrals and differentiability properties of functions, Princeton Univ. Press, Princeton, 1970.

*The Division of Mathematical and Information Sciences  
Faculty of Integrated Arts and Sciences  
Hiroshima University  
Higashi-Hiroshima 739-8521, Japan  
E-mail : mizuta@mis.hiroshima-u.ac.jp*