

# PHYSICAL REVIEW D

## PARTICLES AND FIELDS

THIRD SERIES, VOLUME 53, NUMBER 5

1 MARCH 1996

### RAPID COMMUNICATIONS

*Rapid Communications are intended for important new results which deserve accelerated publication, and are therefore given priority in the editorial office and in production. A Rapid Communication in Physical Review D should be no longer than five printed pages and must be accompanied by an abstract. Page proofs are sent to authors, but because of the accelerated schedule, publication is generally not delayed for receipt of corrections unless requested by the author.*

#### Test of the flavor independence of strong interactions

K. Abe,<sup>29</sup> I. Abt,<sup>14</sup> C. J. Ahn,<sup>26</sup> T. Akagi,<sup>27</sup> N. J. Allen,<sup>27</sup> W. W. Ash,<sup>27,\*</sup> D. Aston,<sup>27</sup> N. Bacchetta,<sup>21</sup> K. G. Baird,<sup>24</sup> C. Baltay,<sup>33</sup> H. R. Band,<sup>32</sup> M. B. Barakat,<sup>33</sup> G. Baranko,<sup>10</sup> O. Bardon,<sup>16</sup> T. Barklow,<sup>27</sup> A. O. Bazarko,<sup>11</sup> R. Ben-David,<sup>33</sup> A. C. Benvenuti,<sup>2</sup> T. Bienz,<sup>27</sup> G. M. Bilei,<sup>22</sup> D. Bisello,<sup>21</sup> G. Blaylock,<sup>7</sup> J. R. Bogart,<sup>27</sup> T. Bolton,<sup>11</sup> G. R. Bower,<sup>27</sup> J. E. Brau,<sup>20</sup> M. Breidenbach,<sup>27</sup> W. M. Bugg,<sup>28</sup> D. Burke,<sup>27</sup> T. H. Burnett,<sup>31</sup> P. N. Burrows,<sup>16</sup> W. Busza,<sup>16</sup> A. Calcaterra,<sup>13</sup> D. O. Caldwell,<sup>6</sup> D. Calloway,<sup>27</sup> B. Camanzi,<sup>12</sup> M. Carpinelli,<sup>23</sup> R. Cassell,<sup>27</sup> R. Castaldi,<sup>23,†</sup> A. Castro,<sup>21</sup> M. Cavalli-Sforza,<sup>7</sup> E. Church,<sup>31</sup> H. O. Cohn,<sup>28</sup> J. A. Coller,<sup>3</sup> V. Cook,<sup>31</sup> R. Cotton,<sup>4</sup> R. F. Cowan,<sup>16</sup> D. G. Coyne,<sup>7</sup> A. D'Oliveira,<sup>8</sup> C. J. S. Damerell,<sup>25</sup> R. De Sangro,<sup>13</sup> P. De Simone,<sup>13</sup> R. Dell'Orso,<sup>23</sup> M. Dima,<sup>9</sup> P. Y. C. Du,<sup>28</sup> R. Dubois,<sup>27</sup> B. I. Eisenstein,<sup>14</sup> R. Elia,<sup>27</sup> D. Falciari,<sup>22</sup> C. Fan,<sup>10</sup> M. J. Fero,<sup>16</sup> R. Frey,<sup>20</sup> K. Furuno,<sup>20</sup> T. Gillman,<sup>25</sup> G. Gladding,<sup>14</sup> S. Gonzalez,<sup>16</sup> G. D. Hallewell,<sup>27</sup> E. L. Hart,<sup>28</sup> Y. Hasegawa,<sup>29</sup> S. Hedges,<sup>4</sup> S. S. Hertzbach,<sup>17</sup> M. D. Hildreth,<sup>27</sup> J. Huber,<sup>20</sup> M. E. Huffer,<sup>27</sup> E. W. Hughes,<sup>27</sup> H. Hwang,<sup>20</sup> Y. Iwasaki,<sup>29</sup> P. Jacques,<sup>24</sup> J. Jaros,<sup>27</sup> A. S. Johnson,<sup>3</sup> J. R. Johnson,<sup>32</sup> R. A. Johnson,<sup>8</sup> T. Junk,<sup>27</sup> R. Kajikawa,<sup>19</sup> M. Kalelkar,<sup>24</sup> I. Karliner,<sup>14</sup> H. Kawahara,<sup>27</sup> H. W. Kendall,<sup>16</sup> M. E. King,<sup>27</sup> R. King,<sup>27</sup> R. R. Kofler,<sup>17</sup> N. M. Krishna,<sup>10</sup> R. S. Kroeger,<sup>18</sup> J. F. Labs,<sup>27</sup> M. Langston,<sup>20</sup> A. Lath,<sup>16</sup> J. A. Lauber,<sup>10</sup> D. W. G. Leith,<sup>27</sup> X. Liu,<sup>7</sup> M. Loreti,<sup>21</sup> A. Lu,<sup>6</sup> H. L. Lynch,<sup>27</sup> J. Ma,<sup>31</sup> G. Mancinelli,<sup>22</sup> S. Manly,<sup>33</sup> G. Mantovani,<sup>22</sup> T. W. Markiewicz,<sup>27</sup> T. Maruyama,<sup>27</sup> R. Massetti,<sup>22</sup> H. Masuda,<sup>27</sup> E. Mazzucato,<sup>12</sup> A. K. McKemey,<sup>4</sup> B. T. Meadows,<sup>8</sup> R. Messner,<sup>27</sup> P. M. Mockett,<sup>31</sup> K. C. Moffeit,<sup>27</sup> B. Mours,<sup>27</sup> G. Müller,<sup>27</sup> D. Muller,<sup>27</sup> T. Nagamine,<sup>27</sup> U. Nauenberg,<sup>10</sup> H. Neal,<sup>27</sup> M. Nussbaum,<sup>8</sup> Y. Ohnishi,<sup>19</sup> L. S. Osborne,<sup>16</sup> R. S. Panvini,<sup>30</sup> H. Park,<sup>20</sup> T. J. Pavel,<sup>27</sup> I. Peruzzi,<sup>13,‡</sup> L. Pescara,<sup>21</sup> M. Piccolo,<sup>13</sup> L. Piemontese,<sup>12</sup> E. Pieroni,<sup>23</sup> K. T. Pitts,<sup>20</sup> R. J. Plano,<sup>24</sup> R. Prepost,<sup>32</sup> C. Y. Prescott,<sup>27</sup> G. D. Punkar,<sup>27</sup> J. Quigley,<sup>16</sup> B. N. Ratcliff,<sup>27</sup> T. W. Reeves,<sup>30</sup> P. E. Rensing,<sup>27</sup> L. S. Rochester,<sup>27</sup> J. E. Rothberg,<sup>31</sup> P. C. Rowson,<sup>11</sup> J. J. Russell,<sup>27</sup> O. H. Saxton,<sup>27</sup> T. Schalk,<sup>7</sup> R. H. Schindler,<sup>27</sup> U. Schneekloth,<sup>16</sup> B. A. Schumm,<sup>15</sup> A. Seiden,<sup>7</sup> S. Sen,<sup>33</sup> V. V. Serbo,<sup>32</sup> M. H. Shaevitz,<sup>11</sup> J. T. Shank,<sup>3</sup> G. Shapiro,<sup>15</sup> S. L. Shapiro,<sup>27</sup> D. J. Sherden,<sup>27</sup> C. Simopoulos,<sup>27</sup> N. B. Sinev,<sup>20</sup> S. R. Smith,<sup>27</sup> J. A. Snyder,<sup>33</sup> P. Stamer,<sup>24</sup> H. Steiner,<sup>15</sup> R. Steiner,<sup>1</sup> M. G. Strauss,<sup>17</sup> D. Su,<sup>27</sup> F. Suekane,<sup>29</sup> A. Sugiyama,<sup>19</sup> S. Suzuki,<sup>19</sup> M. Swartz,<sup>27</sup> A. Szumilo,<sup>31</sup> T. Takahashi,<sup>27</sup> F. E. Taylor,<sup>16</sup> E. Torrence,<sup>16</sup> J. D. Turk,<sup>33</sup> T. Usher,<sup>27</sup> J. Va'vra,<sup>27</sup> C. Vannini,<sup>23</sup> E. Vella,<sup>27</sup> J. P. Venuti,<sup>30</sup> R. Verdier,<sup>16</sup> P. G. Verdini,<sup>23</sup> S. R. Wagner,<sup>27</sup> A. P. Waite,<sup>27</sup> S. J. Watts,<sup>4</sup> A. W. Weidemann,<sup>28</sup> J. S. Whitaker,<sup>3</sup> S. L. White,<sup>28</sup> F. J. Wickens,<sup>25</sup> D. A. Williams,<sup>7</sup> D. C. Williams,<sup>16</sup> S. H. Williams,<sup>27</sup> S. Willocq,<sup>33</sup> R. J. Wilson,<sup>9</sup> W. J. Wisniewski,<sup>5</sup> M. Woods,<sup>27</sup> G. B. Word,<sup>24</sup> J. Wyss,<sup>21</sup> R. K. Yamamoto,<sup>16</sup> J. M. Yamartino,<sup>16</sup> X. Yang,<sup>20</sup> S. J. Yellin,<sup>6</sup> C. C. Young,<sup>27</sup> H. Yuta,<sup>29</sup> G. Zapalac,<sup>32</sup> R. W. Zdarko,<sup>27</sup> C. Zeitlin,<sup>20</sup> Z. Zhang,<sup>16</sup> and J. Zhou<sup>20</sup>

(SLD Collaboration)

<sup>1</sup>Adelphi University, Garden City, New York 11530

<sup>2</sup>Istituto Nazionale di Fisica Nucleare Sezione di Bologna, I-40126 Bologna, Italy

<sup>3</sup>Boston University, Boston, Massachusetts 02215

<sup>4</sup>Brunel University, Uxbridge, Middlesex UB8 3PH, United Kingdom

<sup>5</sup>California Institute of Technology, Pasadena, California 91125

<sup>6</sup>University of California at Santa Barbara, Santa Barbara, California 93106

<sup>7</sup>University of California at Santa Cruz, Santa Cruz, California 95064

<sup>8</sup>University of Cincinnati, Cincinnati, Ohio 45221

<sup>9</sup>Colorado State University, Fort Collins, Colorado 80523

<sup>10</sup>University of Colorado, Boulder, Colorado 80309

<sup>11</sup>Columbia University, New York, New York 10027

<sup>12</sup>Istituto Nazionale di Fisica Nucleare Sezione di Ferrara and Università di Ferrara, I-44100 Ferrara, Italy

<sup>13</sup>Istituto Nazionale di Fisica Nucleare Laboratorio Nazionali di Frascati, I-00044 Frascati, Italy

<sup>14</sup>University of Illinois, Urbana, Illinois 61801

<sup>15</sup>Lawrence Berkeley Laboratory, University of California, Berkeley, California 94720

<sup>16</sup>Massachusetts Institute of Technology, Cambridge, Massachusetts 02139

<sup>17</sup>University of Massachusetts, Amherst, Massachusetts 01003

<sup>18</sup>University of Mississippi, University, Mississippi 38677

<sup>19</sup>Nagoya University, Chikusa-ku, Nagoya 464, Japan

<sup>20</sup>University of Oregon, Eugene, Oregon 97403

<sup>21</sup>Istituto Nazionale di Fisica Nucleare Sezione di Padova and Università di Padova, I-35100 Padova, Italy

<sup>22</sup>Istituto Nazionale di Fisica Nucleare Sezione di Perugia and Università di Perugia, I-06100 Perugia, Italy

<sup>23</sup>Istituto Nazionale di Fisica Nucleare Sezione di Pisa and Università di Pisa, I-56100 Pisa, Italy

<sup>24</sup>Rutgers University, Piscataway, New Jersey 08855

<sup>25</sup>Rutherford Appleton Laboratory, Chilton, Didcot, Oxon OX11 0QX, United Kingdom

<sup>26</sup>Sogang University, Seoul, Korea

<sup>27</sup>Stanford Linear Accelerator Center, Stanford University, Stanford, California 94309

<sup>28</sup>University of Tennessee, Knoxville, Tennessee 37996

<sup>29</sup>Tohoku University, Sendai 980, Japan

<sup>30</sup>Vanderbilt University, Nashville, Tennessee 37235

<sup>31</sup>University of Washington, Seattle, Washington 98195

<sup>32</sup>University of Wisconsin, Madison, Wisconsin 53706

<sup>33</sup>Yale University, New Haven, Connecticut 06511

(Received 16 December 1994)

We present a comparison of the strong couplings of light ( $u$ ,  $d$ , and  $s$ ),  $c$ , and  $b$  quarks determined from multijet rates in flavor-tagged samples of hadronic  $Z^0$  decays recorded with the SLC Large Detector at the SLAC Linear Collider. Flavor separation on the basis of lifetime and decay multiplicity differences among hadrons containing light,  $c$ , and  $b$  quarks was made using the SLD precision tracking system. We find  $\alpha_s^{uds}/\alpha_s^{\text{all}}=0.987\pm 0.027(\text{stat})\pm 0.022(\text{syst})\pm 0.022(\text{theory})$ ,  $\alpha_s^c/\alpha_s^{\text{all}}=1.012\pm 0.104\pm 0.102\pm 0.096$ , and  $\alpha_s^b/\alpha_s^{\text{all}}=1.026\pm 0.041\pm 0.041\pm 0.030$ .

PACS number(s): 13.38.Dg, 11.30.Hv, 12.38.Qk

A fundamental assumption of the theory of strong interactions, quantum chromodynamics (QCD), is that the strong coupling  $\alpha_s$  is independent of quark flavor. This can be tested by measuring the strong coupling in events of the type  $e^+e^- \rightarrow q\bar{q}(g)$  for specific quark flavors  $q$ . Although an absolute determination of  $\alpha_s$  for each quark flavor would have large theoretical uncertainties [1], it is possible to test the flavor independence of QCD precisely by measuring ratios of couplings in which most experimental errors and theoretical uncertainties are expected to cancel. Since it has recently been suggested [2] that a flavor-dependent anomalous quark chromomagnetic moment could modify the probability for the radiation of gluons, comparison of the strong coupling for different quark flavors may also provide information on physics beyond the standard model.

Comparisons of  $\alpha_s$  for  $b$  or  $c$  quarks with  $\alpha_s$  for all flavors made at DESY PETRA [3] were limited in precision to  $\pm 0.41$  ( $c$ ) and  $\pm 0.57$  ( $b$ ) due to small data samples and limited heavy quark tagging capability. LEP measurements of  $\alpha_s^b/\alpha_s^{uds}$  have reached precisions between  $\pm 0.06$  and  $\pm 0.02$  [4]. However, these tests make the simplifying assumption

that  $\alpha_s$  is independent of flavor for all the non- $b$  quarks, and are insensitive to differences between  $\alpha_s$  for these flavors, especially a different  $\alpha_s$  for  $c$  quarks compared with either  $b$  or light quarks. The OPAL Collaboration has measured  $\alpha_s^f/\alpha_s^{\text{all}}$  for all five flavors  $f$  with no assumption on the relative value of  $\alpha_s$  for different flavors [5] to precisions of  $\pm 0.026$  for  $b$  and  $\pm 0.09$  to  $\pm 0.20$  for the other flavors. The kinematic signatures used to tag  $c$  and light quarks suffer from low efficiency and strong biases, due to preferential tagging of events without hard gluon radiation.

The SLC Large Detector (SLD) [6] at the SLAC Linear Collider (SLC) is an ideal environment in which to test the flavor independence of strong interactions. The tracking capability of the central drift chamber (CDC) [7] and the precision CCD vertex detector (VXD) [8], combined with the stable, micron-sized beam interaction point (IP), allows us to select  $Z^0 \rightarrow b\bar{b}(g)$  and  $Z^0 \rightarrow q_1\bar{q}_1(g)$  ( $q_1 = u, d, s$ ) events using their quark decay lifetime signatures with high efficiency and purity, and with low bias against three-jet events, an important advantage of this analysis. Here we present the first precise measurements of  $\alpha_s^b/\alpha_s^{\text{all}}$ ,  $\alpha_s^c/\alpha_s^{\text{all}}$ , and  $\alpha_s^{uds}/\alpha_s^{\text{all}}$  using this technique, and making no assumptions about the relative values of  $\alpha_s^b$ ,  $\alpha_s^c$ , and  $\alpha_s^{uds}$ .

This analysis is based on the  $1.8 \text{ pb}^{-1}$  of  $e^+e^-$  annihilation data collected during the 1993 run of the SLD at the

\*Deceased.

†Also at the Università di Genova.

‡Also at the Università di Perugia.




FIG. 1. The measured distribution of the number of tracks per event with  $d/\sigma_d \geq 3.0$  (points). The histograms show the flavor composition estimated from a Monte Carlo simulation (see text).

LC at a mean center-of-mass energy of  $\sqrt{s} = 91.26$  GeV. The trigger and selection criteria for hadronic  $Z^0$  decays are described in Ref. [1]. The efficiency for selecting a well-contained  $Z^0 \rightarrow q\bar{q}(g)$  event was estimated to be above 96% independent of quark flavor. The selected sample comprised 27 802 events, with an estimated  $0.10 \pm 0.05\%$  background dominated by  $Z^0 \rightarrow \tau^+ \tau^-$  events. This analysis used charged tracks measured in the CDC and in the VXD [1].

We used normalized impact parameters of  $d/\sigma_d$  as the basis for quark flavor tags, where  $d$  is the signed distance of closest approach of a charged track to the IP in the  $(x-y)$  plane transverse to the beam axis, and  $\sigma_d$  is the error on  $d$ . A resolution on  $d$  of  $10.8 \mu\text{m}$  has been measured using  $Z^0 \rightarrow \mu^+ \mu^-$  decays, and the spatial resolution on the average transverse IP position has been measured to be  $7 \mu\text{m}$  [9]. The distributions of  $d$  and  $d/\sigma_d$  are modeled well by the SLD simulation [9]. Tracks used for event flavor tagging were required to have at least one VXD hit; at least 40 CDC hits, with the first hit at a radius less than 39 cm; a combined CDC+VXD fit quality  $\sqrt{2\chi^2} - \sqrt{2N_{\text{DF}} - 1} < 8.0$ ; momentum greater than  $0.5 \text{ GeV}/c$ ;  $\sigma_d < 250 \mu\text{m}$ ; and to miss the IP by less than 0.3 cm in the  $x-y$  plane and by less than 1.5 cm in  $z$ . Tracks from candidate  $K^0$  and  $\Lambda$  decays and  $\gamma$  conversions were removed [9].

Figure 1 shows the distribution of  $n_{\text{sig}}$ , the number of tagging tracks per event with  $d/\sigma_d \geq 3$ . The data are well described by a Monte Carlo simulation of hadronic  $Z^0$  de-

cays [10] with parameter values tuned [11] to hadronic  $e^+e^-$  annihilation data, combined with a simulation of the SLD. For the simulation, the contributions of events of different quark flavors are shown separately. The leftmost bin contains predominantly events containing primary  $u$ ,  $d$ , or  $s$  quarks, while the rightmost bins contain a pure sample of events containing primary  $b$  quarks. The event sample was divided accordingly into three parts: those events with  $n_{\text{sig}} = 0$  were defined to be the  $uds$ -tagged sample; those with  $1 \leq n_{\text{sig}} \leq 3$  were the  $c$ -tagged sample; and those with  $n_{\text{sig}} \geq 4$  were the  $b$ -tagged sample. The hard  $b$  tag yields a sample with very low contamination from charm events, maximizing the sensitivity of the three-flavor test. The light-quark tag does not change the relative flavor composition of the  $uds$  sample. The efficiencies  $\varepsilon$  for selecting events (after cuts) of type  $i$  ( $i = uds, c, b$ ) with tag  $i$ , and the fractions  $\Pi$  of events of type  $i$  in the  $i$ -tagged sample, were calculated from the Monte Carlo simulation to be  $(\varepsilon, \Pi)_{uds} = (75.4 \pm 0.3\%, 86.8 \pm 1.2\%)$ ;  $(\varepsilon, \Pi)_c = (59.7 \pm 0.4\%, 30.0 \pm 0.6\%)$ ;  $(\varepsilon, \Pi)_b = (51.7 \pm 0.2\%, 93.6 \pm 0.6\%)$ ; the errors are discussed below.

Jets were then reconstructed using iterative clustering algorithms. We used the “E,” “E0,” “P,” and “P0” variations of the JADE algorithm, as well as the “Durham” (“D”) and “Geneva” (“G”) algorithms [12]. We divided events into two categories: those containing (1) two jets and (2) three or more jets. The fraction of the event sample in category 2 was defined as the three-jet rate  $R_3$ . This quantity is infrared and collinear safe and has been calculated to  $O(\alpha_s^2)$  in perturbative QCD [12,13]. For each algorithm, the jet resolution parameter  $y_c$  was chosen to be as small as possible subject to the requirement that  $O(\alpha_s^2)$  QCD provides a good description of  $R_3$  measured in our global sample of all flavors [1,14]. This choice maximizes  $R_3$  while avoiding the “Sudakov region” at low  $y_c$  where multiple gluon emission requires that large logarithmic terms of  $1/y_c$  be resummed in order to describe the data [1]. The resulting  $y_c$  values are listed in Table I.

The  $R_3^j$  for each of the  $j$  quark types ( $j = uds, c, b$ ) was extracted from a maximum likelihood fit to  $n_2^i$  and  $n_3^i$ , the number of two-jet and three-jet events, respectively, in the  $i$ -tagged sample:

$$n_2^i = \sum_{j=1}^3 [\varepsilon_{(2 \rightarrow 2)}^{ij} (1 - R_3^j) + \varepsilon_{(3 \rightarrow 2)}^{ij} R_3^j] f^j N, \quad (1)$$

$$n_3^i = \sum_{j=1}^3 [\varepsilon_{(3 \rightarrow 3)}^{ij} R_3^j + \varepsilon_{(2 \rightarrow 3)}^{ij} (1 - R_3^j)] f^j N.$$

TABLE I. Results for  $R_3^j/R_3^{\text{all}}$ , derived from Eq. (1); see text. Errors shown are statistical.

Algorithm	$y_c$	$R_3^{uds}/R_3^{\text{all}}$	$R_3^c/R_3^{\text{all}}$	$R_3^b/R_3^{\text{all}}$	$R_3^c/R_3^u$ factor	$R_3^b/R_3^d$ factor
E	0.080	$0.941 \pm 0.042$	$1.212 \pm 0.173$	$0.980 \pm 0.062$	0.995	0.958
E0	0.050	$0.975 \pm 0.026$	$1.113 \pm 0.145$	$0.981 \pm 0.053$	0.994	0.945
P	0.030	$1.001 \pm 0.027$	$0.985 \pm 0.109$	$1.007 \pm 0.041$	0.992	0.929
P0	0.030	$1.014 \pm 0.026$	$0.899 \pm 0.102$	$1.037 \pm 0.039$	0.992	0.929
D	0.015	$0.989 \pm 0.035$	$1.096 \pm 0.145$	$0.947 \pm 0.049$	0.991	0.921
G	0.030	$1.032 \pm 0.020$	$0.942 \pm 0.079$	$0.952 \pm 0.030$	0.989	0.915




FIG. 2. Values of  $\alpha_s^j/\alpha_s^{\text{all}}$  derived for each of the six jet algorithms for each of the quark flavors  $j$  (see text). The error bars on the averages include the statistical and systematic errors and the total theoretical uncertainty.

Here  $N$  is the total number of selected events corrected for the event selection efficiency, and  $f^j$  is the standard model fractional hadronic width for  $Z^0$  decays to quark type  $j$ . The matrices  $\varepsilon_{(2\rightarrow 2)}^{ij}$  and  $\varepsilon_{(3\rightarrow 3)}^{ij}$  are the efficiencies for an event of type  $j$ , with two- or three-jets at the parton level, to pass all cuts and be tagged as a two- or three-jet event, respectively, of type  $i$ . Matrices  $\varepsilon_{(2\rightarrow 3)}^{ij}$  and  $\varepsilon_{(3\rightarrow 2)}^{ij}$  are the efficiencies for an event of type  $j$ , with two- or three-jets at the parton level, to pass all cuts and be tagged as a three- or two-jet event, respectively, of type  $i$ . This formalism explicitly accounts for modifications of the parton-level three-jet rate due to hadronization, detector effects, and tagging bias. These matrices were calculated from the Monte Carlo simulation. The efficiencies for correctly tagging a two-jet event and a three-jet event differ by an average of 5.7%, 8.3%, and 30.3% for the  $uds$ ,  $c$ , and  $b$  tags, respectively.

Equations (1) were solved using two- and three-jet events defined by each of the six algorithms. The ratios  $R_3^j/R_3^{\text{all}}$ , where  $R_3^{\text{all}}$  is the three-jet rate in the total event sample, are shown in Table I. Averaged over all six algorithms the correlation coefficients from the fit are  $uds$ - $c$ :  $-0.76$ ,  $uds$ - $b$ :  $0.30$ ,  $c$ - $b$ :  $-0.55$ . The statistical errors were calculated using the full covariance matrix.

The three-jet rate in heavy quark ( $b, c$ ) events is expected to be reduced relative to that in light quark events by the diminished phase-space for gluon emission due to the quark masses. We evaluated the suppression factors,  $R_3^c/R_3^u$  and  $R_3^b/R_3^d$ , for each jet algorithm and  $y_c$  value according to Ref. [15], assuming  $b$  ( $c$ ) quark masses of  $4.75 \text{ GeV}/c^2$  ( $1.50 \text{ GeV}/c^2$ ). These factors are listed in Table I, and were used to correct the measured three-jet rate ratios.

To  $O(\alpha_s^2)$  in perturbative QCD,  $R_3(y_c) = A(y_c)\alpha_s + [B(y_c) + C(y_c)]\alpha_s^2$ , where the  $O(\alpha_s^2)$  coefficient includes a term  $B(y_c)$  from three-parton states calculated at

TABLE II. Contributions to the systematic error on  $\alpha_s^j/\alpha_s^{\text{all}}$ .

Source	$\Delta\left(\frac{\alpha_s^{uds}}{\alpha_s^{\text{all}}}\right)$	$\Delta\left(\frac{\alpha_s^c}{\alpha_s^{\text{all}}}\right)$	$\Delta\left(\frac{\alpha_s^b}{\alpha_s^{\text{all}}}\right)$
$b$ physics	0.008	0.060	0.033
$c$ physics	0.017	0.060	0.011
Detector modeling	0.003	0.032	0.017
Monte Carlo statistics	0.011	0.048	0.014
QCD uncertainty	0.003	0.011	0.012

next-to-leading order, and a term  $C(y_c)$  from four-parton states calculated at leading order. Hence, the ratio of the strong coupling of quark type  $j$  to the mean coupling in the sample of all flavors,  $\alpha_s^j/\alpha_s^{\text{all}}$ , can be determined from

$$\frac{R_3^j(y_c)}{R_3^{\text{all}}(y_c)} = \frac{A(y_c)\alpha_s^j + [B(y_c) + C(y_c)](\alpha_s^j)^2}{A(y_c)\alpha_s^{\text{all}} + [B(y_c) + C(y_c)](\alpha_s^{\text{all}})^2}, \quad (2)$$

where  $A(y_c)$ ,  $B(y_c)$ , and  $C(y_c)$  for the different jet-finding algorithms were evaluated using Refs. [12,13]. Using our measured values of  $\alpha_s^{\text{all}}(M_Z^2)$  determined from jet rates [14] we found that for the  $E$ ,  $EO$ ,  $P$ ,  $P0$ , and  $D$  algorithms, the leading-order QCD calculation  $C(y_c)\alpha_s^2$  lies below the experimental four-jet rate by roughly a factor of 2. We increased  $C(y_c)$  *ad hoc* for these algorithms, so as to describe the data. Equation (2) was solved to obtain  $\alpha_s^j/\alpha_s^{\text{all}}$  for each jet algorithm; the results are shown in Fig. 2. The errors include contributions from the statistical error, as well as the experimental systematic errors and theoretical uncertainties.

We considered systematic effects that could modify the tagging efficiencies. In each case the error was evaluated by varying the appropriate parameter in the Monte Carlo simulation, recalculating the matrices  $\varepsilon$ , performing a new fit to Eq. (1) and rederiving  $\alpha_s^j/\alpha_s^{\text{all}}$ . Suitable variation about the world average value of each parameter was considered [9]. The errors are summarized in Table II, where averages over the six algorithms are shown. The largest contributions result from limited knowledge of the heavy quark fragmentation functions and  $B$  decay multiplicity. The uncertainty in  $\mathcal{B}(Z^0 \rightarrow c\bar{c})$  also produces large variations in  $\alpha_s^c/\alpha_s^{\text{all}}$  and  $\alpha_s^{uds}/\alpha_s^{\text{all}}$ . Contributions from  $b$  hadron lifetimes, the fraction of  $D^+$  in  $B$  meson decays,  $b$  baryon production rates, and the charm hadron decay multiplicity are small. The detector systematic error is dominated by the uncertainty in the charged track reconstruction efficiency. No systematic variation of the results was found when the event selection cuts, tag criteria, or  $y_c$  values were changed.

We considered sources of uncertainty in the QCD predictions that affect the values of  $\alpha_s^j/\alpha_s^{\text{all}}$  derived from Eq. (2). For each jet algorithm these include variation of the QCD renormalization scale within the range allowed by our measurements of jet rates in the global sample [14] and variation of the heavy quark masses used in the phase-space correction factors by  $\pm 0.25 \text{ GeV}/c^2$ . In addition, the shifts in  $\alpha_s^j/\alpha_s^{\text{all}}$  due to the *ad hoc* increase of the coefficient  $C(y_c)$  were conservatively assigned as an uncertainty. The variation of the results due to uncertainties in parton production and hadronization was investigated [16] by using the JETSET [10] and

HERWIG [17] event generators and was found to be small. These contributions were added in quadrature to yield the total QCD uncertainties listed in Table II.

There is significant scatter among the  $\alpha_s^j/\alpha_s^{\text{all}}$  values derived from the different jet algorithms. In order to quote a single  $\alpha_s^j/\alpha_s^{\text{all}}$  value for each flavor  $j$ , we made the conservative assumption that the results are completely correlated, and we calculated the unweighted mean values and errors over all six algorithms. We obtained

$$\begin{aligned}\frac{\alpha_s^{uds}}{\alpha_s^{\text{all}}} &= 0.987 \pm 0.027(\text{stat}) \pm 0.022(\text{syst}) \pm 0.022(\text{theory}), \\ \frac{\alpha_s^c}{\alpha_s^{\text{all}}} &= 1.012 \pm 0.104(\text{stat}) \pm 0.102(\text{syst}) \pm 0.096(\text{theory}), \\ \frac{\alpha_s^b}{\alpha_s^{\text{all}}} &= 1.026 \pm 0.041(\text{stat}) \pm 0.041(\text{syst}) \pm 0.030(\text{theory}),\end{aligned}\tag{3}$$

where the theoretical uncertainty is the sum in quadrature of the QCD uncertainty from Table II and the rms of the results

over the six algorithms. These averages are also shown in Fig. 2. The variation of results among jet algorithms, presumably due to different uncalculated  $O(\alpha_s^3)$  QCD contributions, dominates the theoretical uncertainty, is not small compared with experimental errors, and has not been considered in previous analyses [4,5].

In conclusion, we have used hadron lifetime information to separate hadronic  $Z^0$  decays into three flavor samples with high efficiency and purity, and small bias against events containing hard gluon radiation. From a comparison of the rates of multijet events in these samples, we find that the strong coupling is independent of quark flavor within our sensitivity. These are the first such results using a precision vertex detector for flavor separation at the  $Z^0$ . This represents the most precise test for  $uds$  events. Our findings are consistent with measurements performed at the CERN  $e^+e^-$  collider LEP using different flavor-tagging techniques [4,5].

We thank the personnel of the SLAC accelerator department and the technical staffs of our collaborating institutions for their outstanding efforts. We thank C. Ng, T. Rizzo, and E. Maina for their helpful contributions.

- 
- [1] SLD Collaboration, K. Abe *et al.*, Phys. Rev. D **51**, 962 (1995).  
 [2] T. G. Rizzo, Phys. Rev. D **50**, 4478 (1994).  
 [3] TASSO Collaboration, W. Braunschweig *et al.*, Z. Phys. C **42**, 17 (1989); W. Braunschweig *et al.*, *ibid.* **44**, 365 (1989).  
 [4] DELPHI Collaboration, P. Abreu *et al.*, Phys. Lett. B **307**, 221 (1993); L3 Collaboration, B. Adeva *et al.*, *ibid.* **271**, 461 (1991); OPAL Collaboration, M. Z. Akrawy *et al.*, Z. Phys. C **65**, 31 (1995).  
 [5] OPAL Collaboration, R. Akers *et al.*, Z. Phys. C **60**, 397 (1993).  
 [6] SLD Design Report, SLAC Report No. 273, 1984 (unpublished).  
 [7] M. D. Hildreth *et al.*, in *1994 IEEE Conference Record*, Proceedings, Norfolk, Virginia, 1994, edited by R. C. Tredler (IEEE, New York, 1994).  
 [8] SLD Collaboration, M. G. Strauss *et al.*, in *Proceedings of the 27th International Conference on High Energy Physics*, Glasgow, Scotland, 1994, edited by P. J. Bussey and I. G. Knowles (IOP, London, 1995), Vol. 2, p. 1179.  
 [9] SLD Collaboration, K. Abe *et al.*, Phys. Rev. D **53**, 1023 (1996).  
 [10] T. Sjöstrand, Comput. Phys. Commun. **43**, 367 (1987).  
 [11] P. N. Burrows, Z. Phys. C **41**, 375 (1988); OPAL Collaboration, M. Z. Akrawy *et al.*, *ibid.* **47**, 505 (1990).  
 [12] S. Bethke *et al.*, Nucl. Phys. **B370**, 310 (1992).  
 [13] Z. Kunszt *et al.*, in *Z Physics at LEP1*, Proceedings of the Workshop, Geneva, Switzerland, 1989, edited by G. Altarelli *et al.* (CERN, Report No. 89-08, Geneva, 1989), Vol. 1, p. 373.  
 [14] SLD Collaboration, K. Abe *et al.*, Phys. Rev. Lett. **71**, 2528 (1993).  
 [15] A. Ballestrero, E. Maina, and S. Moretti, Phys. Lett. B **294**, 425 (1992).  
 [16] SLD Collaboration, K. Abe *et al.*, Phys. Rev. D **51**, 962 (1995).  
 [17] G. Marchesini *et al.*, Comput. Phys. Commun. **67**, 465 (1992).