

**Fungicidal effect of three new synthetic cationic peptides
against *Candida albicans***

H NIKAWA¹, DDS, PhD., H FUKUSHIMA², DDS, S MAKIHIRA³, DDS, PhD., T, HAMADA⁴,
DDS, PhD and L P SAMARANAYAKE⁵, BDS, DDS, FRCPath

Department of Prosthetic Dentistry, Hiroshima University Faculty of Dentistry, Hiroshima,
Japan.

¹Lecturer, Department of Prosthetic Dentistry.

²Graduate student, Department of Prosthetic Dentistry.

³Instructor, Department of Prosthetic Dentistry.

⁴Professor and Chairman, Department of Prosthetic Dentistry.

⁵Professor, Oral Biosciences, Faculty of Dentistry, University of Hong Kong.

Reprint requests to: Dr. Hiroki Nikawa

Department of Prosthetic Dentistry,

Hiroshima University Faculty of Dentistry,

1-2-3 Kasumi Minami-ku

Hiroshima, 734-8553, Japan.

Fax: +81-82-257-5684,

E-mail: hirocky@hiroshima-u.ac.jp

This study is supported by Grant-in-Aid 13470416, 14370636

Abstract

OBJECTIVE: Peptide antibiotics are considered a new class of antifungal agents. Of these, an alpha-helical, cationic peptide termed Dhvar 4, a relative of salivary histatin has been shown to be an antifungal of relatively high potency. Similarly, lactoferricin B (LFB) and a derivative thereof, LFB(17-30), disrupts the fungal cell membrane and acts against *C. albicans*. As Dhvar 4 and LFB(17-30), exhibit almost identical amino acid sequences at their C-terminal, we hypothesized that laboratory synthesis of peptides with an alpha-helical structure and having similar amphipathic properties could lead to products with candidacidal activity. Hence, three such peptides - JH8194, JH8195 and JH 8944, were synthesized and their antifungal properties compared with recognized antifungals LFB, LFB (17-30), human lactoferricin (LFH), Histatin-5 and Dhvar 4, against two isolates of *C. albicans*

MATERIALS AND METHODS: The antifungal agents were synthesized and their secondary structures evaluated according to a previously described protocol of Situ et al. (2000). *C. albicans* strains were oral isolates from a human immunodeficiency virus (HIV)-infected (isolate A2) and a healthy (A6) individual. A standard concentration of yeasts was exposed to a range of dilutions of the agents for a specific duration and the cell death (viability) in terms of the resultant colony forming units (CFU) /mL was quantified.

RESULTS: Dhvar 4, showed the most alpha-helical propensity, and was the least fungicidal while LFB and LFB(17-30) showed the highest antifungal potential, and demonstrated total kill of A6, and A2 at 5 μ and 10 μ M concentrations, respectively whilst LFH killed both isolates at a 10 μ M concentration. Of the three new synthetic peptides, JH 8194 was the most potent (total

kill of A6/A2 strains at 1.25/ 2.5 μ M), followed by JH 8195 (total kill of A6/A2 strains at 5/ 10 μ M while JH 8944 was the least potent as a 25 μ M concentration was required to kill either strain of *Candida*. On further analyses of the relationship between pI value of the peptides and their anticandidicidal activity, a significant positive correlation was noted. In order to rule out a cytotoxic effect of the new synthetic peptides we compared the fungicidal and hemolytic activities under similar incubation conditions using freshly isolated erythrocytes and all three peptides exhibited no detectable hemolysis upto a concentration of 100 μ M in contrast to the polyene antifungal amphotericin B that elicited significant initiation of hemolysis at a concentration of 5.0 μ M.

CONCLUSION: Our data suggest that laboratory synthesis of agents with an alpha-helical structure and having amphipathic properties similar to known, natural antifungal agents may be a promising avenue to generate products with improved antifungal activity.

Key Words: cationic peptide, antifungal activity, *Candida albicans*, Abbreviations: LF, lactoferricin; LFB, bovine lactoferricin; LFH, human lactoferricin

Introduction

Candida albicans is an opportunistic fungal pathogen that causes mucosal and systemic infections in humans. The AIDS epidemic, improved life-sustaining therapies, and aggressive anticancer therapy have all contributed to the rise in the number of severely immunocompromised patients in the community leading in turn to an increased fungal infections due to this opportunistic yeast (Samaranayake et al 2002). The concomitant escalation in the use of antifungals for prophylaxis and management of these diseases has led to the emergence of drug resistant yeasts worldwide (White et al. 1998). In addition, a number of currently available antifungal agents, such as polyene and azole antimycotics and DNA analogues, have undesirable toxic and other side effects (Situ et al. 2000). Therefore, the search for more effective agents analogous to natural antifungal substances with lesser toxicity is of crucial importance.

A number of natural antifungal agents are found in human mucosal secretions including saliva. These include, secretory IgA, lactoferrin (Nikawa et al. 1993, 1994, 1995), lysozyme (Tobgi et al. 1987, Nikawa et al. 1993), and histatins (Pollock et al. 1984) that are demonstrably candidacidal in nature. Histatins in particular are a group of small, cationic antifungal peptides present both in human saliva (Oppenheim et al. 1988, Pollock et al. 1984) and serum (Murakami et al. 1994) and divided into three major sub-groups, histatin-1, -3 and -5. Of these the latter shows the most potent candidacidal activity (Sternberg 1994), and is thought to play a critical role in primary protection of the oral cavity against candidal infections, especially in compromised hosts (Samaranayake, 2002). In fact, of the aforementioned proteins only the salivary histatins levels show a statistically significant decrease in AIDS patients who develop candidiasis (Tsai et al. 1997). This perhaps is a reason why more than 90 % of AIDS patients develop oral candidiasis over the course of the disease (Samaranayake et al , 1989). We too have recently shown that oral

C. albicans isolates from HIV-infected individuals are less sensitive to Histatin-5 compared with those from healthy controls (Nikawa et al, 2002). For these reasons, there has been intense interest in Histatin-5 and its variant synthetic peptides in order to develop compounds with more efficacious fungicidal properties. Of the synthetic variants of histatins, an alpha-helical peptide, Dhvar 4 has been shown to possess intense fungicidal/static activity (Helmerhorst et al. 1999). Furthermore, other compounds such as lactoferrin B (LFB) and LFB (17-30), derived from bovine lactoferrin, exhibits natural antifungal activity against *C. albicans* (Bellamy et al., 1993, Wakabayashi et al 1998) and so is another peptide derived from the N-terminal region of lactoferrin, lactoferricin (LF) (Bellamy et al., 1993, Jones 1994, Lupetti et al. 2000). These peptide antifungals are considered a new class of promising agents that interact with the yeast cell membranes leading to the disruption of their cellular integrity and cell death (Helmerhorst et al. 1999).

On comparison of the amino acid sequence of the C-terminals of Dhvar 4 and LFB (17-30) we noted a very high congruence (Table 1) and hypothesized that synthetic peptide fragments with identical amino acid sequences and amphipathic properties may yield compounds with improved fungicidal properties. Hence, we synthesized *de novo*, three different peptides - JH8194, JH8195 and JH 8944 with the latter properties, using standard biochemical techniques. Afterwards, the alpha-helicity and the fungicidal properties of the cationic peptides were characterized and their anticandidal activity compared with LFB, LFB(17-30), human lactoferricin (LFH), Histatin-5 and Dhvar 4, against two oral isolates of *C. albicans* derived from a healthy and a HIV-infected individual.

Materials and Methods

Candida isolates and growth conditions

In total two isolates of *C. albicans* were used in the experiment. *C. albicans* A2 was an oral isolate from a male, ethnic Chinese, HIV-infected individual while *C. albicans* A6 was an oral isolate from a healthy female. Both strains were isolated using an oral rinse technique as described previously (Samaranayake et al 1986). The identity of both isolates was reconfirmed with standard sugar assimilation tests using the API 20C system (API Products, Biomerieux, Lyon, France) and the "germ tube" test (Silverman et al. 1990). In a previous study of ours these isolates exhibited differential *in vitro* sensitivity to histatin-5, the isolate A2 being more resistant to histatin than the isolate A6 (Nikawa et al. 2002).

In order to obtain yeast suspensions for the sensitivity assays a loopful of the yeast was inoculated into 100 mL of yeast nitrogen base medium (Difco, Detroit, USA) containing 250 mM glucose and grown aerobically at 37 °C for 24 h. After 18 h incubation, the yeasts were harvested in the mid-exponential growth phase, washed twice with PBS containing 1 mM phosphate buffer (pH 6.8) and resuspended to a final concentration of 10⁷ yeasts/mL by haemocytometric counting (Nikawa et al, 1993, 1996). All yeasts were in the blastospore phase at this stage.

Cationic peptides, peptide design and synthesis

Histatin-5 (DSHEKRHHGYKRKFHEKHHSHRGY; IUPAC-IUB, pI=10.70), bovine lactoferricin (LFB; FKRRWQWRMKKLGAPSITCVRRAF, pI=12.34), LFB(17-30) (LFB(17-30); FKRRWQWRMKKLG, pI=12.23), human lactoferricin (LFH; TKCFWQRNMRKVRGPPVSCIQR, pI= 12.07), Dhvar 4 (KRLFKLLFSLRKY,

pI=11.69), and three new peptides, i.e. JH8194 (KRLFRRWQWRMKKY, pI=12.51), JH8195 (KRLFRLLFSMKKY, pI=12.24), and JH8944 (FKCKKVVISLRRY, pI=10.88) were all synthesized using the T-bag method adapted for 9-fluorenyl methoxycarbonyl (Fmoc)-chemistry. (Helmerhorst et al. 1999).

p-Benzoyloxybenzyl alcohol resins, with the first N-Fmoc-protected amino acids already attached, were included inside the T-bags. The coupling reactions were performed in N, N-dimethylformamide. After completion of the sequence, cleavage from the resin and simultaneous side-chain deprotection was achieved with a mixture of 5% thioanisole, 5% phenol, 5% water, and 85 % trifluoroacetic acid (TANA laboratories., Texas, USA). Purification and purity analysis of peptides were performed using High performance liquid chromatography (HPLC) and reverse phase HPLC. Further confirmation of molecular weight was performed using Mass spectrometry (Matrix Assisted Laser Desorption Ionization-TOF/MS).

Circular dichroism (CD) spectra analysis

CD spectra of the peptides were acquired on a JASCO J-820 spectropolarimeter, calibrated with D10 camphor sulfonic acid. Peptides, other than Histatin-5, LFB and LFH, dissolved in 100 trifluoroethanol (TFE), 75 TFE/distilled water (DS), 50 TFE/DS, 25 TFE/DS and 100 DS, were scanned in the range of 247 to 185 nm for secondary structural analysis. The peptide concentration used was 100 mg/mL. Four scans were acquired from each sample and averaged to improve signal quality. All spectra were expressed in terms of Mean residue ellipticity (MRE), in units of deg cm² /dmol. (Situ et al 2000).

Secondary structure analysis

Far UV CD spectra were analyzed by convex constraint analysis (CCA) according to the method of Situ et al (2000). The ellipticity values at each wavelength in the range of 247 to 185 nm were used for the calculation. The input data set consisted a matrix of 30 proteins, and the spectra were deconvolved in 100 iterations. The spectral assignments of each peptide was performed as described by Perczel and Hollosi (1992) .

Candidacidal assay

Antifungal activity of the peptides was evaluated according to the method of Edgerton et al. (1998) with some modifications. The assays were performed on exponential phase *C. albicans* cells in the presence or absence of increasing concentrations of the peptides (0.325-100 μ M). Briefly, *C. albicans* cells were washed twice with 1 mM sodium phosphate buffer ($\text{Na}_2\text{HPO}_4/\text{NaH}_2\text{PO}_4$; pH 6.8), and resuspended at 1.8×10^5 cells/ml. Twenty microliter of the cell suspension was mixed with 20 ml of 1 mM phosphate buffer containing the test peptide and incubated for 90 min at 37 °C with shaking. Controls were incubated with 20 ml of 1mM phosphate buffer alone. The reaction was stopped by addition of 360 mL of yeast nitrogen base (YNB); 40 uL (microliter) of the suspension were spread onto Sabouraud dextrose agar plates and incubated for 48 h at 37 °C and the resultant colony forming units (CFUs) in the test and control inocula quantified by manual counting. The assays were performed on two independent occasions with quadruplicated specimens on each occasion. Then the percent fungicidal activity of each peptide was calculated as follows.

[Colony Forming Units CFU/ mL from the test suspension with the peptide/ CFU/ mL from the control suspension without the peptide] x 100 .

Assessment of pI value of peptides

The pI (isoelectric point) value of each synthetic peptide was calculated by the use of DNASIS, a DNA & protein analysis software (Hitachi Software Engineering Co., Ltd., Tokyo Japan.) The killing effects of each peptide at 1.25-5.0 μ M were plotted against pI values, and the correlation coefficient evaluated using regression analysis.

Hemolytic assay

Human erythrocytes from a healthy donor were collected in vacuum tubes containing heparin (final concentration 20.4 U/ml) as anti-coagulant. The erythrocytes were prepared for the hemolytic assay as described by Helmerhorst et al. (1999). Briefly, the erythrocytes were harvested by centrifugation for 10 min at 2000 x g at room temperature, and washed three times in PBS (9mM sodium phosphate, pH 7.0 in 150 mM NaCl). From this pellet a 20% (vol/vol) of erythrocytes/PBS suspension was prepared and used directly for the hemolytic assay. A total of 100 μ L of the latter suspension was added in duplicate to 100 μ L of a two-fold serial dilution series of each peptide in the same buffer in a 96-well V-bottomed microtiter plate. Total hemolysis (control) was achieved with Tween-20.

The plates were incubated for 1h at 37°C and then centrifuged for 5 min at 1500 x g at 20°C. Of the supernatant fluid 150 μ l was transferred to a flat-bottom microtiter plate, and the absorbance measured at 450 nm. The percentage hemolysis was calculated as: [(A450 of the peptide treated sample - A450 of buffer treated sample) / (A450 of Tween-20 treated sample - A450 of buffer treated sample)] x 100. To determine the hemolysis by amphotericin B, the same assay was employed with the peptide replaced by amphotericin B.

Statistical analysis

All of the numerical data obtained were analyzed by an one-way analysis of variance (ANOVA) and subjected to Tukey's multiple range test ($p < 0.05$).

Results and Discussion

Purity of synthetic peptides

Purity analysis of peptides using HPLC and Mass spectrometry revealed one major peak for each peptide, and the purity of each peptide was as follows; Histatin-5 : 96.00%, LFB : 100.00%, LFB(17-30) : 98.47%, LFH : 100.00%, Dhvar 4 : 99.30%, JH8194 : 99.55%, JH8195 : 100.00%, and JH8944 : 97.35% .

Conformation analysis of peptides

Secondary structure of antimicrobial peptides, particularly the alpha-helical conformation either in non-aqueous or a hydrophobic environment, is reported to play an important modulating role of their antimicrobial action, particularly the permeabilization of the fungal cell membrane. Thus we analyzed the alpha-helical propensity of each peptide in serially diluted trifluoroethanol (TFE), considered an alpha-helix-promoting solvent. The quantitative shape analysis was performed as described by Perczel and Hollosi (1992) whereas quantitative estimation of the secondary structures was determined by convex constraint analysis (CCA). The latter analysis has been previously employed to estimate the secondary structural content of oligopeptides (Perczel and Hollosi 1992). Since beta-turn is known to involve a specific pair of amino acids, and short oligopeptides usually transit between two phases, we eliminated beta-structures, such

as turns or sheets, in the present study, after which the helix/random coil ratio was calculated (Table 2).

As shown in Fig. 1-a, Dhvar 4 showed very high alpha-helical propensity. In distilled water (DS), CD spectra of Dhvar 4 exhibited a random coil, and in 25% TFE/DS solution, Dhvar 4 comprised 100% alpha-helical conformation and 0% random coil (Fig 1-a: Table 2). alpha-helical conformation of Dhvar 4 matured as the TFE concentration increased from 0 to 25% (Table 2).

In contrast, CD spectra of LFB(17-30) essentially exhibited the random coil conformation, and even in 100% TFE, LFB (17-30) exhibited only 39.0% α -helical conformation and 61.0% random coil (Fig 1-b; Table 2). CD spectra of JH8194 and JH 8195 showed the concentration-dependent conformational changes in TFE. JH8194 formed random coil in distilled water, and incremental increase in their α -helicity with increasing TFE concentration (α -helicity; 2.2% in 25% TFE, 14.5% in 50% TFE, and 34.0% in 75% TFE) and, total alpha-helical conformation in 100% TFE (Fig. 1-c). A similar relationship between alpha-helical conformation and the concentration of TFE was noted with JH8195 (Fig 1-d; Table 2). In contrast, CD spectra of JH 8944 exhibited random coil conformation in 100% TFE, but this compound exhibited 22.7% alpha-helical conformation and 77.3% random coil in 100% TFE (Fig 1-e; Table 2).

Fungicidal activity of peptides

Pilot studies indicated that the fungicidal effect of each peptide gradually increased up to 60 min incubation, and plateaued thereafter (data on file). Hence we examined the fungicidal effect of all the peptides after a standard 90-min period of incubation in all experiments described below.

With regard to Histatin 5 exposure the loss of viability of the two *C. albicans* isolates was dose dependent. Whilst *C. albicans* A6 was more sensitive to Histatin-5, both isolates were rendered totally non-viable at a Histatin-5 concentration of 100 μM . These observations concur with our previous studies where 50 μM Histatin-5, killed almost all (> 95%) of *C. albicans* isolates from healthy individuals but only 66.1 to 75.3 % of two *C. albicans* isolates from HIV-infected individuals. The family of salivary histatins consists of structurally related, low molecular weight histidine rich proteins that contribute to the non-immune defense system of the orol-pharyngeal milieu and, are found in saliva of healthy adults at concentrations ranging from 50 - 425 $\mu\text{g/ml}$ (Edgerton et al. 1998). Histatin-5 is the most potent member of the family and renders most pathogenic *Candida* species non-viable *in vitro* at physiological concentrations (Raj et al 1990).

On the contrary, Dhvar 4, a relative of histatin, which showed the most alpha-helical propensity, was the least effective in killing the two isolates (Figs. 2 & 3). In contrast, LFB and LFB(17-30) exhibited higher potency leading to total kill of A6 and A2 isolates at 5 μM and 10 μM concentrations, respectively (Figs. 2 and 3). The killing effect of LFH was relatively high, and 10 μM of LFH rendered both isolates totally non-viable (Figs. 2 & 3). Notably though, the fungicidal effect of lower concentrations of LFH was less than that of LFB and LFB(17-30).

As compared with the foregoing natural peptides the JH8194, one of the three new synthetic peptides, was exquisitely candidacidal and rendered all A6 and A2 isolates non-viable at 1.25 μM at 2.5 μM concentrations. JH 8195 was of intermediate potency and killed all A2 and A6 isolates at 5 μM and 10 μM concentrations. However, JH 8944 was the least effective fungicidal of the three synthetic peptides, and required 25.0 μM concentrations for a total kill both isolates.

It is generally accepted that many antimicrobial peptides have an alpha-helical structure, and the majority are cationic and amphipathic. Our results suggest that alpha-helicity contributes to the killing effects of peptides, but only to a limited extent. In most microbes, including fungi, both the outer leaflet of the plasma membrane as well as the outer membrane contain anionic molecules oriented towards the exterior of the cell (Shai 1999). As this is not the case for mammalian cell membranes, the cationic antimicrobial peptides preferentially bind to the exposed negative charges of microbial membranes, instead of the zwitterionic amphiphiles present in the exterior surface of the mammalian cells.

We further analysed the relationship between pI value of the peptides and their anticandidal activity. A significant positive correlation between these elements was also observed (Fig. 4). These results imply that higher the pI value of the peptide, the greater is its antifungal activity.

Hemolysis assay

The hemolytic activity of cationic peptides in phosphate buffer is a surrogate marker of their antimicrobial activity, in various growth media (Dathe et al FEBS letts 1997;403; 208-212). Yet the two assays measure different properties, as hemolysis indicates membrane perturbing properties and the latter assay, the growth inhibitory activity. We compared the fungicidal and hemolytic activities under similar incubation conditions using freshly isolated erythrocytes as described by Helmerhorst et al (1999). As shown in Fig. 5, all peptides used in the present study, exhibited no detectable hemolysis upto a concentration of 100 μ M in contrast to the polyene antifungal amphotericin B that elicited significant hemolysis at much lower concentrations (5.0 μ M). As none of the three synthetic peptides were more hemolytic than the widely used polyene

antifungal amphotericin B , these histatin analogues could be considered as safe candidates for putative antifungal agents.

In summary, we have demonstrated that three synthetic peptides, that are histatin analogues have powerful anticandidal activity that may be superior to the naturally occurring antifungals such as histatin 5, lactoferricin B and its derivative LFB(17-30), and the synthetic peptides Dhvar 4. The pI values and the hemolytic properties of these synthetic peptides also indicate that they may be further developed as safe and efficacious agents for treating candidal infections.

Legends to Figures

Fig. 1 Conformational analysis of cationic peptides, i.e. Dhvar 4 (a), LFB (17-30) (b), JH8194 (c), JH8195 (d) and JH8944 (e) by circular dichroism (CD). All spectra are expressed in terms of mean residue ellipticity. The peptide concentration was used was 100 μ M.

Fig. 2 Killing effects of cationic peptides against *C. albicans* A6, an isolate from HIV-negative patient. The data indicate mean viability of survived fungi.

Fig. 3 Cationic peptides induced loss of viability of *C. albicans* A2, an isolate from HIV-positive patient. The data indicate mean viability of survived fungi.

Fig. 4 Relationship between pI values of cationic peptides and their fungicidal effects.

Fig. 5 Hemolytic activity of cationic peptides and amphotericin B.

References

- Bellamy W, Wakabayashi H, Takase M, Kawase K, Shimamura S, Tomita M (1993). Killing of *Candida albicans* by lactoferricin B, a potent antimicrobial peptide derived from the N-terminal region of bovine lactoferrin. *Med Microbiol Immunol* 182:97-105.
- Dathe M, Wieprecht T, Nikolenko H, Handel L, Maloy WL, MacDonald DL, Beyermann M, Bienert M. (1997) Hydrophobicity, hydrophobic moment and angle subtended by charged residues modulate antibacterial and haemolytic activity of amphipathic helical peptides. *FEBS Lett.* 403:208-12.
- Edgerton M, Koshlukova SE, Lo TE, Chrzan BG, Straubinger RM, Raj PA (1998). Candidacidal activity of salivary histatins. Identification of a histatin 5-binding protein on *Candida albicans*. *J Biol Chem* 273:20438-47.
- Groenink J, Walgreen-Weterings E, van 't Hof W, Veerman EC, Nieuw Amerongen AV. (1999) Cationic amphipathic peptides, derived from bovine and human lactoferrins, with antimicrobial activity against oral pathogens. *FEMS Microbiol Lett.* 15;179:217-22.
- Helmerhorst EJ, Reijnders IM, van 't Hof W, Veerman EC, Nieuw Amerongen AV. (1999) A critical comparison of the hemolytic and fungicidal activities of cationic antimicrobial peptides. *FEBS Lett* 23;449:105-10.
- Helmerhorst EJ, Reijnders IM, van't Hof W, Simoons-Smit I, Veerman EC, Amerongen AV. (1999) Amphotericin B- and fluconazole-resistant *Candida* spp., *Aspergillus fumigatus*, and other newly emerging pathogenic fungi are susceptible to basic antifungal peptides. *Antimicrob Agents Chemother* 43:702-4.
- Jones EM, Smart A, Bloomberg G, Burgess L, Millar MR (1992). Lactoferricin, a new antimicrobial peptide. *J Appl Bacteriol.* 77:208-14.

Lal K, Santarpia RP, Xu L, Manssuri F, Pollock JJ (1992). One-step purification of histidine-rich polypeptides from human parotid saliva and determination of anti-candidal activity. *Oral Microbiol Immunol* 7: 44-50.

Lupetti A, Paulusma-Annema A, Welling MM, Senesi S, van Dissel JT, Nibbering PH (2000). Candidacidal activities of human lactoferrin peptides derived from the N terminus. *Antimicrob Agents Chemother.* 44:3257-63.

Murakami Y, Nagata H, Shizukuishi S, Nakashima K, Okawa T, Takigawa M, Tsunemitsu A (1994). Histatin as a synergistic stimulator with epidermal growth factor of rabbit chondrocyte proliferation. *Biochem Biophys Res Commun* 198:274-80.

Nikawa H, Hayashi S, Nikawa Y, Hamada T, Samaranayake LP (1993). Interactions between denture lining material, protein pellicles and *Candida albicans*. *Arch Oral Biol* 38:631-4.

Nikawa H, Samaranayake LP, Tenovuo J, Pang KM, Hamada T (1993). The fungicidal effect of human lactoferrin on *Candida albicans* and *Candida krusei*. *Arch Oral Biol* 38:1057-63.

Nikawa H, Samaranayake LP, Tenovuo J, Hamada T (1994). The effect of antifungal agents on the in vitro susceptibility of *Candida albicans* to apo-lactoferrin. *Arch Oral Biol* 39: 921-3.

Nikawa H, Samaranayake LP, Hamada T (1995). Modulation of the anti-*Candida* activity of apo-lactoferrin by dietary sucrose and tunicamycin in vitro. *Arch Oral Biol* 40: 581-4.

Nikawa H, Nishimura H, Yamamoto T, Hamada T, Samaranayake LP (1996). The role of salivary or serum pellicle in the colonization process of *Candida albicans* on denture acrylic in vitro. *Microbial Ecol Health Dis* 9: 35-48.

Nikawa H, Jin C, Makihiro S, Hamada T, Samaranayake LP (2002). Susceptibility of *Candida albicans* isolates from the oral cavities of HIV-positive patients to histatin-5. *J Prosthet Dent.* 88:263-7.

Oppenheim FG, Xu T, McMillian FM, Levitz SM, Diamond RD, Offner GD, Troxler RF (1988). Histatins, a novel family of histidine-rich proteins in human parotid secretion. Isolation, characterization, primary structure, and fungistatic effects on *Candida albicans*. *J Biol Chem* 263: 7472-7.

Perczel A, Hollosi M (1996). Turns, in *Circular Dichroism and the conformational Analysis of Biomolecules*. Fasman ed. Plenum Press, New York.

Perczel A, Park K, Fasman GD (1992). Analysis of the circular dichroism spectrum of proteins using the convex constraint algorithm: a practical guide. *Anal Biochem.* 203:83-93.

Pollock JJ, Denepitiya L, MacKay BJ, Iacono VJ (1984). Fungistatic and fungicidal activity of human parotid salivary histidine-rich polypeptides on *Candida albicans*. *Infect Immun* 44:702-7.

Samaranayake, L.P., MacFarlane, T.W., Lamey, P-J., and Ferguson, M.M. (1986) A comparison of oral rinse and imprint sampling techniques for the detection of yeast, coliform and *Staphylococcus aureus* carriage in the oral cavity. *J Oral Pathol*, 15: 251-254.


Samaranayake, L. P. (2002) *Essential Microbiology for Dentistry*, Churchill Livingstone, Edinburgh, UK, Second edition.

Samaranayake, L.P. and Holmstrup, P. (1989) Oral candidiasis and the human immunodeficiency virus infection. *J Oral Pathol Med* 18 554-564.


Samaranayake LP, Cheung LK, Samaranayake YH (2002). Candidiasis and other fungal diseases of the mouth *Dermatol Ther* 15: 252-270.

- Shai Y. (1999) Mechanism of the binding, insertion and destabilization of phospholipid bilayer membranes by alpha-helical antimicrobial and cell non-selective membrane-lytic peptides. *Biochim Biophys Acta* 1462:55-70.
- Silverman Jr. S, Migliorati CA, Epstein, JB, Samaranayake LP (1990). Laboratory diagnosis of oral candidosis. in *Oral Candidosis Samaranayake LP & MacFarlane TW ed. London: Butterworth & Co. Ltd; p 213-237.*
- Situ H, Bobek LA (2000). In vitro assessment of antifungal therapeutic potential of salivary histatin-5, two variants of histatin-5, and salivary mucin (MUC7) domain 1. *Antimicrob Agents Chemother* 44:1485-93.
- Stemberg S (1994). The emerging fungal threat. *Science* 266: 1632-4. Tobgi RS, Samaranayake LP, MacFarlane TW (1987). The in vitro susceptibility of *Candida* species to lysozyme. *Oral Microbiol Immunol* 3: 35-9.
- Tsai H, Bobek LA (1997). Human salivary histatin-5 exerts potent fungicidal activity against *Cryptococcus neoformans*. *Biochim Biophys Acta* 1336:367-9.
- Ueta E, Tanida T, Osaki T (2001). A novel bovine lactoferrin peptide, FKRRWQWRM, suppresses *Candida* cell growth and activates neutrophils. *J Pept Res.* 57:240-9.
- Wakabayashi H, Abe S, Teraguchi S, Hayasawa H, Yamaguchi H (1998). Inhibition of hyphal growth of azole-resistant strains of *Candida albicans* by triazole antifungal agents in the presence of lactoferrin-related compounds. *Antimicrob Agents Chemother.* 42:1587-91.
- White TC, Marr KA, Bowden RA (1998). Clinical, cellular, and molecular factors that contribute to antifungal drug resistance. *Clin Microbiol Rev* 11:382-402.
- Xu T, Levitz SM, Diamond RD, Oppenheim FG (1991). Anticandidal activity of major human salivary histatins. *Infect Immun* 59:2549-54.


a) Dhvar4




b) LFBshort



c) JH8194



d) JH8195



e) JH8944




Fig. 1 Nikawa et al.