

REVIEW

A Checklist of the Parasites of Eels (*Anguilla* spp.) (Anguilliformes: Anguillidae) in Japan (1915–2007)

Kazuya NAGASAWA¹⁾, Tetsuya UMINO¹⁾ and Kouki MIZUNO²⁾

¹⁾ Graduate School of Biosphere Science, Hiroshima University
1-4-4 Kagamiyama, Higashi-Hiroshima, Hiroshima 739-8528, Japan
²⁾ Ehime Prefectural Uwajima Fishery High School
1-2-20 Meirin, Uwajima, Ehime 798-0068, Japan

Abstract Information on the protistan and metazoan parasites of three species of eels, the Japanese eel *Anguilla japonica*, the giant mottled eel *A. marmorata* and the European eel *A. anguilla*, in Japan is summarized in the Parasite-Host List and Host-Parasite lists, based on the literature published for 93 years between 1915 and 2007. Both *A. japonica* and *A. marmorata* are native to Japan, whereas *A. anguilla* is an introduced species from Europe. The parasites, including 44 named species and those not identified to species level, are listed by higher taxon as follows: Sarcomastigophora (no named species), Ciliophora (6), Microspora (1), Myxozoa (6), Trematoda (7), Monogenea (7), Cestoda (3), Nematoda (7), Acanthocephala (4), Hirudinida (2), and Copepoda (1). For each taxon of parasite, the following information is given: its currently recognized scientific name, any original combination, synonym(s), or other previous identification used for the parasite occurring in eels; habitat (freshwater, brackish, or marine); site(s) of infection within or on the host; known geographical distribution in Japanese waters; and the published source of each locality record. Of the 44 named species of parasites, 43 are from *A. japonica*, 1 from *A. marmorata*, and 10 from *A. anguilla*. *Gyrodactylus anguillae* (Monogenea) and *Raphidascaris acus* (Nematoda) are the exotic species that were probably introduced on *A. anguilla* from Europe, and it has also been suggested that *G. nipponensis* (Monogenea) was introduced on eels imported from somewhere in the Indo-western Pacific. Of the parasites recorded from *A. japonica*, eight species are marine and/or brackish-water species, of which two (i.e., *Tubulovesicula anguillae*, *Pseudodactylogyrus kamegaii*) are host-specific and occur as adults. This indicates that *A. japonica* serves as the definitive host for these parasites in marine and/or brackish waters where the host species constantly remains.

Key words: *Anguilla anguilla*; *Anguilla japonica*; *Anguilla marmorata*; checklist; eels; parasites

INTRODUCTION

Eels of the genus *Anguilla* are commercially important fishes in Japan, where the Japanese eel *A. japonica* Temminck and Schlegel and the giant mottled eel *A. marmorata* Quoy and Gaimard occur naturally (Matsui, 1972; Hatooka, 2002). *Anguilla japonica* is widely distributed in Japan and is one of the most important fishes in freshwater culture and commercial catch. The distribution of *A. marmorata* in Japan is confined to the southern region and its catch small. On the other hands,

since the late 1960's, as many as 10 species of eels have been imported to Japan from other countries for culture in ponds or stocking in rivers and lakes (Tabeta et al., 1977). Of these species, some fish of the European eel *A. anguilla* (Linnaeus), the American eel *A. rostrata* (Lesueur) and the shortfin eel *A. australis* Richardson, 1841 have been actually recorded from natural waters in Japan (Tabeta et al., 1976, 1977; Zang et al., 1999; Aoyama et al., 2000; Okamura et al., 2001, 2002; Miyai et al., 2004).

In 1915, Dr. S. Ishii published a paper dealing with myxozoans found on *A. japonica*, which is the first report on the parasites of eels in Japan (Ishii, 1915). Since then, due to the importance of eels in aquaculture and fisheries, numerous studies and investigations have been conducted on the eel parasites in the country. In the present checklist, based on the literature published for 93 years between 1915 and 2007, information on the parasites of eels in Japan is compiled in two lists, Parasite-Host List and Host-Parasite List. In total, 44 named species of parasites are listed along with those not identified to species level. The parasites listed herein are from three species of eels (*A. japonica*, *A. marmorata* and *A. anguilla*). No parasites have been reported from the other eels in Japan.

In the **PARASITE-HOST LIST**, parasites are arranged by higher taxon in the following order: Sarcomastigophora, Ciliophora, Microspora, Myxozoa, Trematoda, Monogenea, Cestoda, Nematoda, Acanthocephala, Hirudinida, and Copepoda. The format is almost the same as in the checklists of Margolis and Arthur (1979) and McDonald and Margolis (1995). Within each higher taxon, genera and species are listed alphabetically. For each taxon of parasite, the following information is provided:

1) The current **scientific name**, including author(s) and date(s), followed by any original combination, recognized synonym(s), or other identification(s) that have been used in establishing records from eels in Japan. No attempt has been made to evaluate the taxonomic validity of the published reports.

2) The **habitat** in which the parasite was acquired and normally completes its life cycle is given as FW for fresh waters, B for brackish waters, and M for marine waters.

3) The **Site(s) of infection** of the parasite in or on its host. If the site was not given in the original record, the likely site was determined from other records and is enclosed in square brackets.

4) The **Distribution** of the parasite is indicated by prefecture (boundaries shown in Fig. 1), in geographical order from northeast to southwest. For marine or brackish-water species, the name of the prefecture nearest the collection site is given.

5) The **Record(s)**. The authors responsible for the records are listed in chronological order. If a parasite has been reported more than once, the references are numbered, but not when there has been only one record of the parasite. Each reference is followed by the locality or localities given in two parts, first the prefecture(s) and then the detailed collection locality or localities from which the parasite was reported. If no locality record was given, the geographical locality is shown by a dash (—). When all records are from the same prefecture, only the detailed collection locality or localities are listed.

6) Under **Remarks**, explanatory comments are given on systematics, nomenclature, useful references, and notes on specific items such as tentative parasite identifications in the original reports.

7) The **References** section includes works directly cited in the Parasite-Host List; the **Supplementary References** are publications dealing with various aspects of parasites of eels in Japan but not containing original parasite records. If only a Japanese title was given by the original author(s), our translation of the title into English is provided in square brackets.

In the **HOST-PARASITE LIST**, hosts are listed alphabetically. In each higher taxon, parasites




Fig. 1. Map of Japan showing the prefectoral boundaries. The following prefectoral names are arranged in alphabetical order: Aichi-24; Akita-7; Aomori-5; Chiba-15; Ehime-41; Etorofu Island-1; Fukui-26; Fukuoka-43; Fukushima-10; Gifu-23; Gunma-14; Hiroshima-37; Hokkaido-4; Hyogo-33; Ibaraki-12; Ishikawa-25; Iwate-6; Kagawa-39; Kagoshima-49; Kanagawa-18; Kochi-42; Kumamoto-47; Kunashiri Island-3; Kyoto-29; Mie-28; Miyagi-8; Miyazaki-48; Nagano-20; Nagasaki-45; Nara-30; Niigata-11; Oita-46; Okayama-35; Okinawa-50; Osaka-31; Saga-44; Saitama-16; Shiga-27; Shikotan Island-2; Shimane-36; Shizuoka-21; Tochigi-13; Tokushima-40; Tokyo-17; Tottori-34; Toyama-22; Wakayama-32; Yamagata-9; Yamaguchi-38; and Yamanashi-19.

are also listed in alphabetical order, and after the name of each parasite, its geographical distribution is given in parenthesis.

This checklist is the sixth in the following series of published synopses of the parasites of commercially important fishes and shellfishes in Japan: Nagasawa et al. (1987) for the parasites of salmonids; Nagasawa et al. (1989) for the parasites of freshwater fishes in Hokkaido; Nagasawa (1993a) for the parasites of squids and cuttlefishes; Nagasawa (1993b) for the parasites of gadids; and Nagasawa et al. (2007a) for the parasites of ayu (*Plecoglossus altivelis altivelis*).

PARASITE-HOST LIST

SARCOMASTIGOPHORA

Cryptobia sp.

(FW)

Hosts: *Anguilla anguilla*

Anguilla japonica

Sites of infection: skin, fins
 Distribution: unknown
 Record: Niwa 1979 (—)

Ichthyobodo sp.

(FW)

Includes: *Costia* sp. (erroneously as "Chostia") of Niwa, 1979

Hosts: *Anguilla anguilla*

Anguilla japonica

Sites of infection: skin, fins
 Distribution: unknown
 Record: Niwa 1979 (—)

Trypanosoma sp.

(FW)

Host: *Anguilla japonica*

Site of infection: blood

Distribution: Shizuoka

Records: 1. Hoshina and Sano 1957 (Yoshida); 2. Egusa 1967 (Yoshida)

CILIOPHORA*Ambiphrya* sp.

(FW)

Host: *Anguilla japonica*

Sites of infection: gills, skin

Distribution: unknown

Record: Egusa 1978 (—)

Aplosoma sp.

(FW)

Includes: *Glossatella* sp. of Nishio et al., 1970; Egusa, 1970; Hatai and Egusa, 1973; Niwa, 1979

Hosts: *Anguilla anguilla* (3, 5)

Anguilla japonica (1, 2, 4)

Site of infection: gills

Distribution: Shizuoka

Records: 1. Nishio et al. 1970 (Yoshida); 2. Egusa 1970 (Yoshida); 3. Hatai and Egusa 1973 (Yaizu, Yoshida); 4. Egusa 1978 (—); 5. Niwa 1979 (—)

Capriniata piscium (Buetschli, 1889) Jankowski, 1973

(FW)

Previous identification: *Trichophrya piscium* of Egusa, 1978

Includes: *Trichophrya* sp. of Egusa and Ahmed, 1970; Nishio et al., 1970; Egusa, 1970, 1971

Hosts: *Anguilla anguilla* (1, 2, 4, 5)

Anguilla japonica (1, 3, 4)

Site of infection: gills

Distribution: Shizuoka

Records: 1. Egusa and Ahmed 1970 (Yaizu); 2. Nishio et al. 1970 (Yoshida); 3. Egusa 1970 (Yoshida); 4. Egusa 1971 (—); 5. Egusa 1978 (—)

Remarks: Matsui (1972: 577–578, figs. 27.44, 27.45) reported the occurrence of two ciliates, “*Sayphidia* or *Sayphydia* sp.” and “*Sudonia* sp.” on the gills of *A. japonica*, as well as *Caprinia piscium* (as *Trichophrya* sp.). His identification of the first two species, is definitely not correct.

Carchesium polypinum Linnaeus, 1758

(FW)

Host: *Anguilla japonica*

Site of infection: skin

Distribution: Tokushima

Record: Naruto Station, Fish. Exp. St. Tokushima Pref. 1966 (—)

Chilodonella sp.

(FW)

Hosts: *Anguilla anguilla* (2)

Anguilla japonica (1, 2)

Sites of infection: gills, skin

Distribution: unknown

Records: 1. Egusa 1978 (—); 2. Niwa 1979 (—)

Ichthyophthirius multifiliis Fouquet, 1876

(FW)

Hosts: *Anguilla anguilla* (1, 3, 4, 5, 6, 7, 8, 9)

Anguilla japonica (1, 2, 6, 7, 8, 9)

Sites of infection: skin, fins, gills, buccal cavity

Distribution: Shizuoka

Records 1. Egusa et al. 1970 (Yaizu); 2. Nishio et al. 1970 (Yoshida); 3. Egusa 1971 (—); 4. Oka 1973a (near Lake Hamana); 5. Oka 1973b (—); 6. Egusa 1978 (—); 7. Egusa 1979 (—); 8. Niwa 1979 (—); 9. Egusa 1983 (—)

Trichodina acuta Lom, 1961

(FW)

Host: *Anguilla japonica*

Site of infection: gills

Distribution: Mie

Record: Imai et al. 1991 (Tsu)

Trichodina jadranica Haider, 1964

(FW)

Host: *Anguilla japonica*

Site of infection: gills

Distribution: Mie

Record: Imai et al. 1991 (Tsu)

Remarks: This trichodinid was reported from the gills of *A. japonica* cultured in freshwater ponds in central Japan (Imai et al., 1991). However, it was later found on marine fishes (Japanese flounder *Paralichthys olivaceus* and stone flounder *Kareius bicoloratus*) in China (Xu et al., 2001), suggesting that *T. jadranica* is a euryhaline species.

Trichodina japonica Imai, Miyazaki and Nomura, 1991

(FW)

Host: *Anguilla japonica*

Site of infection: gills

Distribution: Mie

Record: Imai et al. 1991 (Tsu)

Remarks: This trichodinid was described from the gills of *A. japonica* cultured in freshwater ponds in central Japan (Imai et al., 1991). However, it also occurs on marine fishes (Japanese seaperch *Lateolabrax japonicus* and red seabream *Chrysophryrys major*) and a brackish-water fish (barramundi *Lates calcarifer*) in China and India, respectively (Xu et al., 1999, 2001; Mitra and Bandyopadhyay 2005), indicating that *T. japonica* is a euryhaline species, like *T. jadranica* (see above).

Trichodina sp.

(FW)

Hosts: *Anguilla anguilla* (3, 6, 7, 8) *Anguilla japonica* (1, 2, 3, 4, 5, 6, 8)

Sites of infection: gills

Distribution: Shizuoka

Records: 1. Egusa 1967 (Yoshida); 2. Egusa 1968 (Yoshida); 3. Nishio et al. 1970 (Yoshida); 4. Egusa 1970 (Yoshida); 5. Egusa et al. 1971 (Yoshida); 6. Egusa 1971 (—); 7. Hatai and Egusa 1973 (Yaizu, Yoshida); 8. Niwa 1979 (—)

MICROSPORA

Heterosporis anguillarum (Hoshina, 1951) Lom, Dyková, Körting and Klinger, 1989

(FW)

Original combination: *Plistophora anguillarum* Hoshima, 1951Includes: *Plistophora* sp. of Niwa, 1979Hosts: *Anguilla anguilla* (6) *Anguilla japonica* (1, 2, 3, 4, 5, 6, 7, 8, 9)

Site of infection: musculature

Distribution: Hokkaido, Kanagawa, Shizuoka, Aichi, Kagoshima

Records: 1. Hoshina 1951a (Kangawa: near Odawara; Shizuoka: Yoshida); 2. Hoshima 1972 (Kanagawa:—; Shizuoka:—; Aichi:—); 3. Awakura 1974 (Hokkaido: Shikabe); 4. Hashimoto and Takinami 1976 (Shizuoka: Hamanko Branch of Shizuoka Pref. Fish. Exp. St.); 5. Hashimoto et al. 1976 (Shizuoka: Hamanko Branch of Shizuoka Pref. Fish. Exp. St.); 6. Niwa 1979 (Shizuoka:—; Aichi:—; Kagoshima:—); 7. Kano and Fukui 1982 (—); 8. Kano et al. 1982 (—); 9. Buchmann et al. 1992 (Shizuoka:—)

Remarks: The present species was transferred from the genus *Pleistophora* to *Heterosporis* by Lom et al. (1989). Although Awakura (1974) found this parasite in Hokkaido, the infected fish had been transported from Shizuoka. The species is known to infect *A. japonica* in Taiwan (T'sui and Wang, 1988; T'sui et al., 1988; Tsai et al., 2002) and Korea (Suh and Chun, 1988; Joh et al., 2007) as well. Hoshima (1972) reported the presence of this parasite in young *A. japonica* imported from Taiwan to Japan.

Unidentified Microspora

(FW)

Host: *Anguilla japonica*

Site of infection: gills

Distribution: Shizuoka

Record: Egusa 1967 (Yoshida)

MYXOZOA

Myxidium giardi Cépède, 1906

(FW)

Synonyms: *Myxidium anguillae* Ishii, 1915; *Myxidium enchelypterygii* Hoshina, 1952

Includes: *Myxidium* sp. of Ishii, 1916; Iwata, 1972

Hosts: *Anguilla anguilla* (5)

Anguilla japonica (1, 2, 3, 4, 6)

Sites of infection: skin, fins, gills

Distribution: Tokyo, Shizuoka, Miyazaki

Records 1. Ishii 1915a (Shizuoka: Numazu); 2. Ishii 1916b (Tokyo:—); 3. Hoshina 1952 (Shizuoka: Yoshida Fish-Cultural Laboratory); 4. Iwata 1972 (Miyazaki: Hosoda River); 5. Hine 1980 (—); 6. Oka and Egusa 1989 (Shizuoka: Hamamatsu)

Remarks: Although Hoshina (1952) reported that the spores of *Myxidium enchelypterygii* were clearly differentiated from those of *M. anguillae* by their size and shape, Hine (1980) regarded both taxa as identical, which was supported by Oka and Egusa (1989). Hine (1980: table 1) listed a record of *M. giardi* found in the gall bladder and musculature of the American eel *Anguilla rostrata* from Japan, but this record is not herein included because no references were found to support it.

Myxidium lentiforme Fujita, 1929

(FW)

Synonym: *Myxidium fusiforme* Fujita, 1927

Host: *Anguilla japonica*

Site of infection: kidney

Distribution: Shiga

Record: Fujita 1927 (Lake Biwa)

Remarks: This parasite had been originally described by Fujita (1927) as *M. fusiforme*, but it was later renamed as *Myxidium lentiforme* by Fujita (1929: 249–250) because the former had been preoccupied.

Myxidium matsuii Fujita, 1929

(FW)

Host: *Anguilla japonica*

Site of infection: skin

Distribution: Kanagawa, Shizuoka, Aichi

Records: 1. Fujita 1929 (Shizuoka: near Lake Hamana; Aichi: Toyohashi); 2. Hoshina 1952 (Kanagawa: Odawara); 3. Egusa 1978 (—); 4. Hine 1980 (—)

Myxidium uchiyamae Fujita, 1927

(FW)

Host: *Anguilla japonica*

Site of infection: kidney

Distribution: Shiga

Record: Fujita 1927 (Lake Biwa)

Myxidium sp.

(FW)

Hosts: *Anguilla anguilla* (3, 4, 6) *Anguilla japonica* (1, 2, 5, 6)

Sites of infection: gills, kidney, liver

Distribution: Shizuoka

Records: 1. Egusa 1967 (Yoshida); 2. Egusa 1970 (Yoshida); 3. Oka 1973a (near Lake Hamana); 4. Oka 1973b (—); 5. Ushiyama and Misaki 1977 (Hamamatsu); 6. Niwa 1979 (—)

Remarks: There is no information on the morphology and identification of this myxozoan. Niwa (1979) reported that the spores of *Myxidium* sp. are more commonly found in the kidney of *A. anguilla* than *A. japonica*.*Myxobolus dermatobius* (Ishii, 1915) Landsberg and Lom, 1991

(FW)

Original combination: *Lentospora dermatobia* Ishii, 1915Previous identification: *Myxosoma (Lentospora) dermatobia* (Ishii, 1915)Host: *Anguilla japonica*

Site of infection: skin

Distribution: Tochigi, Shizuoka

Records: 1. Ishii 1915b (Shizuoka: Numazu); 2. Hoshina 1952 (Tochigi: Lake Chuzenji)

Remarks: The present species, initially described as *Lentospora dermatobia* by Ishii (1915b), was transferred to the genus *Myxobolus* by Landsberg and Lom (1991).*Myxobolus fujitai* (Fujita, 1929) Eiras, Molnár and Lu, 2005

(FW)

Synonym: *Lentospora anguillae* Fujita, 1929Host: *Anguilla japonica*

Site of infection: skin

Distribution: Ibaraki

Record: Fujita 1929 (Lake Hinuma)

Remarks: The present species was initially described as *Lentospora anguillae* by Fujita (1929) but it was later renamed as *Myxobolus anguilli* by Landsberg and Lom (1991). However, because of the preoccupation of the latter name, Eiras et al. (2005) proposed a new name, *Myxobolus fujitai*, for it.

Unidentified Myxozoa

(FW)

Host: *Anguilla japonica*

Site of infection: gills

Distribution: Shizuoka, Gifu

Records: 1. Nishio et al. 1970 (Shizuoka: Yoshida); 2. Nishio et al. 1971 (Shizuoka: Yoshida); 3.

Anonymous 2002 (Gifu: a tributary of the Kiso River)

TREMATODA

Azygia gotoi (Ariake, 1922) Shimazu, 1979 (FW)

Original combination: *Azygia anguillae* Ozaki, 1924

Includes: *Azygia gotoi*-like trematodes of Shimazu, 1979

Host: *Anguilla japonica*

Site of infection: stomach

Distribution: Aomori, Ibaraki, Tokyo, Nagano, Shiga

Records: 1. Ozaki 1924 (Tokyo:—); 2. Yamaguti 1934a (Ibaraki: Lake Kasumiga-ura [as “Kasumiga-ura”]); 3. Shimazu 1979 (Aomori: Lake Hira-numa; Nagano: Lake Kizaki, Lake Suwa; Shiga: Lake Biwa)

Remarks: The taxonomy and life history of this trematode was reported in details by Shimazu (1979). Information on the species is available from Shimazu (1999a, 2003).

Centrocestus formosanus (Nishigori, 1924) Price, 1932 (metacercaria) (FW)

Host: *Anguilla japonica*

Habitat: gills

Distribution: Kagoshima

Records: 1. Yanohara and Kagei 1983 (Tanegashima Island); 2. Kagei and Yanohara 1995 (Tanegashima Island)

Genarchopsis goppo Ozaki, 1925 (FW)

Synonym: *Genarchopsis anguillae* Yamaguti, 1938

Host: *Anguilla japonica*

Site of infection: intestine

Distribution: Ibaraki, Shiga

Records: 1. Yamaguti 1938 (Ibaraki: Tsuchiura [as Tutiura]); 2. Shimazu 1995 (Ibaraki: Tsuchiura; Shiga: Lake Biwa)

Remarks: Moravec et al. (2003) regarded *Genarchopsis anguillae* as a junior synonym of *G. goppo*. Shimazu and Urabe (2005) later made a similar suggestion. Information on the species is available from Shimazu (1999a, 2003).

Hemuiridae gen. sp. (FW?)

Host: *Anguilla japonica*

Site of infection: stomach

Distribution: Tokyo

Record: Ozaki 1924 (—)

Remarks: When *Azygia gotoi* (as *A. anguillae*) was described, Ozaki (1924: 426) reported that another trematode belonging to the family Hemuiridae was in the stomach of *A. japonica*. No description of this trematode is yet available.

Lasiotocus sp. (B or FW)

Host: *Anguilla japonica*

Site of infection: intestine

Distribution: Aomori

Record: Shimazu 2005 (Lake Ogawara)

Lecithochrimum musculus (Looss, 1907) Nasir and Diaz, 1971 (M)

Synonym: *Sterrurus musculus* Looss, 1907

Host: *Anguilla japonica*

Site of infection: stomach

Distribution: Mie, unspecified prefecture facing the Seto Inland Sea

Record: Yamaguti 1934a (Mie: Ise Bay; unspecified prefecture: Seto Inland Sea [as Inland Sea])

Remarks: The identification of this trematode by Yamaguti (1934) needs confirmation (Gibson and Bray, 1986: 83–90).

Metagonimus spp. (metacercaria) (FW)

Host: *Anguilla japonica*

Site of infection: fins

Distribution: Shizuoka

Record: Ito and Mochizuki 1968 (Tenryu River)

Phyllodistomum anguiae Long and Wai, 1958 (FW)

Host: *Anguilla japonica*

Sites of infection: urinary bladder, intestine

Distribution: Aomori, Ibaraki

Record: Shimazu 2005 (Aomori: Lake Ogawara; Ibaraki: Tsuchiura)

Procotrematoides pisodontophidis Yamaguti, 1938 (M)

Host: *Anguilla japonica*

Site of infection: intestine

Distribution: Chiba

Record: Hoshina 1951b (Urayasu)

Tubulovesicula anguillae Yamaguti, 1934 (M)

Host: *Anguilla japonica*

Site of infection: stomach

Distribution: Miyagi

Record: Yamaguti 1934a (Matsushima Bay [as Matusima Bay])

MONOGENEA

Gyrodactylus anguillae Ergens, 1960 (FW)

Host: *Anguilla anguilla*

Sites of infection: skin, gills

Distribution: Shizuoka

Record: Ogawa and Egusa 1980 (Maisaka)

Remarks: Ogawa and Egusa (1980) suggested that this species was introduced into Japan from France on *A. anguilla*. Hayward et al. (2001) showed the current worldwide distribution of the species. Ogawa and Egusa (1978) redescribed it based on the specimens from England.

Gyrodactylus egusai Ogawa and Hioki, 1986

(FW)

Host: *Anguilla japonica*

Site of infection: skin

Distribution: Shizuoka

Record: Ogawa and Hikoki (Yoshida)

Gyrodactylus joi Ogawa and Hioki, 1986

(FW)

Host: *Anguilla japonica*

Site of infection: skin

Distribution: Shizuoka

Record: Ogawa and Hikoki (Yoshida)

Gyrodactylus nipponensis Ogawa and Egusa, 1978

(FW or B)

Host: *Anguilla japonica*

Site of infection: gills

Distribution: Chiba, Shizuoka, Tokushima, Miyazaki

Records: 1. Ogawa and Egusa 1978 (Shizuoka:—; Tokushima:—); 2. Ogawa and Egusa 1980 (Chiba:—; Shizuoka:—; Tokushima:—; Miyazaki:—); 3. Hayward et al. 2001 (Chiba: Minato River, Shizuoka: Lake Hamana)

Remarks: Hayward et al. (2001: 422) suggested that this monogenean was introduced into Japan on eels imported from somewhere in the Indo-western Pacific.

Gyrodactylus sp.

(FW)

Host: *Anguilla japonica*

Site of infection: gills

Distribution: Shizuoka

Record: Ushiyama and Misaki 1977 (Hamamatsu)

Remarks: There is no information on the morphology and taxonomy of this gyrodactylid. Identification needs to be confirmed in comparison with the four species of *Gyrodactylus* reported from eels in Japan.

Pseudodactylogyrus anguillae (Yin and Sproston, 1948) Gusev, 1965

(FW)

Synonym: *Pseudodactylogyrus microrchis* Ogawa and Egusa, 1976

Hosts: *Anguilla anguilla* (1, 2, 3, 5, 6)

Anguilla japonica (2)

Anguilla sp. (4)

Site of infection: gills

Distribution: Chiba, Shizuoka, Aichi, Tokushima, Kagoshima

Records: 1. Ogawa and Egusa 1976 (Chiba:—; Shizuoka:—); 2. Ogawa et al. 1985 (Chiba:—; Aichi:—; Tokushima:—); 3. Iwashita et al. 2002 (Shizuoka: Maisaka); 4. Hayward 2004 (Aichi:—; Kagoshima: Yaku Island); 5. Yoshikawa 2005 (Shizuoka: Hamanako Branch of Shizuoka Pref. Fish. Exp. St.); 6. Umeda et al. 2006 (Kagoshima: Ibusuki Branch of Kagoshima Pref. Fish. Center);

Remarks: Ogawa et al. (1985) synonymised *P. microrchis* as a junior synonym of *P. anguillae*.

Pseudodactylogyrus bini (Kikuchi, 1929) Gusev, 1965

(FW)

Original combination: *Dactylogyrus bini* Kikuchi, 1929

Hosts: *Anguilla anguilla* (2, 4)

Anguilla japonica (1, 5)

Anguilla sp. (3)

Site of infection: gills

Distribution: Chiba, Shizuoka, Aichi, Kagoshima

Records: 1. Kikuchi 1929 (—); 2. Ogawa and Egusa 1976 (Chiba:—; Shizuoka:—); 3. Hayward 2004 (Aichi:—; Kagoshima: Yaku Island); 4. Umeda et al. 2006 (Kagoshima: Ibusuki Branch of Kagoshima Pref. Fish. Center); 5. Sato and Tanaka 2007 (Shizuoka: near Lake Hamana)

Pseudodactylogyrus kamegaii Iwashita, Hirata and Ogawa, 2002

(B)

Host: *Anguilla japonica*

Site of infection: gills

Distribution: Chiba

Record: Iwashita et al. 2002 (Minato River)

Remarks: This species was found on eels collected in brackish waters (Iwashita et al., 2002).

Pseudodactylogyrus spp.

(FW)

Includes: *Dactylogyrus* sp. of Kikuchi, 1929; Egusa and Ahmed, 1970; Egusa, 1970, 1971; Oka, 1973a; Hatai and Egusa, 1973; Ushiyama and Misaki, 1977

Pseudodactylogyrus bini or *P. anguillae* of Tanaka and Sato, 2007; Sato and Tanaka, 2007
“*Pseudodactylogyrus* sp. ang. 4” of Hayward, 2004

Hosts: *Anguilla anguilla* (2, 4, 5, 6, 8, 10)

Anguilla japonica (1, 3, 7, 8, 11, 12)

Anguilla sp. (9)

Site of infection: gills

Distribution: Shizuoka, Kagoshima

Records: 1. Kikuchi 1929 (—); 2. Egusa and Ahmed 1970 (Shizuoka: Yaizu); 3. Egusa 1970 (Shizuoka: Yoshida); 4. Egusa 1971 (—); 5. Oka 1973a (Shizuoka: near Lake Hamana); 6. Hatai and Egusa 1973 (Shizuoka: Yaizu, Yoshida); 7. Ushiyama and Misaki 1977 (Shizuoka: Hamamatsu); 8. Niwa 1979 (—); 9. Hayward 2004 (Kagoshima: Yaku Island); 10. Yoshikawa et al. 2006 (Shizuoka: Hamana Branch of Shizuoka Pref. Fish. Exp. St.); 11. Tanaka and Sato 2007 (Shizuoka: near Lake Hamana); 12. Sato and Tanaka 2007 (Shizuoka: near Lake Hanama)

Unidentified Monogenea

(FW)

Includes: *Gyrodactylus* sp. or *Dactylogyrus* sp. of Nishio et al., 1970

“Monogenetic trematodes” of Shimazu, 1979

Hosts: *Anguilla anguilla* (1) *Anguilla japonica* (1, 2)

Site of infection: gills

Distribution: Nagano, Shizuoka

Records: 1. Nishio et al. 1970 (Shizuoka: Yoshida); 2. Shimazu 1979 (Nagano: Lake Kizaki)

CESTODA

Bothriocephalus claviceps (Goeze, 1782) Rudolphi, 1810

(FW)

Hosts: *Anguilla japonica* (?)(2) *Anguilla maromorata* (1, 2)

Site of infection: intestine

Distribution: Shiga, Kagoshima

Records: 1. Luo et al. 2002 (Kagoshima: Yaku Island [as Yako Island]); 2. Scholz et al. 2004 (Shiga: Lake Biwa; Kagoshima: Yaku Island)

Remarks: Identification of the eel from Lake Biwa examined by Scholz et al. (2004) was uncertain: these authors tentatively identified the fish as *A. japonica* but it may be identified as *A. anguilla*. If the eel was actually the latter species, the cestode may have been introduced into the lake by the import of *A. anguilla* from overseas (Scholz et al., 2004).*Bothriocephalus japonicus* Yamaguti, 1934

(FW)

Host: *Anguilla japonica*

Site of infection: intestine

Distribution: Ibaraki, Nagano, Gifu, Shiga

Records: 1. Yamaguti 1934b (Ibaraki: Lake Kasumiga-ura [as “Kasumiga-ura”]); 2. Anonymous 2002 (Gifu: a tributary of the Kiso River); 3. Scholz et al. 2004 (Ibaraki: Kasumiga-ura, Nagano: Lake Suwa, Shiga: Lake Biwa)

Remarks: Information on this cestode is available from Shimazu (1997). The scientific name was misspelled “*japonicum*” in Anonymous (2002).*Bothriocephalus* sp.

(FW)

Host: *Anguilla japonica*

Site of infection: intestine

Distribution: Nagano

Record: Shimazu 1979 (Lake Kizaki)

Remarks: There is no morphological and taxonomic information on this cestode (Shimazu, 1979: 230, footnote).

Nybelinia anguillicola Yamaguti, 1952 (larva)

(M)

Previous identification: *Nybelinia* sp. of Yamaguti, 1934

Host: *Anguilla japonica*

Site of infection: encysted in submucosa of intestine

Distribution: Mie

Records: 1. Yamaguti 1934b (Kuki); 2. Yamaguti 1952 (Kuki)

Unidentified Cestoda

(FW)

Host: *Anguilla japonica*

Site of infection: intestine

Distribution: Shizuoka

Record: Ushiyama and Misaki 1977 (Hamamatsu)

Remarks: There is no information on the morphology and identification of this cestode: it was frequently found from June to September in cultured *A. japonica* (Ushiyama and Misaki, 1977).

NEMATODA

Anguillicola globiceps Yamaguti, 1935

(FW)

Hosts: *Anguilla japonica* (1, 2, 3, 4, 5, 7)

Anguilla sp. (*A. japonica* ?)(6)

Site of infection: swimbladder

Distribution: Aomori, Nagano, Chiba, Shizuoka, Aichi, Okayama

Records: 1. Yamaguti 1935b (Shizuoka: Lake Hamana); 2. Suyehiro 1957 (Okayama:—); 3. Egusa 1978 (Shizuoka:—; Aichi:—); 4. Egusa 1979 (—); 5. Shimazu 1979 (Aomori: Lake Hira-numa, Nagano: Lake Kizaki); 6. Moravec and Taraschewski 1988 (—); 7. Hirose et al. 1998 (Chiba: Tone River)

Remarks: The biology of this nematode was reviewed by Nagasawa et al. (1994) and Shimazu (1998). A brief review on *Anguillicola* is available in Salati (1987). Although Egusa et al. (1969) reported *A. globiceps* from Japanese eels cultured in Shizuoka, Hirose et al. (1976: 27, footnote) reported that Egusa et al.'s worms were not *A. globiceps* but *A. crassus*. The latter authors also mentioned that the morphology of the worms collected at an eel farm in Mishima, Shizuoka was similar to that of *A. globiceps*.

Anguillicoloides crassus (Kuwahara, Niimi and Itagaki, 1974) Moravec and Taraschewski, 1988 (FW)

Original combination: *Anguillicola crassa* Kuwahara, Niimi and Itagaki, 1974

Previous identification: *Anguillicola globiceps* of Egusa et al. 1969

Includes: *Anguillicola japonica* of Matsui, 1972

Anguillicola sp. of Egusa and Ahmed, 1970; Ushiyama and Misaki, 1977

"Swimbladder nematodes" of Egusa, 1970

Hosts: *Anguilla anguilla* (1, 2, 5, 9, 10)

Anguilla japonica (1, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17)

Site of infection: swimbladder

Distribution: Chiba, Shizuoka, Gifu, Aichi, Mie, Okayama, Tokushima, Oita, Miyazaki, Okinawa

Records: 1. Egusa et al. 1969 (Shizuoka: Yoshida); 2. Egusa and Ahmed 1970 (Shizuoka: Yaizu); 3. Egusa 1970 (Shizuoka: Yoshida); 4. Matsui 1972 (—); 5. Kuwahara et al. 1974 (Shizuoka: Hamamatsu); 6. Hirose et al. 1976 (Shizuoka:—; Aichi:—); 7. Ushiyama and Misaki 1977

(Shizuoka: Hamamatsu); 8. Egusa 1978 (Chiba: eel farm, Lake Inba-numa, Tone River; Shizuoka: eel farm, Lake Hamana; Aichi: eel farm; Mie: eel farm; Okayama: Kojima Bay; Tokushima: eel farm; Oita: eel farm; Miyazaki: Oyodo River; Miyazaki: eel farm; Okinawa: eel farm); 9. Egusa 1979 (—); 10. Niwa 1979 (—); 11. Moravec and Taraschewski 1988 (Shizuoka [as "Shizuka"]:—); 12. Inui et al. 1998 (Shizuoka:—); 13. Hirose et al. 1998 (Aichi: Mikawa); 14. Ushikoshi et al. 1999 (—); 15. Inui et al. 1999 (Shizuoka:—); 16. Anonymous 2002 (Gifu: a tributary of the Kiso River); 17. Moravec et al. 2005 (Aichi: Isshiki)

Remarks: The biology of this nematode was reviewed by Nagasawa et al. (1994) and Moravec (2006). Information on the species is also available from Shimazu (1998). A brief note on the nematode is also published by Salati (1987). Although Matsui (1972: 571) stated infection of "*Anguillilicola japonica*" in the "gall bladder" of *Anguilla japonica*, the worm is identifiable as *A. crassus*, based on a picture (fig. 27.36) shown by him (see Nagasawa et al., 1994: 128). The records reported by Inui et al. (1998, 1999) were based on the species from *A. japonica* imported from Taiwan to Japan. Information on the life cycle of the nematode in Japan is available in Hirose et al. (1976) and Moravec et al. (2005).

Cucullanus filiformis Yamaguti, 1935

(M)

Host: *Anguilla japonica*

Site of infection: intestine

Distribution: Mie

Record: Yamaguti 1941 (Hamajima)

Remarks: This nematode was originally reported from conger eels *Conger myriaster* in Japan (Yamaguti, 1935).

Gnathosoma spinigerum Owen, 1836 (larva)

(FW)

Host: *Anguilla japonica*

Site of infection: musculature

Distribution: unspecified prefecture in Kyushu

Records: 1. Kikuchi 1956 (—); 2. Miyazaki 1963 (unspecified prefecture in Kyushu:—)

Heliconema longissimum (Ortlepp, 1923)

(B)

Synonym: *Heliconema anguillae* Yamaguti, 1935

Host: *Anguilla japonica*

Site of infection: stomach

Distribution: unknown

Records: 1. Yamaguti 1935b (—); 2. Matsui 1972 (—)

Remarks: Matsui (1972: fig. 27.33) showed pictures of the stomach of *A. japonica* heavily infected by this nematode. Information on the nematode is available from Shimazu (1998). Moravec et al. (2007) recently redescribed the species from *Pisodonophis boro* (Ophichthidae) in Thailand.

Heliconema sp.

(?)

Host: *Anguilla japonica*

Site of infection: digestive tract

Distribution: Okayama

Record: Suyehiro 1957

Remarks: The morphology of this nematode is different from that of *H. longissimum* (as *H. anguillae*) (Suyehiro, 1957).

Philometroides anguillae (Ishii, 1916) Rasheed, 1963

(FW)

Original combination: *Filaria anguillae* Ishii, 1916

Host: *Anguilla japonica*

Site of infection: orbit

Distribution: Tokyo, Aichi

Records: 1. Ishii 1916a (Tokyo: Fukagawa-Fuyuki; Aichi: Toyohashi); 2. Ishii 1931 (Tokyo: Fukagawa-Fuyuki; Aichi: Toyohashi)

Remarks: Yamaguti (1935) suggested that the species described by Ishii (1916a) as “*Filaria anguillae*” should be placed in the genus *Philometra*. Later Rasheed (1963) transferred it to the genus *Philometroides*. Matsui (1972: 584) mistakenly reported the species as “*Philometra parasiluri*.” Information on the species is available from Shimazu (1998) and Moravec (2006: 425–427).

Raphidascaris acus (Bloch, 1779) Railliet and Henry, 1915

(FW)

Host: *Anguilla japonica*

Site of infection: intestine

Distribution: Shiga

Record: Grygier and Urabe 2003 (Lake Biwa)

Remarks: This nematode is not a parasite of Japanese fishes. It has been suggested that the nematode was introduced into Japan by the import of *A. anguilla* from overseas (Grygier and Urabe, 2003).

ACANTHOCEPHALA

Acanthocephalus gotoi Van Cleave, 1925

(FW)

Host: *Anguilla japonica*

Site of infection: intestine

Distribution: Tokyo, Aichi

Records: 1. Van Cleave 1925 (Tokyo: fish market); 2. Yamaguti 1935a (various localities in Japan); 3. Fukui and Morisita 1936 (Aichi:—)

Remarks: Information on this acanthocephalan is available from Shimazu (1999b).

Echinorhynchus cottii Yamaguti, 1935

(FW)

Host: *Anguilla japonica*

Site of infection: [intestine]

Distribution: Shiga

Record: Amin et al. 2007 (Lake Biwa)

Remarks: Information on this acanthocephalan is available from Shimazu (1999b).

Longicollum alemniscus (Harada, 1935) Fukui and Morisita, 1937 (immature worm) (M)

Host: *Anguilla japonica*

Site of infection: [intestine]

Distribution: Aichi

Record: Fukui and Morisita 1937 (—)

Remarks: Information on this species is available in Fukui and Morisita (1938) as well. Petrochenko (1956) considered this species as a junior synonym of *Longicollum pagrosomi* but his suggestion has not been supported by Yamaguti (1963), Golvan (1969) and Amin (1985). The species is thus herein treated as a valid species.

Pseudorhadinorhynchus samegaiensis Nakajima and Egusa, 1975 (FW)

Host: *Anguilla japonica*

Site of infection: [intestine]

Distribution: Shiga

Record: Amin et al. 2007 (Lake Biwa)

Remarks: Information on this acanthocephalan is available from Shimazu (1999b).

HIRUDINIDA

Batracobdella smaragdina (Oka, 1910) (FW)

Host: *Anguilla japonica*

Habitat: skin

Distribution: Aichi, Kagoshima

Record: Ogawa et al. 1985 (Aichi: Isshiki; Kagoshima:—)

Hemiclepsis marginata (O. F. Müller, 1774) (FW and B)

Host: *Anguilla japonica*

Site of infection: skin

Distribution: Aichi

Record: Nagasawa and Miyakawa 2006 (river near Akabane Port)

Remarks: Although this species usually occurs in fresh waters (Burreson, 2006), Nagasawa and Miyakawa (2006) found the specimens on elvers caught in brackish waters.

COPEPODA

Lernaea cyprinacea Linnaeus, 1758 (FW)

Original combination: *Lernaea (Lernaeocera) elegans* Leigh-Sharpe, 1925

Previous identification: *Lernaea elegans* Leigh-Sharpe, 1925

Includes: *Lernaea* sp. of Niwa, 1979

Hosts: *Anguilla anguilla* (10)

Anguilla japonica (1, 2, 3, 4, 5, 6, 7, 8, 9, 10)

Sites of infection: buccal cavity, nostril, orbit, fins

Distribution: Chiba, Shizuoka, Aichi, Mie, Okayama, Hyogo, Shimane, Miyazaki

Records: 1. Leigh-Sharpe 1925 (Aichi: Kitajima near Toyohashi); 2. Matsui and Kumada 1928 (Shizuoka: on the coast of Lake Hamana; Aichi: near Toyohashi, Hekikai County); 3. Nakai and Kokai 1931 (Chiba:—); 4. Yamaguti 1939 (Shizuoka [as "Sizuoka"]:—); 5. Kasahara 1957 (—); 6. Kasahara 1958 (Shizuoka:—; Aichi: Toyohashi; Mie:—); 7. Kasahara 1959 (—); 8. Kasahara 1962 (Shizuoka:—; Aichi:—; Mie:—; Okayama:—; Hyogo:—; Shimane:—; Miyazaki:—); 9. Tsutsumi 1978 (—); 10. Niwa 1979 (—)

Remarks: Information on this copepod as a parasite of *A. japonica* is available from Matsui (1972). The bibliography of the copepod in Japan was published by Nagasawa et al. (2007b).

HOST-PARASITE LIST

<i>Anguilla anguilla</i> (Linnaeus, 1758)	European eel, "yōroppa-unagi" (in Japanese)
Sarcostigiphora	
<i>Cryptobia</i> sp. (—)	
<i>Ichthyobodo</i> sp. (—)	
Ciliophora	
<i>Aplosoma</i> sp. (Shizuoka)	
<i>Caprinata piscium</i> (Shizuoka)	
<i>Chilodonella</i> sp. (—)	
<i>Ichthyophthirius multifiliis</i> (Shizuoka)	
<i>Trichodina</i> sp. (Shizuoka)	
Microspora	
<i>Heterosporis anguillarum</i> (Shizuoka, Aichi, Kagoshima)	
Myxozoa	
<i>Myxidium giardi</i> (—)	
<i>Myxidium</i> sp. (Shizuoka)	
Monogenea	
<i>Gyrodactylus anguillae</i> (Shizuoka)	
<i>Pseudodactylogyrus anguillae</i> (Chiba, Shizuoka, Aichi, Tokushima)	
<i>Pseudodactylogyrus bini</i> (Chiba, Shizuoka, Tokushima)	
<i>Pseudodactylogyrus</i> spp. (Shizuoka)	
Unidentified Monogenea (Shizuoka)	
Nematoda	
<i>Anguillicoloides crassus</i> (Shizuoka)	
Copepoda	
<i>Lernaea cyprinacea</i> (—)	
Remarks: The parasites listed herein are all from cultured <i>A. anguilla</i> . There is no record of parasites from this host occurring in natural waters of Japan.	
<i>Anguilla japonica</i> Temminck and Schlegel, 1847	Japanese eel, "unagi" (in Japanese)
Sarcostigiphora	
<i>Cryptobia</i> sp. (—)	
<i>Ichthyobodo</i> sp. (—)	
<i>Trypanosoma</i> sp. (Shizuoka)	

Ciliophora

- Ambiphrya* sp. (—)
Apiosoma sp. (Shizuoka)
Capriniata piscium (Shizuoka)
Carchesium polypinum (Tokushima)
Chilodonella sp. (—)
Ichthyophthirius multifiliis (Shizuoka)
Trichodina acuta (Mie)
Trichodina jadranica (Mie)
Trichodina japonica (Mie)
Trichodina sp. (Shizuoka)

Microspora

- Heterosporis anguillarum* (Hokkaido, Kanagawa, Shizuoka, Aichi, Kagoshima)
Unidentified Microspora (Shizuoka)

Myxozoa

- Myxidium giardi* (Tokyo, Shizuoka, Miyazaki)
Myxidium lentiforme (Shiga)
Myxidium matsuii (Kanagawa, Shizuoka, Aichi)
Myxidium uchiyamae (Shiga)
Myxidium sp. (Shizuoka)
Myxobolus dermatobius (Tokyo, Shizuoka)
Myxobolus fujitai (Ibaraki)
Unidentified Myxozoa (Shizuoka, Gifu)

Trematoda

- Azygia gotoi* (Aomori, Ibaraki, Tokyo, Nagano, Shiga)
Centrocestus formosanus (Kagoshima)
Genarchopsis goppo (Ibaraki, Shiga)
Hemiuridae gen. sp. (Tokyo)
Lasiotocus sp. (Aomori)
Lecithochrimum musculus (Mie, unspecified prefecture facing the Seto Inland Sea)
Metagonimus spp. (Shizuoka)
Phyllodistomum anguilae (Aomori, Ibaraki)
Proctotrematoides pisodontophidis (Chiba)
Tubulovesicula anguillae (Miyagi)

Monogenea

- Gyrodactylus egusai* (Shizuoka)
Gyrodactylus joi (Shizuoka)
Gyrodactylus nipponensis (Chiba, Shizuoka, Tokushima, Miyazaki)
Gyrodactylus sp. (Shizuoka)
Pseudodactylogyrus anguillae (Chiba, Aichi, Tokushima)
Pseudodactylogyrus bini (Shizuoka)
Pseudodactylogyrus kamegaii (Chiba)
Pseudodactylogyrus spp. (Shizuoka)

Unidentified Monogenea (Shizuoka, Nagano)

Cestoda

Bothriocephalus claviceps (Shiga)

Bothriocephalus japonicus (Ibaraki, Nagano, Gifu, Shiga)

Bothriocephalus sp. (Nagano)

Nybelinia anguillicola (Mie)

Unidentified Cestoda (Shizuoka)

Nematoda

Anguillicola globiceps (Aomori, Nagano, Chiba, Shizuoka, Aichi, Okayama)

Anguillicoloides crassus (Chiba, Shizuoka, Gifu, Aichi, Mie, Okayama, Tokushima, Oita, Miyazaki, Okinawa)

Cucullanus filiformis (Mie)

Gnathosoma spinigerum (unspecified prefecture in Kyushu)

Heliconema longissimum (-)

Heliconema sp. (Okayama)

Philometroides anguillae (Tokyo, Aichi)

Raphidascaris acus (Shiga)

Acanthocephala

Acanthocephalus gotoi (various localities including Tokyo and Aichi)

Echinorhynchus cotti (Shiga)

Longicollum alemniscus (Aichi)

Pseudorhadinorhynchus samegaiensis (Shiga)

Hirudinida

Batracobdella smaragdina (Aichi, Kagoshima)

Hemiclepsis marginata (Aichi)

Copepoda

Lernaea cyprinacea (Chiba, Shizuoka, Aichi, Mie, Okayama, Hyogo, Shimane, Miyazaki)

Remarks: The above list shows that 42 named species of parasites have been recorded from *A. japonica*. Based on their habitat, these species are categorized into two groups: 34 freshwater (FW) parasites, and 8 marine (M) and/or brackish-water (B) parasites (see the Parasite-Host List). The following six species in the latter group occur as adults: *Lecithochrimum musculus*, *Proctotrematoides pisodontophidis*, *Tubulovesicula anguillae*, *Pseudodactylogyrus kamegaii*, *Cucullanus filiformis*, and *Heliconema longissimum*. This clearly supports the current knowledge on the ecology of *A. japonica* (Tsukamoto et al., 1998; Tsukamoto and Arai, 2001; Arai et al., 2003a, 2003b): the species commonly occurs in marine coastal waters as "sea eels" and "estuarine eels" as well as in freshwater waters as "river eels." Of the six species of parasites of marine and/or brackish-water origin, two species, *Tubulovesicula anguillae* and *Pseudodactylogyrus kamegaii*, are host-specific. It is thus apparent that *A. japonica* serves as the definitive host for the two parasites in marine and/or brackish waters, where the eel constantly remains.

Anguilla marmorata Quoy and Gaimard, 1824

Giant mottled eel, "ō-unagi" (in Japanese)

Cestoda

Bothriocephalus claviceps (Kagoshima)

Anguilla sp.

Monogenea

Pseudodactylogyrus anguillae (Kagoshima)*Pseudodactylogyrus bini* (Aichi, Kagoshima)*Pseudodactylogyrus* sp. (Kagoshima)

Nematoda

Anguillicola globiceps (-)

ACKNOWLEDGEMENTS

We thank Mr. Hideo P. Oka, Port of Nagoya Public Aquarium, Dr. Makoto Iwashita, Japan Fisheries Resources Conservation Association, and Dr. Makoto Tanaka, Shizuoka Prefectural Fisheries Experiment Station, for their help with the literature. Part of the present work was supported by a Grant-in-Aid for Scientific Research (B)(No. 18380116) from the Japan Society for the Promotion of Science.

REFERENCES

- Amin, O., 1985. Classification. In "Biology of Acanthocephala" (ed. by D. W. T. Crompton and B. B. Nickol), Cambridge University Press, Cambridge: 27-72.
- Amin, O. M., Nagasawa, K., Grygier, M. J., 2007. Host and seasonal distribution of fish acanthocephalans from the Lake Biwa basin, Japan. *Comparative Parasitology*, **74**: 244-253.
- Anonymous, 2002. [The parasite fauna of fishes in an experimental river and its characteristics]. *Annual Report of the Aqua Restoration Research Center 2001*: 206-218. (In Japanese).
- Aoyama, J., Watanabe, S., Miyai, T., Sasai, S., Nishida, M., Tsukamoto, K., 2000. The European eel, *Anguilla anguilla* (L.), in Japanese waters. *Dana*, **12**: 1-5.
- Arai, K., Kotake, A., Ohji, M., Miyazaki, N., Tsukamoto, K., 2003a. Migratory history and habitat use of Japanese eel *Anguilla japonica* in the Sanriku Coast of Japan. *Fisheries Science*, **69**: 813-818.
- Arai, K., Kotake, A., Ohji, M., Miller, M. J., Tuskamoto, K., Miyazaki, N., 2003b. Occurrence of sea eels of *Anguilla japonica* along the Sanriku Coast of Japan. *Ichthyological Research*, **50**: 78-81.
- Awakura, T., 1974. Studies on the microsporidian infection in salmonids fishes. *Scientific Reports of the Hokkaido Fish Hatchery*, **29**: 1-95. (In Japanese with English abstract).
- Buchmann, K., Ogawa, K., Lo, C.-F., 1992. Immune response of the Japanese eel (*Anguilla japonica*) against major antigens from the microsporean *Pleistophora anguillarum* Hoshina, 1951. *Fish Pathology*, **27**: 157-161.
- Burreson, E. M. (2006): Phylum Annelida: Hirudinea as vectors and disease agents. In "Fish Diseases and Disorders, Vol. 1. Protozoan and Metazoan Infections. Second Edition" (ed. by P. T. K. Woo), CAB International, Oxfordshire: 566-591.
- Egusa, S., 1967. Gill damage of pond cultured eels. *Fish Pathology*, **1**(2): 72-77. (In Japanese).
- Egusa, S., 1968. Gill damage of pond cultured eels, second series. *Fish Pathology*, **2**: 184-186. (In Japanese).
- Egusa, S., 1970. Branchiophratitis prevailed among eel populations in farm ponds in the winter of 1969-70. *Fish Pathology*, **5**: 51-66. (In Japanese).
- Egusa, S., 1971. [Some problems in recent eel culture]. *Chemistry and Biology*, **9**: 385-389. (In Japanese).
- Egusa, S., 1978. [The Infectious Diseases of Fishes]. Koseisha Koseikaku, Tokyo: 554 p. (In Japanese).

- Egusa, S., 1979. Notes on the culture of the European eel (*Anguilla anguilla* L.) in Japanese eel-farming ponds. *Rapports et Proces-verbaux des Reunions/Conseil Permanent International pour l'exploration de la Mer*, **174**: 51–58.
- Egusa, S., Ahmed, A. T. A., 1970. A suctorian parasite of eels in farm ponds. *Fish Pathology*, **4**: 172–175. (In Japanese with English abstract).
- Egusa, S., Ahmed A. T. A., Kubota, S., 1970. Ichthyophthriasis of elvers imported from France. *Fish Pathology*, **4**: 176–179. (In Japanese with English abstract).
- Egusa, S., Hirose, H., Wakabayashi, H., 1971. A report of investigations on branchionephritis of cultured eels-II. Conditions of the gills and serum ion conditions. *Fish Pathology*, **6**: 57–61. (In Japanese).
- Egusa, S., Kira, K., Wakabayashi, H., 1969. On the occurrence of *Anguillicola globiceps* Yamaguti, a swimbladder roundworm, in pond-cultured eels. *Fish Pathology*, **4**: 52–58. (In Japanese).
- Eiras, J. C., Molnar, K., Lu, Y. S., 2005. Synopsis of the species of *Myxobolus* Bütschli, 1882 (Myxozoa: Myxosporea: Myxobolidae). *Systematic Parasitology*, **61**: 1–46.
- Fujita, T., 1927. Studies on Myxosporidia of Japan. 5. On Myxosporidia in fishes of Lake Biwa. *Journal of the College of Agriculture, Hokkaido Imperial University*, **16**: 229–247, 1 pl.
- Fujita, T., 1929. The skin-disease of the eel. *Annotationes Zoologicae Japonenses*, **12**: 245–250.
- Fukui, T., Morisita, T., 1936. Three new species of Acanthocephala from Japan (a preliminary report). *Dobutsugaku Zasshi (Zoological Magazine)*, **48**: 759–764. (In Japanese with English abstract).
- Fukui, T., Morisita, T., 1937. [Studies on Japanese Acanthocephala]. *Zikken Igaku Zasshi*, **21**: 36–41. (In Japanese).
- Fukui, T., Morisita, T., 1938. Notes on the acanthocephalan fauna of Japan. *Annotationes Zoologicae Japonenses*, **17**: 567–576.
- Gibson, D. I., Bray, R. A., 1986. The Hemiuridae (Digenea) of fishes from the north-east Atlantic. *Bulletin of the British Museum (Natural History), Zoology Series*, **51**: 1–111.
- Golvan, Y. J., 1969. Sytématische des Acanthocéphales (Acanthocephala Rudolphi 1801), L'ordre des Palaeacanthocephala Meyer, 1931, La super-famille des Echinorhynchidea (Cobbald 1876) Golvan et Houin 1973. *Mémoires du Muséum National D'Histoire Naturelle, Série A*, **47**: 5–373.
- Grygier, M. J., Urabe, M., 2003. [Fish parasites introduced into Lake Biwa from overseas]. *Umindo (Quarterly Newsletter of the Lake Biwa Museum)*, **8**(1): 6. (In Japanese).
- Hashimoto, K., Takinami, K., 1976. Electron microscopic observations of the spores of *Plistophora anguillarum*, a microsporidian parasite of the eel. *Bulletin of the Japanese Society of Scientific Fisheries*, **42**: 411–419. (In Japanese with English abstract).
- Hashimoto, K., Sasaki, Y., Takinami, K., 1976. Conditions for extrusion of the polar filament of the spore of *Plistophora anguillarum*, a microsporidian parasite in *Anguilla japonica*. *Bulletin of the Japanese Society of Scientific Fisheries*, **42**: 837–845. (In Japanese with English abstract).
- Hatai, K., Egusa, S., 1973. A note on gill parasites of European eels cultured in ponds. *Fish Pathology*, **8**: 102–105. (In Japanese).
- Hatooka, K., 2002. Anguillidae. In “*Fishes of Japan with Pictorial Keys to the Species, English Edition*” (ed. by T. Nakabo), Tokai University Press, Tokyo: 192.
- Hayward, C. J., 2004. [Fish and parasites—speciation, geographical distribution, migration and dispersion]. In “*Aquaparasitology in the Field in Japan*” (ed. by K. Nagasawa), Tokai University Press, Hadano: 313–325, 344. (In Japanese).

- Hayward, C. J., Iwashita, M., Ogawa, K., Ernst, I., 2001. Global spread of the eel parasite *Gyrodactylus anguillae* (Monogenea). *Biological Invasions*, **3**: 417–424.
- Hine, P. M., 1980. A review of some species of *Myxidium* Bütschli, 1882 (Myxosporea) from eels (*Anguilla* spp.). *Journal of Protozoology*, **27**: 260–267.
- Hirose, H., Sekino, T., Egusa, S., 1976. Notes on the egg deposition, larval migration and intermediate host of the nematode *Anguillicola crassa* parasitic in the swimbladder of eels. *Fish Pathology*, **11**: 27–31. (In Japanese with English abstract).
- Hirose, H., Yabu, T., Hirono, I., Aoki, T., 1998. The phylogeny of *Anguillicola crassus* and *A. globiceps* based on partial 18S ribosomal RNA sequences. *Journal of Fish Diseases*, **21**: 265–271.
- Hoshina, T., 1951a. On a new microsporidian, *Plistophora anguillarum* n. sp., from the muscle of the eel, *Anguilla japonica*. *Journal of Tokyo University of Fisheries*, **38**: 35–46, 2 pls.
- Hoshina, T., 1951b. Zur Entwicklungsgeschichte von, *Proctotrematoides pisodontophidis* Yamaguti, 1938. I. Mitteilung, Agamodistoma und ihre Entwicklung. *Journal of Tokyo University of Fisheries*, **38**: 247–257, 1 pl.
- Hoshina, T., 1952. Notes on some myxosporidian parasites of fishes of Japan. *Journal of Tokyo University of Fisheries*, **39**: 69–89.
- Hoshina, T., 1972. *Plistophora anguillarum* infection found in young eels imported from Formosa. *Fish Pathology*, **6**: 120. (In Japanese).
- Hoshina, T., Sano, T., 1957. On a trypanosome of eel. *Journal of Tokyo University of Fisheries*, **43**: 67–69.
- Imada, R., Muroga, K., 1977. *Pseudodactylogyrus microrchis* (Monogenea) on the gills of cultured eels –I. Seasonal changes in abundance. *Bulletin of the Japanese Society of Scientific Fisheries*, **43**: 1397–1401. (In Japanese with English abstract).
- Imada, R., Muroga, K., 1978. *Pseudodactylogyrus microrchis* (Monogenea) on the gills of cultured eels –II. Oviposition, hatching and development on the host. *Bulletin of the Japanese Society of Scientific Fisheries*, **44**: 571–576. (In Japanese with English abstract).
- Imada, R., Muroga, K., 1979. *Pseudodactylogyrus microrchis* (Monogenea) on the gills of cultured eels –III. Experimental control by trichlorfon. *Bulletin of the Japanese Society of Scientific Fisheries*, **45**: 25–29. (In Japanese with English abstract).
- Imai, S., Miyazaki, H., Nomura, K., 1991. Trichodinid species from the gill of cultured Japanese eel, *Anguilla japonica*, with the description of a new species based on light and scanning electron microscopy. *European Journal of Protistology*, **27**: 79–84.
- Inui, T., Nogami, S., Hirose, H., 1998. Antigenicity of swimbladder Nematoda, *Anguillicola crassus* in Japanese eel, *Anguilla japonica*. *Suisanzoshoku*, **46**: 151–155. (In Japanese with English abstract).
- Inui, T., Ushikoshi, R., Nogami, S., Hirose, H., 1999. A competitive-ELISA for the serodiagnosis of anguillicolosis in Japanese eel, *Anguilla japonica*. *Fish Pathology*, **34**: 25–31. (In Japanese with English abstract).
- Ishii, S., 1915a. [Myxosporidiosis of Japanese eel]. *Dobutsugaku Zasshi (Zoological Magazine)*, **27**: 372–382, 1 pl. (In Japanese).
- Ishii, S., 1915b. [Lentospora-disease of the eel]. *Dobutsugaku Zasshi (Zoological Magazine)*, **27**: 471–474. (In Japanese).
- Ishii, S., 1916a. [On a new species of Nematoda, *Filaria anguillae*, n. sp., from the orbit of Japanese eels]. *Dobutsugaku Zasshi (Zoological Magazine)*, **28**: 214–220. (In Japanese).

- Ishii, S., 1916b. [On a myxosporidian parasitic in the fin of Japanese eel]. *Dobutsugaku Zasshi (Zoological Magazine)*, **28**: 271–273. (In Japanese).
- Ishii, S., 1931. [Parasites of fishes of Japan]. In "Biology, Vol. 18", Iwanami, Tokyo: 179–207. (In Japanese).
- Ito, J., Mochizuki, H., 1968. An epidemiologic study of human helminthes in Shizuoka Prefecture VI. The metacercarial fauna in fresh and brackish water fish. *Japanese Journal of Parasitology*, **17**: 69–74. (In Japanese with English abstract).
- Iwashita, M., Hirata, J., Ogawa, K., 2002. *Pseudodactylogyrus kamegaii* sp. n. (Monogenea: Pseudodactylogyridae) from wild Japanese eel, *Anguilla japonica*. *Parasitology International*, **51**: 337–342.
- Iwata, K., 1972. A case of gill infection of *Myxidium* sp. in wild eels. *Fish Pathology*, **7**: 77–78. (In Japanese).
- Joh, S.-J., Kwon, Y.-K., Kim, M.-C., Kim, M.-J., Kwon, H.-M., Park, J.-W., Kwon, J.-H., Kim, J.-H., 2007. *Heterosporis anguillarum* infections in farm cultured eels (*Anguilla japonica*) in Korea. *Journal of Veterinary Science*, **8**: 147–149.
- Kagei, N., Yanohara, Y., 1995. Epidemiological study on *Centrocestus formosanus* (Nishigori, 1924)—Surveys of its infection in Tanegashima, Kagoshima Prefecture, Japan—. *Japanese Journal of Parasitology*, **44**: 154–160. (In Japanese with English abstract).
- Kano, T., Fukui, H., 1982. Studies on *Plesitophora* infection in eel, *Anguilla japonica*-I. Experimental induction of microsporidiosis and fumagillin efficacy. *Fish Pathology*, **16**: 193–200. (In Japanese with English abstract).
- Kano, T., Okauchi, T., Fukui, H., 1982. Studies on *Plesitophora* infection in eel, *Anguilla japonica*-II. Preliminary tests for application of fumagillin. *Fish Pathology*, **17**: 107–114. (In Japanese with English abstract).
- Kasahara, S., 1957. [On the effect on the control of the anchor worm by Dipterex]. *Noyaku Kenkyu*, **4**(2): 1–6. (In Japanese).
- Kasahara, S., 1958. [A method to control the anchor worm by Dipterex]. *Noyaku Kenkyu*, **5**(2): 26–32. (In Japanese).
- Kasahara, S., 1959. [On the control of the anchor worm]. *Suisanzoshoku*, **6**: 140–148. (In Japanese).
- Kasahara, S., 1962. Studies on the biology of the parasitic copepod *Lernaea cyprinacea* Linnaeus and the methods for controlling this parasite in fish-culture ponds. *Contributions of the Fisheries Laboratory, Faculty of Agriculture, University of Tokyo*, **3**: 103–196. (In Japanese with English abstract).
- Kikuchi, H., 1929. Two new species of Japanese trematodes belonging to Gyrodactylidae. *Annotationes Zoologicae Japonenses*, **12**: 175–186.
- Kikuchi, T., 1929. An investigation into the geographical distribution of *Gnathostoma spinigerum* and an experimental study of its route of infection. *Acta Medica*, **26**: 2943–2970. (In Japanese with English abstract).
- Kuwahara, A., Niimi, A., Itagaki, H., 1974. Studies on a nematode parasitic in the air bladder of the eel I. Description of *Anguillicola crassa* n. sp. (Philometridae, Anguillicolidae). *Japanese Journal of Parasitology*, **23**: 275–279.
- Landsberg, J. H., Lom, J., 1991. Taxonomy of the genera of the *Myxobolus/Myxosoma* group (Myxobolidae: Myxosporea), current listing of species and revision of synonyms. *Systematic*

- Parasitology*, **18**: 165–186.
- Leigh-Sharpe, W. H., 1925. *Lernaea (Lernaeocera) elegans* n. sp., a parasitic copepod of *Anguilla japonica*. *Parasitology*, **17**: 245–251.
- Lom, J., Dyková, I., Körtting, W., Klinger, H., 1989. *Heteosporis schuberti* n. sp., a new microsporidian parasite of aquarium fish. *European Journal of Protistology*, **25**: 129–135.
- Luo, H. Y., Nie, P., Zhang, Y. A., Wang, G. T., Yao, Y. J., 2002. Molecular variation of *Bothriocephalusacheilognathi* Yamaguti, 1934 (Cestoda: Pseudophyllidea) in different fish host species based on ITS rDNA sequences. *Systematic Parasitology*, **52**: 159–166.
- Margolis, L., Arthur, J. R., 1979. Synopsis of the parasites of fishes of Canada. *Bulletin of the Fisheries Research Board of Canada*, **199**: 269 p.
- Matsui, I., 1972. [Science of Eels. Part I: Biological Research, Part II: Culture Techniques]. Koseisha Koseikaku, Tokyo: 737 p. (In Japanese).
- McDonald, T. E., Margolis, L., 1995. Synopsis of the parasites of fishes of Canada: supplement (1978–1993). *Canadian Special Publication of Fisheries and Aquatic Sciences*, **122**: 265 p.
- Mitra, A. K., Bandyopadhyay, P. K., 2005. First record of *Trichodina japonica* Imai, Miyazaki et Nomura 1991 and *Trichodina mutabilis* Kazubski et Migala 1968 (Ciliophora, Trichonidae) from Indian fishes. *Protistology*, **4**: 121–127.
- Miyai, T., Aoyama, J., Sasai, S., Inoue, J. G., Miller, M. J., Tsukamoto, K., 2004. Ecological aspects of the downstream migration of introduced European eels in the Uono River, Japan. *Environmental Biology of Fishes*, **71**: 105–114.
- Moravec, F., 2006. *Drancucunloid and Anguillicoloid Nematodes Parasitic in Vertebrates*. Academia, Prague: 634 p.
- Moravec, F., Taraschewski, H., 1988. Revision of the genus *Anguillicola* Yamaguti, 1935 (Nematoda: Anguillicolidae) of the swimbladder of eels, including descriptions of two new species, *A. novaezelandiae* sp. n. and *A. papernai* sp. n. *Folia Parasitologica*, **35**: 125–146.
- Moravec, F., Nagasawa, K., Miyakawa, M., 2005. First record of ostracods (Ostracoda) as natural intermediate hosts of *Anguillicola crassus* (Nematoda: Anguillicolidae), a pathogenic swimbladder parasite of eels (*Anguilla* spp.). *Diseases of Aquatic Organisms*, **66**: 171–173.
- Moravec, F., Nie, P., Scholz, T., Wang, G., 2003. Some trematodes and cestodes of fishes mainly from Hupei Province, central China. *Acta Societatis Zoologicae Bohemicae*, **67**: 161–174.
- Moravec, F., Taraschewski, H., Anantaphruti, M. T., Maipanich, W., Laoprasert, T., 2007. *Heliconema longissimum* (Ortlepp, 1923) (Nematoda: Physalopteridae) from *Pisodonophis boro* (Teleostei: Ophichthidae) in Thailand, with remarks on the taxonomy of the Prolepidae Schulz, 1927. *Systematic Parasitology*, **66**: 73–80.
- Nagasawa, K., 1993a. Review of human pathogenic parasites in the Japanese common squid (*Todarodes pacificus*). In “Recent Advances in Cephalopod Fisheries Biology” (ed. by T. Okutani, R. K. O’dor and T. Kubodera), Tokai University Press, Tokyo: 293–312.
- Nagasawa, K., 1993b. Parasites of gadid fishes in Japanese waters (review and bibliography). *Scientific Reports of the Hokkaido Fisheries Experimental Station*, **42**: 69–89. (In Japanese with English abstract).
- Nagasawa, K., Miyakawa, M., 2006. Infection of Japanese eel *Anguilla japonica* elvers by *Hemiclepsis marginata* (Hirudinida: Glossiphonidae). *Journal of the Graduate School of Biosphere Science, Hiroshima University*, **45**: 15–19.

- Nagasawa, K., Urawa, S., Awakura, T., 1987. A checklist and bibliography of parasites of salmonids of Japan. *Scientific Reports of the Hokkaido Salmon Hatchery*, **41**: 1–75.
- Nagasawa, K., Awakura, T., Urawa, S., 1989. A checklist and bibliography of parasites of freshwater fishes of Hokkaido. *Scientific Reports of the Hokkaido Fish Hatchery*, **44**: 1–49.
- Nagasawa, K., Kim, Y.-G., Hirose, H., 1994. *Anguillicola crassus* and *A. globiceps* (Nematoda: Dracunculidae) parasitic in the swimbladder of eels (*Anguilla japonica* and *A. anguilla*) in East Asia: a review. *Folia Parasitologica*, **41**: 127–137.
- Nagasawa, K., Umino, T., M. J. Grygier, 2007a. A checklist of the parasites of ayu (*Plecoglossus altivelis altivelis*) (Salmoniformes: Plecoglossidae) in Japan (1912–2007). *Journal of the Graduate School of Biosphere Science, Hiroshima University*, **46**: 59–89.
- Nagasawa, K., Inoue, A., Myat, S., Umino, T., 2007b. New host records for *Lernaea cyprinacea* (Copepoda), a parasite of freshwater fishes, with a checklist of the Lernaeidae in Japan (1915–2007). *Journal of the Graduate School of Biosphere Science, Hiroshima University*, **46**: 21–33.
- Nakai, N., Kokai, E., 1931. On the biological study of a parasitic copepod, *Lernaea elegans* Leigh-Sharpe, infesting on Japanese fresh water fishes. *Journal of the Imperial Fisheries Experimental Station*, **2**: 93–121, 1 pl. (In Japanese with English abstract).
- Naruto Station, Fisheries Experimental Station of Tokushima Prefecture, 1966. Damage from fish diseases and control measures taken in Tokushima Pref. *Fish Pathology*, **1**(1): 62. (In Japanese).
- Nishio, K., Hioki, M., Shiraishi, Y., 1970. Ectoparasitoses occurred in young eels in farm ponds in the winter of 1970. *Fish Pathology*, **5**, 48–50. (In Japanese).
- Nishio, K., Hioki, M., Takeno, N., Shiraishi, Y., Takano, H., Shiraishi, S., Kawamura, E., Toshida, S., Taketani, K., 1971. A report of investigations on branchionephritis of cultured eels—I. Health and environmental conditions of eels and serum ion conditions. *Fish Pathology*, **6**: 47–56. (In Japanese).
- Niwa, M., 1979. [Parasitic diseases]. In “*Culture of European Eel (Anguilla anguilla)*” (ed. by the Eel Culture Research Association), Japan Fisheries Resources Conservation Association, Tokyo: 109–112. (In Japanese).
- Ogawa, K., Egusa, S., 1976. Studies on eel pseudodactylogyrosis—I. Morphology and classification of three eel dactylogyrids with a proposal of a new species, *Pseudodactylogyrus microrchis*. *Bulletin of the Japanese Society of Scientific Fisheries*, **42**: 395–404.
- Ogawa, K., Egusa, S., 1978. Seven species of *Gyrodactylus* (Monogenea: Gyrodactylidae) from *Plecoglossus altivelis* (Plecoglossidae), *Cyprinus carpio* (Cyprinidae) and *Anguilla* spp. (Anguillidae). *Bulletin of the Japanese Society of Scientific Fisheries*, **44**: 613–618.
- Ogawa, K., Egusa, S., 1980. *Gyrodactylus* infestations of cultured eels (*Anguilla japonica* and *A. anguilla*) in Japan. *Fish Pathology*, **15**: 95–99. (In Japanese with English abstract).
- Ogawa, K., Hioki, M., 1986. Two new species of *Gyrodactylus* (Monogenea: Gyrodactylidae) of eel, *Anguilla japonica*, with some data on the occurrence of gyrodactylids in greenhouse culture at Yoshida, Shizuoka Prefecture, Japan. *Fish Pathology*, **21**: 89–94.
- Ogawa, K., Chung, H.-Y., Kou, G.-H., Imada, R., 1985. On the validity of an eel monogenean *Pseudodactylogyrus microrchis* Ogawa et Egusa, 1976. *Bulletin of the Japanese Society of Scientific Fisheries*, **51**: 381–385.
- Ogawa, K., Uno, S., Ito, S., 1985. Infection of cultured eel with *Bartacobdella smaragdina* (Hirudinea: Glossiphoniidae). *Fish Pathology*, **20**: 67–68. (In Japanese).

- Oka, H. P., 1973a. Observations on white spot disease in youngs of the European eel reared in ponds. *Fish Pathology*, **8**: 32–36. (In Japanese with English abstract).
- Oka, H. P., 1973b. A note on the spores of *Myxidium* sp. observed in an “Ich”. *Fish Pathology*, **8**: 37–40. (In Japanese with English abstract).
- Oka, H. P., Egusa, S., 1983. On the *Myxidium* parasitic in fins and the skin of a Japanese eel, *Anguilla japonica*. In “Proceedings of the ROC-Japan Cooperative Scientific Seminar on Fish Diseases” (ed. by G.-H. Kou and S. Egusa), Taipei: 68–72.
- Okamura, A., Zhang, H., Yamada, Y., Utoh, T., Mikawa, N., Horie, N., Tanaka, S., Motonobu, T., 2001. Identification of two eel species, *Anguilla japonica* and *A. anguilla* by discriminant function analysis. *Nippon Suisan Gakkaishi*, **67**: 1056–1060.
- Okamura, A., Yamada, Y., Mikawa, N., Tanaka, A., Oka, H. P., 2002. Exotic silver eels *Anguilla anguilla* in Japanese waters: seaward migration and environmental factors. *Aquatic Living Resources*, **15**: 335–341.
- Ozaki, Y., 1924. [On a new species of trematode of the genus *Azygia*]. *Dobutsugaku Zasshi (Zoological Magazine)*, **36**: 426–435. (In Japanese).
- Ozaki, Y., 1925a. [Preliminary notes on a trematode with anus]. *Dobutsugaku Zasshi (Zoological Magazine)*, **37**: 416–423, 1 pl. (In Japanese).
- Ozaki, Y., 1925b. Preliminary notes on a trematode with anus. *Journal of Parasitology*, **12**: 51–53, 1 pl.
- Rasheed, S., 1963. A revision of the genus *Philometra* Costa, 1845. *Journal of Helminthology*, **37**: 89–130.
- Salati, F., 1987. La parassitosi da *Anguillicola* sp. in Giappone. *Rivista Italiana Piscicoltura e Ittiopatologia*, **22**: 115–117.
- Sato, T., Tanaka, M., 2007. [Control of pseudodactylogyrids in glass eels-II. Occurrence and treatment by temperature change in rearing water]. *Hamana*, **518**: 1–4. (In Japanese).
- Shimazu, T., 1979. Developmental stages of *Azygia gotoi* (Digenea, Azygiidae). *Bulletin of the National Science Museum, Series A (Zoology)*, **5**: 225–234.
- Shimazu, T., 1995. Trematodes of the genus *Genarchoopsis* (Digenea, Derogenidae, Halipeginae) from freshwater fishes of Japan. *Proceedings of the Japanese Society of Systematic Zoology*, **54**: 1–18.
- Shimazu, T., 1997. Cestodes of freshwater earthworms and freshwater fishes in Japan: a review. *Journal of Nagano Prefectural College*, **52**: 9–17. (In Japanese with English abstract).
- Shimazu, T., 1998. Nematodes of freshwater fishes in Japan: a review. *Journal of Nagano Prefectural College*, **53**: 1–19. (In Japanese with English abstract).
- Shimazu, T., 1999a. [Turbellarians and trematodes of freshwater animals in Japan]. In “*Progress of Medical Parasitology in Japan*” [Japanese edition] (ed. by M. Otsuru, S. Kamegai and S. Hayashi), Meguro Parasitological Museum, Tokyo, **6**: 65–86. (In Japanese).
- Shimazu, T., 1999b. Acanthocephalans of freshwater fishes in Japan: a review. *Journal of Nagano Prefectural College*, **54**: 21–29. (In Japanese with English abstract).
- Shimazu, T., 2003. Turbellarians and trematodes of freshwater animals in Japan. In “*Progress of Medical Parasitology in Japan*” [English edition] (ed. by M. Otsuru, S. Kamegai and S. Hayashi), Meguro Parasitological Museum, Tokyo, **7**: 63–86.
- Shimazu, T., 2005. Digeneans found in fresh- and brackish-water fishes of Lake Ogawara in Aomori Prefecture, Japan. *Bulletin of the National Science Museum, Series A (Zoology)*, **31**: 137–150.

- Shimazu, T., Urabe, M., 2005. Digeneans found freshwater fishes of the Uji River at Uji, Kyoto Prefecture, and the Takami River at Higashiyoshino, Nara Prefecture, Japan. *Journal of Nagano Prefectural College*, **60**: 1-14.
- Scholz, T., Škeříková, A., Shimazu, T., Grygier, M. J., 2004. A taxonomic study of species of *Bothrioccephalus* Rudolphi, 1808 (Cestoda: Pseudophyllidea) from eels in Japan: morphological and molecular evidence for the occurrence of *B. claviceps* (Goeze, 1782) and confirmation of the validity of *B. japonicus* Yamaguti, 1934. *Systematic Parasitology*, **57**: 87-96.
- Suh, J.-W., Chun, S.-K., 1988. The infection experiment of *Pleistophora* to eels, *Anguilla japonica* and the histopathological investigation of the infection development. *Bulletin of the Korean Society of Fish Pathology*, **1**: 51-57. (In Korean with English abstract).
- Suyehiro, Y., 1957. [On the ecology of parasites of Japanese eel]. In "Suisangaku Shusei" (ed. By Y. Suyehiro, Y. Oshima and Y. Hiyama), Tokyo University Press, Tokyo: 415-418. (In Japanese).
- Tabeta, O., Takai, T., Matsui, I., 1976. Record of short finned eel from Nagata River, Shimonoseki, Japan. *Bulletin of the Japanese Society for Scientific Fisheries*, **42**: 1333-1338.
- Tabeta, O., Takai, T., Matsui, I., 1977. [On the foreign eel species in Japan]. *Suisanzoshoku*, **24**: 116-122. (In Japanese).
- Tanaka, M., Sato, T., 2007. [Control of pseudodactylogyrids in glass eels]. *Hamana*, **517**: 1-3. (In Japanese).
- Tsai, S.-J., Kou, G.-H., Lo, C.-F., Wang, C.-H., 2002. Complete sequence and structure of ribosomal RNA gene of *Heterosporis anguillarum*. *Diseases of Aquatic Organisms*, **49**: 199-206.
- T'sui, W.-H., Wang, C.-H., 1988. On the *Plistophora* infection in eel. I. Histopathology, ultrastructure, and development of *Plistophora anguillarum* in eel, *Anguilla japonica*. *Bulletin of the Institute of Zoology, Academia Sinica*, **27**: 159-166.
- T'sui, W.-H., Wang, C.-H., Lo, C.-F., 1988. On the *Plistophora* infection in eel. II. The development of *Plistophora anguillarum* in experimentally infected elvers, *Anguilla japonica*. *Bulletin of the Institute of Zoology, Academia Sinica*, **27**: 249-258.
- Tsukamoto, K., Arai, T., 2001. Facultive catadromy of the eel *Anguilla japonica* between freshwater and seawater habitats. *Marine Ecology Progress Series*, **220**: 265-276.
- Tsukamoto, K., Nakai, I., Tesch, F. W., 1998. Do all freshwater eels migrate? *Nature*, **396**: 635-636.
- Tsutsumi, T., 1978. [Treatment of fish diseases at aquaria. Part 7. The anchor worm, a parasite of freshwater fishes, and its control]. *Noyaku Kenkyu*, **12**: 21-22. (In Japanese).
- Umeda, N., Nibe, H., Hara, T., Hirazawa, N., 2006: Effects of various treatments on hatching of eggs and viability of oncomiracidia of the monogenean *Pseudodactylogyrus anguillae* and *Pseudodactylogyrus bini*. *Aquaculture*, **253**: 148-153.
- Ushikoshi, R., Inui, T., Mano, N., Hirose, H., 1999. A method for specific antibody detection from Japanese eel by indirect ELISA using cuticular antigen of *Anguillicola crassus*. *Fish Pathology*, **34**: 81-82.
- Ushiyama, M., Misaki, S., 1977. Seasonal change of physiological conditions in the eel of a typical culture pond. *Bulletin of the Shizuoka Prefectural Fisheries Experimental Station*, **11**: 25-32. (In Japanese).
- Suh, J.-W., Chun, S.-K., 1988. The infection experiment of *Pleistophora* to eels, *Anguilla japonica* and the histopathological investigation of the infection development. *Bulletin of the Korean Society of Fish Pathology*, **1**: 51-57. (In Korean with English abstract).

- Van Cleave, H. J., 1925. Acanthocephala from Japan. *Parasitology*, **17**: 149–156.
- Xu, K., Song, W., Warren, A., 1999. Trichodinid ectoparasites (Ciliophora: Petritrichida) from the gills of cultured marine fishes in China, with the description of *Trichodina lomi* n. sp. *Systematic Parasitology*, **42**: 219–227.
- Xu, K., Song, W., Warren, A., Choi, J. K., 2001. Trichodinid ectoparasites (Ciliophora: Petritrichida) of some marine fishes from coastal regions of the Yellow Sea and Bohai Sea. *Systematic Parasitology*, **50**: 69–79.
- Yamaguti, S., 1934a. Studies on the helminth fauna of Japan. Part 2. Trematodes of fishes, I. *Japanese Journal of Zoology*, **5**: 249–541.
- Yamaguti, S., 1934b. Studies on the helminth fauna of Japan. Part 4. Cestodes of fishes. *Japanese Journal of Zoology*, **6**: 1–112.
- Yamaguti, S., 1935a. Studies on the helminth fauna of Japan. Part 8. Acanthocephala. *Japanese Journal of Zoology*, **6**: 247–278.
- Yamaguti, S., 1935b. Studies on the helminth fauna of Japan. Part 9. Nematodes of fishes, I. *Japanese Journal of Zoology*, **6**: 337–386.
- Yamaguti, S., 1938. Studies on the helminth fauna of Japan. Part 21. Trematodes of fishes, IV. Published by the author: 139 p., 1 pl.
- Yamaguti, S., 1939. Parasitic copepods from fishes of Japan. Part 5. Caligoida, III. *Volumen Jubilare pro Professore Sadao Yoshida*, **2**: 443–487, 33 pls.
- Yamaguti, S., 1941. Studies on the helminth fauna of Japan. Part 33. Nematodes of fishes, II. *Japanese Journal of Zoology*, **9**: 343–396.
- Yamaguti, S., 1963. *Systema Helminthum. Volume V. Acanthocephala*. Interscience Publishers, New York and London: 423 p.
- Yanohara, Y., Kagei, N., 1983. Studies on metacercariae of *Centrocestus formosanus* (Nishigori, 1924)–I. Parasitism of metacercariae in gills of young rearing eels, and abnormal deaths of the hosts. *Fish Pathology*, **17**: 237–241. (In Japanese with English abstract).
- Yoshikawa, M., 2005. [Test of control drugs for pseudodactylogyrids of eels]. *Hamana*, **509**: 1–5. (In Japanese).
- Yoshikawa, M., Tanaka, M., Iinuma, N., Uemura, N., Suzuki, K., 2006. Rearing, and male and female of European eel *Anguilla anguilla*. *Bulletin of the Shizuoka Prefectural Fisheries Experiment Station*, **41**: 63–76. (In Japanese).
- Zang, H., Mikawa, N., Yamada, Y., Horie, N., Okamura, A., Utoh, T., Tanaka, S., Motonobu, T., 1999. Foreign eel species in the natural waters of Japan detected by polymerase chain reaction of mitochondrial cytochrome b region. *Fisheries Science*, **65**: 684–686.

SUPPLEMENTARY REFERENCES

- Egusa, S., ed., 1983. [*Fish Pathology—Infectious and Parasitic Diseases*]. Koseisha Koseikaku, Tokyo: 352 p. (In Japanese).
- Egusa, S., Nakajima, K., 1973. A bibliography of diseases of fishes in Japan. Part 1. Parasitic diseases and parasites. *Fish Pathology*, **7**: 137–229. (In Japanese).
- Fisheries Agency of Japan, ed., 1975. [*A Guide to Diagnosis of Fish Diseases. Carp, Eel, and Yellowtail*]. Japan Fisheries Resources Conservation Association, Tokyo: 159 p. (In Japanese).
- Hatai, K., Ogawa, K., eds., 2006. *New Atlas of Fish Diseases*. Midori Shobo, Tokyo: 295 p. (In Japanese).

- Kobayashi, H., 1935. [Japanese Freshwater Fishes and Their Parasites]. Yokendo, Tokyo: 145 p. (In Japanese).
- Muroga, K., Egusa, S., eds. 1996. [General Fish Pathology]. Koseisha Koseikaku, Tokyo: 177 p. (In Japanese).
- Nagasawa, K., 2002. [Parasitic Animals of Aquatic Organisms]. Seizando Shoten, Tokyo: 184 p. (In Japanese).
- Ogawa, K., 2005. *Parasitology of Fishes in Aquaculture*. University of Tokyo Press, Tokyo: 215 p. (In Japanese).
- Tominaga, O., Yamazaki, T., Nishio, K., Takahashi, K., 1977. [Diseases of Freshwater Fishes: Diagnoses, Causes, and Countermeasures]. Noson-gyoson Bunka Kyokai, Tokyo: 184 p. (In Japanese).
- Wakabayashi, H., Muroga, K., eds., 2004. *Infectious and Parasitic Diseases of Fish and Shellfish*. Koseisha Koseikaku, Tokyo: 424 p. (In Japanese).

日本産ウナギ類の寄生虫目録（1915～2007年）

長澤 和也¹⁾・海野 徹也¹⁾・水野 晃秀²⁾

¹⁾ 広島大学大学院生物圏科学研究所 739-8528 東広島市鏡山1-4-4

²⁾ 愛媛県立宇和島水産高等学校 798-0068 宇和島市明倫町1-2-20

要旨 1915～2007年の93年間に出版された文献に基づき、日本産ウナギ属魚類3種（ウナギ *Anguilla japonica*, オオウナギ *Anguilla marmorata*, ヨーロッパウナギ *Anguilla anguilla*）の寄生虫に関する情報を2つのリスト（寄生虫一宿主リスト、宿主一寄生虫リスト）に整理して目録を作成した。ウナギとオオウナギは在来種であり、ヨーロッパウナギは国外からの移入種である。本目録には、44種の寄生虫（織毛虫類6種、微胞子虫類1種、ミクソゾア類6種、吸虫類7種、単生類7種、条虫類3種、線虫類7種、鉤頭動物4種、ヒル類2種、カイアシ類1種）に加えて、学名がまだ決定していない寄生虫の情報が含まれる。寄生虫一宿主リストでは、各寄生虫は高位分類群ごとに配列され、最新の学名、シノニム、寄生部位、地理的分布および報告者の情報が示されている。上記44種のうち、ウナギから43種、オオウナギから1種、ヨーロッパウナギから10種の寄生虫が報告されている。単生類の *Gyrodactylus anguillae* と線虫類の *Raphidascaris acus* はヨーロッパウナギとともにヨーロッパから、また *Gyrodactylus nipponensis* もインド・西太平洋地域から持ち込まれたと推察されている。ウナギから報告された寄生虫のうち、8種は海産または汽水産で、2種（*Tubulovesicula anguillae*, *Pseudodactylogyrus kamegaii*）の成虫がウナギに特異的に寄生しているため、海洋と汽水域において、ウナギはそれら寄生虫の固有宿主の役割を果たしていると言える。

キーワード：ウナギ；オオウナギ；ヨーロッパウナギ；寄生虫；目録；*Anguilla anguilla*；*Anguilla japonica*；*Anguilla marmorata*