

Remarkable Site Difference of Vibrational Energy Relaxation in Benzene Dimer: Picosecond Time-Resolved IR-UV Pump-Probe Spectroscopic Study**

Ryoji Kusaka and Takayuki Ebata*

Benzene dimer (Bz_2) is one of the most fundamental molecular systems and shows intriguing properties attributed to aromatic-aromatic weak interactions. The structure of Bz_2 in the ground electronic state (S_0) has been extensively studied experimentally^[1] and theoretically.^[2] Several experimental studies on Bz_2 such as the molecular-beam electric deflection^[1(a), 1(b)] and Raman-UV double resonance^[1(h)] and UV-UV hole-burning spectroscopy^[1(g)] strongly suggested that the two benzene (Bz) molecules are not symmetrically equivalent. In addition, recent high level *ab initio* calculations suggested that a floppy T-shaped structure is a global minimum, although there are several possible isomers in nearly isoenergetic region.^[2] The rotational constant obtained from the Fourier-transform microwave spectrum agreed well with the predicted value for the T-shaped form.^[1(i)]

An important subject strongly related to the structure is the vibrational dynamics. In the T-shaped Bz_2 , the two Bz molecules are at different symmetrical conditions; one is at the Stem-site, and the other is at the Top-site. Therefore, the rate constant as well as the process of the vibrational energy relaxation of the two molecules will be different. In this sense, the study of a site-specified vibrational energy relaxation will provide us with a more detailed picture of the structure and vibrational dynamics of Bz_2 . Several studies on the vibrational energy relaxation of Bz_2 have been reported.^[3] Lee and co-workers estimated the lifetime of the CH stretching region to be ~ 3 ps based on the band-width measurement of the IR spectrum.^[3(c)] Felker and co-workers suggested that the ν_2 (CH stretching, Wilson's numbering) level of the Stem-site Bz relaxes faster than that of the Top-site Bz , according to the difference in the band-width of the Raman spectra.^[1(h)] However, in both cases the measured band-width was not that of a single rovibrational level. It is also not clear how the band-width is related to the vibrational energy relaxation dynamics, such as intramolecular (intracluster) vibrational energy redistribution (IVR) and vibrational predissociation (VP). Moreover, there has been no report on the time-resolved spectroscopy for the site-specified Bz_2 .

Here we study the dynamics of the vibrational energy relaxation of the T-shaped Bz_2 in the CH stretching energy region by using picosecond time-resolved IR-UV pump-probe spectroscopy. It is known that for the isolated Bz monomer having a D_{6h} symmetry the IR active CH vibration (ν_{20}) with the e_{1u} symmetry is anharmonically coupled with the $\nu_8 + \nu_{19}$ and $\nu_1 + \nu_6 + \nu_{19}$




Figure 1. Energy level diagram of isotope-substituted benzene dimers [$h^*(\text{Stem})d(\text{Top})$ and $d(\text{Stem})h^*(\text{Top})$] and an excitation scheme. The dimers are vibrationally excited to the CH stretching region by a picosecond IR pulse, and the time evolution is probed by a picosecond UV pulse. The $h^*(\text{Stem})$ and $h^*(\text{Top})$ benzene components (an asterisk represents the vibrationally excited site) show different vibronic structures. Thus, the vibrationally excited dimers can be separately monitored by a tunable picosecond UV pulse. In this figure, VER means vibrational energy relaxation.

combination bands, forming so-called Fermi-triad. All the vibrational modes coupled with the CH stretching are in-plane modes, having stretching or bending character. Their symmetries and frequencies in the isolated Bz molecule are as follow: ν_8 (e_{2g}) = 1600 cm^{-1} , ν_{19} (e_{1u}) = 1484 cm^{-1} , ν_1 (a_{1g}) = 993 cm^{-1} , and ν_6 (e_{2g}) = 608 cm^{-1} .^[4] The Fermi-triad appears at 3048, 3079, and 3101 cm^{-1} in the IR spectrum, and these levels were precisely investigated by Lee and co-workers using IR-UV double resonance spectroscopy.^[4] They also investigated the CH stretching vibrational level for Bz_2 and suggested that the Fermi-triad structure is very similar to that of the monomer, although the vibrational frequencies are slightly shifted from those of the monomer to 3047, 3078, and 3099 cm^{-1} .^[3(c)] Each band contains the transitions of the Bz molecules at different sites. In addition, the Bz molecules at different sites have different symmetries. That is, the Stem-site Bz has a C_{2v} symmetry, and the Top-site Bz has a C_6 symmetry, assuming a nearly free rotational motion of the Stem-site Bz along the C_6 axis of the Top site Bz .^[2(a)]

In the present study, we investigate the vibrational energy relaxation of Bz_2 in order to answer the three fundamental problems; (i) How fast are the IVR and VP in Bz_2 ?, (ii) Are there any site-difference in the rate constants?, and (iii) If it is the case, how is the difference related to the structure? To accomplish the site-specific IR excitation, we use two isotope-substituted heterodimers: (1) $h(\text{Stem})d(\text{Top})$ and (2) $d(\text{Stem})h(\text{Top})$, where $h = C_6H_6$ and $d = C_6D_6$. Figure 1 shows the energy level diagram of the T-shaped dimers and an excitation scheme. A picosecond IR pulse pumps the dimers to one of the CH stretching Fermi-triad. For the hd dimer, the Fermi-triad vibrational frequencies are reported to be 3045, 3077,

[*] JSPS Research Fellow R. Kusaka, Prof. T. Ebata
Department of Chemistry, Graduate School of Science
Hiroshima University
Higashi-Hiroshima 739-8526, Japan
Fax: +81-82-424-7407
E-mail: tebata@hiroshima-u.ac.jp
Homepage: <http://home.hiroshima-u.ac.jp/ebatalab/>

[**] This work is supported by MEXT for the Scientific Research on Priority Area "Molecular Science for Supra Functional Systems" (No. 477). R. K. is supported by JSPS Research Fellowships for Young Scientists.




Figure 2. Transient (1+1) REMPI spectra at several delay times after the IR excitation of the band at 3077 cm^{-1} . Here the $[hd]^+$ mass channel is monitored. The spectra are shown as a function of ν_{UV} (UV frequency, lower axis) and ν'_{vib} (vibrational energy in $S_1 = \nu_{IR} + \nu_{UV} - 0_0^0$, upper axis).

and 3099 cm^{-1} ,^[1(k)] and we chose the band at 3077 cm^{-1} in the present study. After a certain delay time (Δt), a UV pulse probes the population of the vibrationally excited dimers, $h^*(Stem)d(Top)$ and $d(Stem)h^*(Top)$, by (1+1) resonance enhanced multiphoton ionization (REMPI) through the S_1 state. The S_1 - S_0 transition from the level at 3077 cm^{-1} occurs mostly by the $\Delta\nu_1 = -1$ and $\Delta\nu_6 = -1$ transitions because the main character of this level is $\nu_1 + \nu_6 + \nu_{19}$.^[4] The $h^*(Stem)$ exhibits a strong vibronic band while $h^*(Top)$ shows a weak progression in the higher frequency side of the band of $h^*(Stem)$.^[1(c),1(g),1(h)]

Figure 2 shows the mass-resolved transient (1+1) REMPI spectra at several delay times after the IR excitation of hd at 3077 cm^{-1} with the $[hd]^+$ mass channel being monitored. The spectra are shown as a function of ν_{UV} (UV frequency, lower axis) and ν'_{vib} (vibrational energy in $S_1 = \nu_{IR} + \nu_{UV} - 0_0^0$, upper axis). The spectrum at the delay time of 15 ps shows a strong peak at 36390 cm^{-1} and a weak one at 36440 cm^{-1} , and they decay with different time constants. In the present picosecond time-resolved study, the two Bz_2 isomers are simultaneously excited to the vibrational levels $[h^*(Stem)d(Top)$ and $d(Stem)h^*(Top)]$ since their vibrational levels are separated by only 1 cm^{-1} .^[1(k)] So, the transitions of both $h^*(Stem)d(Top)$ and $d(Stem)h^*(Top)$ are overlapped in the REMPI spectrum. However, as described above, they exhibit different vibronic transitions; the Stem-site Bz exhibits a strong vibronic band while the Top-site Bz shows a weak progression with the 15 cm^{-1} interval in the higher frequency side of the strong band. Therefore, the bands at 36390 and 36440 cm^{-1} can be assigned to the transitions from $h^*(Stem)$ and $h^*(Top)$, respectively. Furthermore, since the frequency of the mode 19 (ν'_{19}) of the isolated Bz in S_1 is 1405 cm^{-1} , and the IR-excited bands mainly involve the $\nu_1 + \nu_6 + \nu_{19}$ character,^[4] the transitions at 36390 and 36440 cm^{-1} can be assigned to $19^1 \leftarrow h^*(Stem)$ and $19^1 X^n \leftarrow h^*(Top)$ resonance transitions, respectively, where X represents an intermolecular vibration. A very weak broad background in the 36250 - 36350 cm^{-1} region is due to larger clusters, which generate the $[hd]^+$ fragment ion after the photoionization.^[1(c),1(d),1(f),5] As seen in the figure, the contribution of the larger clusters is very small in the present condition.

The black and red curves in Figure 3(a) show the time profiles of the pump-probe signals obtained by fixing the UV frequencies to the bands at 36390 and 36440 cm^{-1} , respectively. This is the first real-





Figure 3. Picosecond IR-UV pump-probe signals of the vibronic bands at a) 36390 cm^{-1} (black) and 36440 cm^{-1} (red) and the broad band at b) 37600 cm^{-1} (blue) after the IR excitation of the band at 3077 cm^{-1} .

time observation of the site-specified vibrational energy relaxation of Bz_2 . It is clear that the intensity of the band at 36390 cm^{-1} (black curve) decays much faster than that of the band at 36440 cm^{-1} (red curve). The decay curves can be fitted by single exponential decay function

$$I(t) = I_0 \exp(-\Delta t / \tau), \quad (1)$$

convoluted with the 12 ps widths of the laser pulses. One point to be mentioned in Fig. 3(a) is that the two curves do not completely reach to zero at long delay time because of the unavoidable fragmentation from the larger clusters after the ionization. Therefore, we fitted the decay curves by including the nonzero baseline as a step function. Finally, we obtain the decay lifetimes $\tau_{Stem} = 110 \text{ ps}$ ($k_{Stem} = 9.0 \times 10^9 \text{ s}^{-1}$) and $\tau_{Top} = 500 \text{ ps}$ ($k_{Top} = 2.0 \times 10^9 \text{ s}^{-1}$) for $h^*(Stem)d(Top)$ and $d(Stem)h^*(Top)$, respectively. Thus, the lifetime of the Stem-site Bz is 4.5 times shorter than that of the Top-site Bz, in spite of the fact that their vibrational frequencies differ by only 1 cm^{-1} . It was reported that the isolated Bz does not show vibrational energy relaxation in this energy region,^[4,8(a)] which may be because of its high symmetry (D_{6h}). In the dimer, on the other hand, the two Bz molecules are at lower symmetrical environments; the Stem- and Top-site benzene molecules belong to C_{2v} and C_6 symmetry, respectively. The lowering of the symmetry may cause the vibrational energy relaxation in the T-shaped Bz_2 . Moreover, the present study indicates that “the initial level \leftrightarrow dark state” coupling strength of the Stem-site Bz is 4.5 larger than that of the Top-site Bz. The larger coupling of the Stem-site Bz is partly explainable by the nature of the CH stretching mode and the structure of Bz_2 . In the Stem-site Bz, the CH stretching motion is parallel to the intermolecular stretching motion, hence an efficient “CH \leftrightarrow intermolecular modes” coupling may occur through the CH $\cdots\pi$ interaction. In the Top-site Bz, on the other hand, the coupling of the CH stretching motion with the intermolecular modes is negligibly small. A theoretical treatment is strongly demanded to support this hypothesis.

Another important issue remains; “What dynamics is the observed decay attributed to?” Since the binding energy of Bz_2 is reported to be 500 - 800 cm^{-1} ,^[6] and the input energy of 3077 cm^{-1} is much larger than the binding energy, the vibrationally excited Bz_2 dissociates finally. Two dissociation pathways have to be considered,





In scheme (2a), the dimer sequentially predissociates through IVR while in scheme (2b) the dimer directly predissociates from the initial level. One way to answer the above question is to observe the electronic transitions from the redistributed levels (hd_{bath}). Actually, we observed a broad transition at $\sim 37600 \text{ cm}^{-1}$, which is assignable to the hot band transitions from the bath modes (the UV spectrum is not shown here). Figure 3(b) shows the IR-UV pump-probe time profile obtained by fixing the UV frequency to 37600 cm^{-1} . The intensity of the broad transition becomes maximum at $\Delta t = 200\text{-}300 \text{ ps}$ and decays with the time scale of $\sim 1 \text{ ns}$. The time reaching to the maximum, $\Delta t = 200\text{-}300 \text{ ps}$, is close to the time scale of the disappearance of the Stem-site Bz signal, meaning that the rise corresponds to IVR, and the decay with the long time scale is attributed to VP. Thus, we can say that at least scheme (2a) occurs. At present, however, we cannot conclude that the sequential process (2a) occurs solely. Zewail and his group investigated the vibrational energy relaxation processes in detail for the van der Waals complexes of stilbene with Helium, Neon, and Argon in S_1 by picosecond time-resolved photofragment spectroscopy. They concluded that IVR precedes VP and the complexes dissociate through the sequential process.^[7] A similar sequential process was also reported for the dissociation from the excited OH stretching vibration of hydrogen-bonded clusters of phenol.^[8(c)] For further confirmation, the detection of the Bz fragment and its time profile measurement are necessary, and these issues are to be addressed in our near-future work. We also performed the experiments for the other bands forming Fermi-triad (3045 and 3099 cm^{-1}) and found that they show the very similar relaxation rate constants. This result is understandable because this Fermi-triad is composed of the same set of the vibrations.

In conclusion, we investigated the site-specified vibrational energy relaxation of the benzene dimer by picosecond time-resolved IR-UV pump-probe spectroscopy. By using isotope-substituted species, the IR excitation of the site-specified benzene components and time-resolved probe of the vibrational energy relaxation have become possible for the first time. It was found that the CH stretching vibrational level of the Stem-site benzene relaxes 4.5 times faster than that of the Top-site benzene, although their vibrational energies differ by only 1 cm^{-1} . The result indicates an importance of the $\text{CH}\cdots\pi$ interaction not only to the structure of the aromatic systems but also to the vibrational energy relaxation dynamics, which will trigger numerous theoretical investigations. The studies of the vibrational energy relaxation of homodimer (hh) and that starting from the CD stretching vibration of hd and dd are now in progress.

Experimental Section

The experimental setup for the picosecond time-resolved IR-UV pump-probe spectroscopy is the upgraded version of that described in our previous papers.^[6] Briefly, a tunable picosecond IR pulse is obtained by difference frequency generation (DFG) between the signal and the idler output of an optical parametric generation/optical parametric amplifier (OPG/OPA) system (Ekspra PG401/DFG2-10P) pumped by a mode-locked picosecond Nd:YAG laser (Ekspra PL2143S). A tunable picosecond UV pulse is obtained by second harmonic generation (SHG) of the output of an OPG/OPA (Ekspra PG401SH) pumped by the same Nd:YAG laser. The spectral

resolutions of the IR and UV lights are 5 cm^{-1} , and the time resolution of the two pulses is 12 ps . The delay time between IR and UV pulses is changed by an optical delay line. For generating hd heterodimers, a gaseous mixture of h and d [50:50(v/v)] diluted with helium carrier gas at a total pressure of 3 bar is expanded into vacuum by a pulsed valve (General valve, Series 9). The molecular beam is obtained by skimming the center of the expansion. To reduce the formation of the clusters larger than the dimer, the container of the sample is maintained at $-20 \text{ }^\circ\text{C}$, and the width and timing of the pulse valve are carefully controlled. The IR and UV laser lights cross the molecular beam, and the photoionized benzene clusters are mass-analyzed by a time-of-flight mass spectrometer and detected by a channeltron (Burle 4900).

Received: ((will be filled in by the editorial staff))

Published online on ((will be filled in by the editorial staff))

Keywords: benzene dimer · IVR · laser chemistry · time-resolved spectroscopy · vibrational energy relaxation

- [1] a) K. C. Janda, J. C. Hemminger, J. S. Winn, S. E. Novick, S. J. Harris, W. Klemperer, *J. Chem. Phys.* **1975**, *63*, 1419; b) J. M. Steed, T. A. Dixon, W. Klemperer, *J. Chem. Phys.* **1979**, *70*, 4940; c) J. B. Hopkins, D. E. Powers, R. E. Smalley, *J. Phys. Chem.* **1981**, *85*, 3739; d) P. R. R. Langridge-Smith, D. V. Brumbaugh, C. A. Haynam, D. H. Levy, *J. Phys. Chem.* **1981**, *85*, 3742; e) K. O. Börnsen, H. L. Selzle, E. W. Schlag, *J. Chem. Phys.* **1986**, *85*, 1726; f) A. Kiermeier, B. Ernstberger, H. J. Neusser, E. W. Schlag, *Z. Phys. D –Atoms, Molecules and Clusters* **1988**, *10*, 311; g) W. Scherzer, O. Krätzschmar, H. L. Selzle, E. W. Schlag, *Z. Naturforsch., A: Phys. Sci.* **1992**, *47*, 1248; h) B. F. Henson, G. V. Hartland, V. A. Venturo, P. M. Felker, *J. Chem. Phys.* **1992**, *97*, 2189; i) E. Arunan, H. S. Gutowsky, *J. Chem. Phys.* **1993**, *98*, 4294; j) V. A. Venturo, P. M. Felker, *J. Chem. Phys.* **1993**, *99*, 748; k) U. Erlekam, M. Frankowski, G. Meijer, G. von Helden, *J. Chem. Phys.* **2006**, *124*, 171101.
- [2] a) P. Hobza, H. L. Selzle, E. W. Schlag, *J. Chem. Phys.* **1990**, *93*, 5893; b) P. Hobza, H. L. Selzle, E. W. Schlag, *J. Phys. Chem.* **1996**, *100*, 18790; c) V. Špirko, O. Engkvist, P. Soldán, H. L. Selzle, E. W. Schlag, P. Hobza, *J. Chem. Phys.* **1999**, *111*, 572; d) S. Tsuzuki, T. Uchmaru, K. Matsumura, M. Mikami, K. Tanabe, *Chem. Phys. Lett.* **2000**, *319*, 547; e) W. Wang, M. Pitonák, P. Hobza, *ChemPhysChem* **2007**, *8*, 2107; f) E. C. Lee, D. Kim, P. Jurecka, P. Tarakeshwar, P. Hobza, K. S. Kim, *J. Phys. Chem. A* **2007**, *111*, 3446.
- [3] a) M. F. Vernon, J. M. Lisy, H. S. Kwok, D. J. Krajnovich, A. Tramer, Y. R. Shen, Y. T. Lee, *J. Phys. Chem.* **1981**, *85*, 3327; b) G. Fischer, *Chem. Phys. Lett.* **1987**, *139*, 316; c) R. H. Page, Y. R. Shen, Y. T. Lee, *J. Chem. Phys.* **1988**, *88*, 4621.
- [4] R. H. Page, Y. R. Shen, Y. T. Lee, *J. Chem. Phys.* **1988**, *88*, 5362.
- [5] a) T. Iimori, Y. Ohshima, *J. Chem. Phys.* **2002**, *117*, 3656; b) T. Iimori, Y. Aoki, Y. Ohshima, *J. Chem. Phys.* **2002**, *117*, 3675.
- [6] a) J. R. Grover, E. A. Walters, E. T. Hui, *J. Phys. Chem.* **1987**, *91*, 3233; b) A. Kiermeier, B. Ernstberger, H. J. Neusser, E. W. Schlag, *J. Phys. Chem.* **1988**, *92*, 3785; c) H. Krause, B. Ernstberger, H. Neusser, *J. Chem. Phys. Lett.* **1991**, *184*, 411.
- [7] a) D. H. Semmes, J. S. Baskin, A. H. Zewail, *J. Am. Chem. Soc.* **1987**, *109*, 4104; b) D. H. Semmes, J. S. Baskin, A. H. Zewail, *J. Chem. Phys.* **1990**, *92*, 3359.
- [8] a) T. Ebata, M. Kayano, S. Sato, N. Mikami, *J. Phys. Chem. A* **2001**, *105*, 8623; b) Y. Yamada, J. Okano, N. Mikami, T. Ebata, *Chem. Phys. Lett.* **2006**, *432*, 421; c) Y. Yamada, Y. Katsumoto, T. Ebata, *Phys. Chem. Chem. Phys.* **2007**, *9*, 1170; d) Y. Yamada, N. Mikami, T. Ebata, *Proc. Natl. Acad. Sci.* **2008**, *105*, 12690.