

# Asymmetric nature of two subunits of RAD18, a RING-type ubiquitin ligase E3, in the human RAD6A–RAD18 ternary complex

Yuji Masuda<sup>1,\*</sup>, Miki Suzuki<sup>1</sup>, Hidehiko Kawai<sup>2</sup>, Fumio Suzuki<sup>2,3</sup> and Kenji Kamiya<sup>1</sup>

<sup>1</sup>Department of Experimental Oncology, <sup>2</sup>Department of Molecular Radiobiology and <sup>3</sup>Department of International Radiation Emergency Medicine, Research Institute for Radiation Biology and Medicine, Hiroshima University, 1-2-3 Kasumi, Minami-ku, Hiroshima 734-8553, Japan

Received May 31, 2011; Revised September 7, 2011; Accepted September 13, 2011

## ABSTRACT

**RAD18, a RING-type ubiquitin ligase (E3) that plays an essential role in post-replication repair, possesses distinct domains named RING, UBZ, SAP and the RAD6-binding domain (R6BD) and forms a dimer. RAD6, an ubiquitin-conjugating enzyme (E2), stably associates with R6BD in the C-terminal portion. In this study, we established a method to distinguish between the two subunits of RAD18 by introduction of different tags, and analyzed mutant complexes. Our results, surprisingly, demonstrate that RAD6A and RAD18 form a ternary complex, RAD6A–(RAD18)<sub>2</sub> and the presence of only one R6BD in the two RAD18 subunits is sufficient for ternary complex formation and the ligase activity. Interestingly, ligase activity of a mutant dimer lacking both R6BDs is not restored even with large amounts of RAD6A added in solution, suggesting a requirement for precise juxtaposition via interaction with R6BD. We further show that mutations in both subunits of either RING or SAP, but not UBZ, strongly reduce ligase activity, although inactivation in only one of two subunits is without effect. These results suggest an asymmetric nature of the two RAD18 subunits in the complex.**

## INTRODUCTION

Ubiquitin ligases (E3s) catalyze the transfer of ubiquitin from E2 (ubiquitin-conjugating enzyme)–ubiquitin conjugates to lysine residues in target proteins. A subset of E3s contains a RING (*really interesting new gene*) domain, which binds to E2–ubiquitin conjugates and seems to activate thioester bonds (1,2). Some RING-type E3s are known to form heterodimers such as BRCA1–BARD1

(3–5), Ring1b–Bmi1 (6,7) and MDM2–MDMX (8–10), while others like cIAP2 (11) and RNF4 (12) act as homodimers. The heterodimeric RING-type E3s are composed of active and inactive RING domains, and dimerization enhances the ligase activity, suggesting that the pairing itself is very important for enzyme function (5,6,9).

Through a RAD6-binding domain (R6BD) located in its C-terminal region, the RAD18 RING-type ubiquitin ligase forms a stable complex with a specific E2, RAD6 (13–17). Since RAD6 also contacts the RING domain near the N-terminal of RAD18 for catalytic function (2,15,18), it could interact with two distinct domains of RAD18 simultaneously. Such interactions between an E2 and an E3 are quite unique to RAD6–RAD18. Recently, it has been reported that interaction between R6BD and RAD6 blocks the intrinsic activity of RAD6 in forming ubiquitin chains, ensuring mono-ubiquitination of PCNA (19). Previous studies have also reported that RAD18 forms a dimer and suggested the RAD6–RAD18 complex to be a dimer of RAD6–RAD18 heterodimer (15,20,21), hereafter designated as (RAD6–RAD18)<sub>2</sub>.

RAD6 and RAD18 play an essential role in post-replication repair of damaged DNA *via* ubiquitination of proliferating cell nuclear antigen (PCNA) at Lys164 (22,23). The RAD6–RAD18 complex itself catalyzes mono-ubiquitination of PCNA *in vitro* (24–28). The mono-ubiquitinated PCNA appears to enhance lesion bypass replication by stimulation of entry of translesion DNA polymerases at stalled 3'-ends, through interactions between ubiquitin-binding domains of the polymerases and ubiquitin moieties of mono-ubiquitinated PCNA (24,26,29–31). Furthermore, RAD18 features a pol η-binding domain that is important for recruiting pol η to stalled 3'-ends (17,32).

RAD18 has two other domains UBZ (ubiquitin-binding zinc finger) (33–35) and SAP (SAF-A/B, Acinus and

\*To whom correspondence should be addressed. Tel: +81 82 257 5893; Fax: +81 82 257 5843; Email: masudayu@hiroshima-u.ac.jp

PIAS) (15,36–38). The former is required for accumulation of RAD18 at damage sites (34,37,39) and has affinity for Ub chains (33,34) suggesting specific binding to damage-associated poly-ubiquitinated proteins (39). The SAP domain possesses DNA-binding activity (15,38), although it appears to be unnecessary for accumulation of RAD18 at damage sites (37,39). Rather, it is crucial for pol  $\eta$  focus formation (37) possibly depending on PCNA ubiquitination, which is attributed to an essential role of the SAP domain in ligase activity (38,39).

Although multiple domains of RAD18 are clearly concerned with ligase activity, it is unknown how the distinct entities of the two RAD18 subunits interact with each other for enzyme function. In the present study, we established a method to analyze the structure and functions of the human RAD6A–RAD18 complex and demonstrated an asymmetric nature of the two RAD18 molecules in the complex.

## MATERIALS AND METHODS

### Plasmids

Expression plasmids for RAD6A–RAD18, E1, ubiquitin, PCNA and RFC were as described previously (26,40–42). For overproduction of human RAD6A, RAD6A–<sup>His</sup>RAD18, <sup>FLAG</sup>RAD6A–<sup>His</sup>RAD18, the genes were cloned into pET20b(+) (Novagen) to yield pET-RAD6A and pET-RAD6A/hisRAD18, pET-flagRAD6A/hisRAD18, respectively. To make an expression plasmid compatible with pET plasmids in *Escherichia coli* cells, the entire coding unit of pET-RAD6A/RAD18 (42) was cloned into pACYC Duet1 (Novagen) to yield pAC-RAD6A/RAD18. Expression plasmids for overproduction of RAD6A–<sup>FLAG</sup>RAD18 and <sup>FLAG</sup>RAD6A–<sup>FLAG</sup>RAD18, pAC-RAD6A/flagRAD18 and pAC-flagRAD6A/flagRAD18 were generated, respectively. In those plasmids, His-tagged sequences were taken from pET15 and all tagged sequences were attached to immediately before the start codons of the respective genes. Expression plasmids in human cells for *RAD6* were cloned in pCMV, and for *RAD18* in pCDNA3 flag and pCAGGS (43).

### Proteins

Proteins used in this study were overproduced in *E. coli* cells. During all purification steps, monitoring was done by SDS–PAGE followed by staining with Coomassie Brilliant Blue R-250 (CBB), or western blotting. Protein concentrations were determined by Bio-Rad protein assay using BSA (Bio-Rad) as the standard. PCNA, RFC, E1, ubiquitin and RAD6A–RAD18 complex were purified as described previously (26,40,41). Detailed procedures for purification of recombinant proteins established in this study are described in Supplementary Data.

### Sucrose density gradient sedimentation

Sucrose density gradient sedimentation was performed as described earlier (44). Purified RAD6A–RAD18 complexes (1.7  $\mu$ g), were sedimented through 2 ml of

10–40% sucrose gradient in buffer A containing 300 mM NaCl by centrifugation at 55 000 rpm for 20 h in a TLS 55 rotor (Beckman) at 4°C and fractions (100  $\mu$ l) were collected from the bottom of the tube and analyzed by SDS–PAGE. Gel bands were stained with CBB and quantified using Multi Gauge software Version 3.0 (FUJIFILM). Sedimentation coefficients were determined relative to those of standard proteins sedimented in parallel gradients.

### Antibodies

To obtain polyclonal antibodies against RAD18, truncated His-tagged RAD18 proteins (127–255 amino acids) were expressed in BL21 (DE3) (45), purified and used to immunize rabbits. Anti-Penta-His monoclonal (Qiagen, 34660), anti-FLAG M2 monoclonal (Sigma, F3165), anti-RAD6 polyclonal (Abcam, ab31917) and anti-PCNA polyclonal (Santa Cruz, sc-7907) antibodies were purchased.

### PCNA-mono-ubiquitination assays

The standard reaction mixture (25  $\mu$ l) contained 20 mM HEPES–NaOH (pH 7.5), 50 mM NaCl, 0.2 mg/ml BSA, 1 mM DTT, 10 mM MgCl<sub>2</sub>, 1 mM ATP, 100 ng poly(dA)-oligo(dT), PCNA (1 pmol), RFC (350 fmol), E1 (850 fmol), ubiquitin (170 pmol) and the indicated amounts of RAD6A–RAD18 complex. Reaction mixtures were prepared on ice then incubated at 30°C for 10 min. The reactions were terminated by addition of 2 $\times$  SDS sample buffer containing 25 mM EDTA. Ubiquitination of PCNA was assessed by western analysis, detected by an ECL chemiluminescence kit (GE Healthcare Life Science).

## RESULTS

### Physicochemical properties of the human RAD6A–RAD18 complex

To study the subunit composition of RAD6A–RAD18 complex, we first determined the Stokes' radius and sedimentation coefficient of the purified RAD6A–RAD18 complex (26) by gel filtration and sucrose density gradient centrifugation, respectively (Table 1). The obtained value of Stokes' radius (62 Å) corresponds to an apparent molecular mass of 490 kDa, but that of the

**Table 1.** Physicochemical properties of the RAD6A–RAD18 complex

Stokes' radius <sup>a</sup> (Å)	Sedimentation coefficient <sup>b</sup> $\times 10^{-13}$ s ( $S_{20,w}$ )	Molecular mass <sup>c</sup> (kDa)
62	5.0	131

<sup>a</sup>Determined by Superdex 200 gel filtration using the size markers ferritin (61.0 Å), aldolase (48.1 Å), albumin (35.5 Å) ovalbumin (30.5 Å) and ribonuclease A (16.4 Å), and the data were based on  $A_{280}$  values monitored during the chromatography.

<sup>b</sup>Determined with ferritin (17.6 S), catalase (11.3 S), aldolase (7.4 S), albumin (4.2 S) and ribonuclease A (1.8 S) as standards, and the data were based on the SDS–PAGE gel profile.

<sup>c</sup>Estimated from the Stokes' radius and the sedimentation coefficient assuming a partial specific volume of 0.73 (46).

sedimentation coefficient (5.0S) corresponds to an apparent molecular mass of 70 kDa. The large difference between these two values suggests that the complex does not have a compact globular shape. Employing the method described by Siegel and Monty (46), we estimated the molecular mass of the RAD6A–RAD18 complex to be 131 kDa (Table 1). Since calculated molecular masses of RAD6A and RAD18 are 17.3 and 56.2 kDa, respectively, the total of molecular mass should be 73.5 kDa for RAD6A–RAD18 and 147 kDa for (RAD6A–RAD18)<sub>2</sub>. Although the value estimated from the experiments was close to that of (RAD6A–RAD18)<sub>2</sub>, we further investigated the stoichiometry of RAD6A and RAD18 in the complex.

### Direct evidence that the RAD6A–RAD18 complex contains two RAD18 molecules


To obtain direct evidence that the RAD6A–RAD18 complex contains two molecules of RAD18, we co-expressed both His-tagged *RAD18* and FLAG-tagged *RAD18* genes together with *RAD6A* gene in the same *E. coli* cells. Three different complexes containing His<sup>6</sup>RAD18–His<sup>6</sup>RAD18, FLAG<sup>3</sup>RAD18–FLAG<sup>3</sup>RAD18 or His<sup>6</sup>RAD18–FLAG<sup>3</sup>RAD18 would be expected if we assume that the complex should contain two molecules of RAD18. When the cell lysate was loaded to a Ni-chelating column, we found some FLAG<sup>3</sup>RAD18 to be absorbed to the column and eluted together with approximately equal amounts of His<sup>6</sup>RAD18 at a lower imidazole concentration (Figure 1A, lane 2), whereas the remainder of the His<sup>6</sup>RAD18 was eluted at a higher imidazole concentration (see Supplementary Materials and Methods section). Since it was expected that the former was His<sup>6</sup>RAD18–FLAG<sup>3</sup>RAD18 hetero complex and the latter was His<sup>6</sup>RAD18 homo complex, the former was further purified through a heparin column (Figure 1A, lane 3) and analyzed by gel filtration (Figure 1A, lane 4). The elution profile was the same as that of untagged

RAD6A–RAD18 complex (Table 1 and see also Figure 6C), suggesting the tag sequences do not affect the overall structure of the complex. While western blotting with anti-Penta-His and anti-FLAG antibodies specifically detected His<sup>6</sup>RAD18 and FLAG<sup>3</sup>RAD18 proteins, respectively (Figure 1B), blotting with anti-RAD18 antibodies detected the two proteins equally, thereby indicating that the purified complex contains His<sup>6</sup>RAD18 and FLAG<sup>3</sup>RAD18 at a 1:1 ratio. To further verify that RAD6A, His<sup>6</sup>RAD18 and FLAG<sup>3</sup>RAD18 form a complex, a fraction eluted from gel filtration was applied to FLAG-affinity chromatography. The result demonstrated that the three proteins were adsorbed to the anti-FLAG M2 affinity gel and eluted with the FLAG peptide (Figure 1A, lane 5). We thus conclude that the RAD6A–RAD18 complex contains two RAD18 molecules.

### Subunit composition of the RAD6A–RAD18 complex is RAD6A–(RAD18)<sub>2</sub>


To examine the stoichiometry of RAD6A in the complex, we directly compared amounts of RAD6A in the complex with purified RAD6A protein as a reference. We applied 500 ng of the purified complex, RAD6A–His<sup>6</sup>RAD18–FLAG<sup>3</sup>RAD18, in parallel with different amounts (65 or 115 ng) of RAD6A monomers to SDS–PAGE. The amounts of RAD6A in the complex should be 65 or 115 ng, if the complex is RAD6A–(RAD18)<sub>2</sub> or (RAD6A–RAD18)<sub>2</sub>, respectively (Figure 2A). The results of CBB staining and western blotting showed the amount of RAD6A in the complex to be closer to 65 ng (Figure 2A), thus suggesting that the complex contains RAD6A and RAD18 at the ratio of 1:2. To confirm this, the FLAG-tag was introduced to RAD6A and the FLAG<sup>3</sup>RAD6A–His<sup>6</sup>RAD18–FLAG<sup>3</sup>RAD18 complex was purified as described above. We applied 505 ng of the purified complex to SDS–PAGE, in parallel with 500 ng of RAD6A–His<sup>6</sup>RAD18–FLAG<sup>3</sup>RAD18 as a reference (Figure 2B). Western blotting with anti-RAD6 antibodies confirmed that both complexes contained equivalent amounts of RAD6A (Figure 2B), and blotting with an anti-FLAG antibody clearly demonstrated that the molecular ratio of FLAG<sup>3</sup>RAD6A and FLAG<sup>3</sup>RAD18 was 1:1 (Figure 2B), suggesting strongly the ratio of RAD6A to RAD18 to be 1:2.

R6BD is located within amino acid residues 340–395 of RAD18 (Figure 3D) (17,19). Using a deletion mutant consisting of amino acid residues 1–341 of RAD18 (hereafter designated as RAD18<sup>ΔC1</sup>, Figure 3D), we further examined the stoichiometry of RAD6 and RAD18. When His<sup>6</sup>RAD18<sup>ΔC1</sup> and FLAG<sup>3</sup>RAD18<sup>ΔC1</sup> were co-produced with RAD6A in the same *E. coli* cells, a complex containing His<sup>6</sup>RAD18<sup>ΔC1</sup> and FLAG<sup>3</sup>RAD18<sup>ΔC1</sup> was purified similarly as described above, but it did not contain RAD6A (Figure 2C, lane 3). This result indicates that the R6BD is required for complex formation with RAD6 but is dispensable for dimerization. In contrast, when His<sup>6</sup>RAD18<sup>ΔC1</sup> was co-produced with FLAG<sup>3</sup>RAD18 and RAD6A in the same *E. coli* cells, we successfully obtained a RAD6A–His<sup>6</sup>RAD18<sup>ΔC1</sup>–FLAG<sup>3</sup>RAD18 ternary complex



**Figure 1.** Purification of the RAD6A–His<sup>6</sup>RAD18–FLAG<sup>3</sup>RAD18 complex. (A) Pooled fractions eluted from respective columns were analyzed by SDS–PAGE followed by staining with CBB. Lane 1, cell lysate; lane 2, Ni-chelating column; lane 3, heparin column; lane 4, gel-filtration column; lane 5, anti-FLAG affinity column. Molecular masses of each marker (lane M) are shown to the left of the gel. (B) Western analysis of the pooled fraction eluted from the gel filtration column. Membranes were probed with the indicated antibodies.






**Figure 2.** Subunit composition of the RAD6A-RAD18 complex and deletion analysis of RAD18. (A) The indicated amounts of purified RAD6A-RAD18 complex and RAD6A were analyzed by SDS-PAGE followed by staining with CBB and western blotting probed with anti-RAD6 antibodies. Relative chemiluminescence signals detected with a CCD camera are shown. Amounts of each subunit calculated for two different postulated subunit compositions are shown in the table. (B and C) Indicated amounts of purified RAD6A-RAD18 complexes were analyzed by SDS-PAGE followed by staining with CBB and western blotting probed with anti-FLAG and anti-RAD6 antibodies. Relative chemiluminescence signals detected with a CCD camera are shown. Amounts of each subunit calculated for the postulated subunit compositions are shown in the tables. (D) Schematic representations of the structures of the complexes for respective RAD6A-RAD18 complexes. (E) Ligase activities of the respective RAD6A-RAD18 complexes. Increasing amounts of the complexes (0.5, 1 and 2 pmol) were subjected to standard assays. The reaction products were analyzed by western blotting with anti-PCNA antibodies. I, RAD6A-(HisRAD18-FLAGRAD18); II, RAD6A-(HisRAD18<sup>ΔC1</sup>-FLAGRAD18); III, HisRAD18<sup>ΔC1</sup>-FLAGRAD18<sup>ΔC1</sup>. (F) Titration of RAD6A in the reaction with HisRAD18<sup>ΔC1</sup>-FLAGRAD18<sup>ΔC1</sup>. Indicated amounts of RAD6A were incubated with 2 pmol of HisRAD18<sup>ΔC1</sup>-FLAGRAD18<sup>ΔC1</sup> (III) under standard assay conditions. As control reactions, 1 pmol of RAD6A-(HisRAD18-FLAGRAD18) (I) was incubated in the presence or absence of additional RAD6A. (G) Analysis of complex formation and ligase activity of another C-terminal deletion mutant of RAD18. Purified complexes (3.7 pmol as a trimer) were analyzed by SDS-PAGE followed by staining with CBB. Lane 1, RAD6A-(HisRAD18-FLAGRAD18) (I); lane 2, RAD6A-(HisRAD18<sup>ΔC2</sup>-FLAGRAD18<sup>ΔC2</sup>) (IV). Structure of the mutant complex (IV) was represented schematically. Assays were performed as described in (E).

(Figure 2C, lane 2), indicating that only one R6BD in the two RAD18 subunits could form the complex with RAD6A. Again, amounts of each protein in these complexes were compared by CBB staining and western blotting. For Figure 2C, we applied 500 ng of RAD6A-HisRAD18-FLAGRAD18 (lane 1) and 432 ng of the RAD6A-HisRAD18<sup>ΔC1</sup>-FLAGRAD18 complex (lane 2) to SDS-PAGE. If stoichiometry of each component were 1:1:1 in both complexes, amounts of each FLAGRAD18 and each RAD6A in the two complexes should be identical (Figure 2C). The results of CBB staining and western blotting proved to be in good agreement with our estimations (Figure 2C, lanes 1 and 2), suggesting that the

RAD18 dimer, as well as the RAD18 monomer, is capable of accommodating only one RAD6 molecule (Figure 2D). Very interestingly, the hetero complex RAD6A-(HisRAD18<sup>ΔC1</sup>-FLAGRAD18) was catalytically active in terms of PCNA ubiquitination (Figure 2E). As expected, HisRAD18<sup>ΔC1</sup>-FLAGRAD18<sup>ΔC1</sup> was inactive (Figure 2E), and its ligase activity was hardly restored by addition of excess amounts of RAD6A in solution (Figure 2F). To see whether the defect is due to the loss of function of the C-terminal portion, another truncated RAD18 mutant consisting of amino acid residues 1-388 (hereafter designated as RAD18<sup>ΔC2</sup>, Figure 3D), was examined. It was expected that RAD18<sup>ΔC2</sup> could form a




**Figure 3.** RAD18 mutants analyzed in this study. (A) Multiple alignments of the RING domains of human RAD18 and its orthologs. Mutants analyzed in this study are shown below the alignments. Mutants shown in parentheses were eluted in void volumes of gel filtration chromatography. (B) Multiple alignments of UBZ of human RAD18 and its orthologs. (C) Multiple alignments of the SAP domains of human RAD18 and its orthologs. (D) Schematic representations of the structures of RAD18. Positions of the mutations are shown with designated names of the mutant proteins.

complex with RAD6, since it retained R6BD (Figure 3D) (15). Indeed, it was successfully reconstituted into a ternary complex, RAD6A<sup>-His</sup>RAD18<sup>ΔC2</sup>-FLAG-RAD18<sup>ΔC2</sup>, and the complex exhibited ligase activity similar to the wild-type level (Figure 2G), indicating that the C-terminal region of 389–495 amino acid residues is dispensable for ligase activity. From all these results (Figures 1 and 2), together with the fact that RAD6 is essentially a monomer (see Supplementary Materials and Methods section) (19,47), we conclude the subunit composition of the RAD6A–RAD18 complex to be RAD6A–(RAD18)<sub>2</sub> with a molecular mass of 130 kDa. Thus, the overall structure of the complex is asymmetric. Notably, this well matched the 131 kDa estimated molecular mass from the Stokes' radius and the sedimentation coefficient (Table 1).

#### Evidence of the ternary complex, RAD6A–(RAD18)<sub>2</sub>, *in vivo*

To obtain evidence of ternary complex formation *in vivo*, HA-tagged RAD6A and FLAG-tagged RAD18 were expressed together in human cells, and

HA-RAD6A-FLAG-RAD18 complexes were isolated using FLAG affinity and then HA affinity gels. The complexes were a mixture of unmodified RAD18 and mono-ubiquitinated RAD18, as reported previously (Figure 4A) (48). Subunit composition of the purified complexes was determined by western blotting with anti-RAD6 and anti-RAD18 antibodies using RAD6A–(RAD18)<sub>2</sub> complex purified from *E. coli* cells (26) as references. Ratio of signals from HA-RAD6A and FLAG-RAD18 well fitted those from RAD6A–(RAD18)<sub>2</sub> complexes (Figure 4A), suggesting the subunit composition to be RAD6A–(RAD18)<sub>2</sub>. This was further confirmed using partially purified untagged RAD6A–RAD18 complexes generated by conventional column chromatography from RAD6A and RAD18 expressing human cells. We found that RAD6A, RAD18 and mono-ubiquitinated RAD18 all eluted together from a gel filtration column (Figure 4B), similar to the RAD6A–(RAD18)<sub>2</sub> complex (Table 1 and Figure 6C). Again, the subunit composition of the eluted fraction corresponded to RAD6A–(RAD18)<sub>2</sub> (Figure 4B). These results strongly suggest ternary complex formation *in vivo*.



**Figure 4.** Ternary complex formation of RAD6A and RAD18 *in vivo*. (A)  $^{HA}RAD6-^{FLAG}RAD18$  complexes isolated from human cells were analyzed by SDS-PAGE (lane 1) together with  $RAD6A-(RAD18)_2$  isolated from *E. coli* (lanes 2–4) followed by western blotting probed with anti-RAD18 and anti-RAD6 antibodies. Relative chemiluminescence signals of  $^{HA}RAD6A$  and  $^{FLAG}RAD18$  (sum of unmodified  $^{FLAG}RAD18$  and  $Ub-^{FLAG}RAD18$ ) detected with a CCD camera were plotted with those of RAD6A and RAD18 purified from *E. coli* as references. The theoretical 1:1 ratio of RAD6 and RAD18 is shown as a broken line. (B) Untagged RAD6–RAD18 complexes isolated from human cells were analyzed by SDS-PAGE as (A).

### Analysis of UBZ roles in complex formation and ligase activity


Our system to purify RAD6–RAD18 complex with two different tags is a useful tool for analysis of structure–function relationships in RAD18. First, we applied this system to test whether UBZ is required for dimerization of RAD18 using two loss-of-function mutants,  $RAD18^{C207F}$  and  $RAD18^{D221A}$  (34,37) (Figure 3B and D). The results demonstrated that these mutants were successfully reconstituted into the complexes,  $RAD6A-(RAD18^{C207F})_2$  and  $RAD6A-(RAD18^{D221A})_2$  (Figure 5A) and their ligase activities were similar to that of wild-type complex (Figure 5B). These results indicate UBZ to be dispensable for complex formation and ligase activity, in line with recent reports (15,39), demonstrating reliability of our system for analysis of the structure and function of RAD6–RAD18 complexes.

### Functional interaction between RAD6 and the RING domains of the two subunits of RAD18 in the complex


Next, we addressed the question whether the N-terminal part containing a RING domain mediates dimerization. We generated another truncated RAD18 mutant consisting of the 1–115 amino acid residues of RAD18 (hereafter designated as  $RAD18^{\Delta C3}$ , Figure 3D). When  $^{His}RAD18^{\Delta C3}$  and  $^{FLAG}RAD18^{\Delta C3}$  were co-produced with RAD6A in the same *E. coli* cells, we found the  $^{His}RAD18^{\Delta C3}-^{FLAG}RAD18^{\Delta C3}$  dimer to be reconstituted, but that RAD6A did not co-purify with the dimer. The observed elution profile of the dimer from a gel filtration column is shown in Figure 6A.  $^{His}RAD18^{\Delta C3}$  and  $^{FLAG}RAD18^{\Delta C3}$ , confirmed by western blotting (Figure

6B), co-eluted with an apparent molecular mass of ~37 kDa, slightly larger than the calculated molecular mass of 29 kDa as a dimer (Figure 6A and B). As a complementary experiment, an N-terminal deletion mutant of RAD18 consisting of 113–495 amino acid residues (hereafter designated as  $RAD18^{\Delta N1}$ , Figure 3D), was generated. When  $^{His}RAD18^{\Delta N1}$  was coproduced with  $^{FLAG}RAD18$  and RAD6A,  $^{His}RAD18^{\Delta N1}$  co-purified with RAD6A but not with  $^{FLAG}RAD18$  (Figure 6D). As mentioned above, the  $RAD6A-(^{His}RAD18-^{FLAG}RAD18)$  ternary complex eluted at a position corresponding to 490 kDa in gel filtration (Figure 6C). In contrast, the complex of  $RAD6A-^{His}RAD18^{\Delta N1}$  eluted at a position corresponding to 220 kDa, which is much smaller than the ternary complex (Figure 6C). Furthermore the molecular ratio of RAD6A and  $^{His}RAD18^{\Delta N1}$  determined by western blotting, compared with the ternary complex as a reference, was close to 1:1 (Figure 6D). These results suggest that the  $RAD6A-^{His}RAD18^{\Delta N1}$  complex is a dimer composed of one RAD6A and one  $^{His}RAD18^{\Delta N1}$  molecule and imply that  $^{His}RAD18^{\Delta N1}$  neither self-associates nor forms a heterodimer with  $^{FLAG}RAD18$ . From these data shown in Figure 6, we conclude that the N-terminal region (1–115) is necessary and sufficient for dimerization, while UBZ and SAP domains are dispensable.

In general, the RING domains of E3s have an essential function in ligase activity by mediating interactions with E2s (1,2). It is of great interest to clarify how the two RING domains interact with one RAD6 subunit in the  $RAD6A-(RAD18)_2$  complex. To address this point, we made several mutants in which one or two conserved amino acid residues in the RING domain were replaced



**Figure 5.** Analysis of ligase activity and complex formation with UBZ mutants of RAD18. (A) Purified complexes (3.7 pmol) were analyzed by SDS-PAGE followed by staining with CBB. I, RAD6A-(HisRAD18-FLAGRAD18); II, RAD6A-(HisRAD18<sup>C207F</sup>-FLAGRAD18<sup>C207F</sup>); III, RAD6A-(HisRAD18<sup>D221A</sup>-FLAGRAD18<sup>D221A</sup>). Structures are represented schematically, UBZ domains with a mutation being shown in white boxes. (B) Ligase activity of the respective RAD6A-RAD18 complexes. Increasing amounts of the complexes (0.5, 1 and 2 pmol) shown in (A) were subjected to standard assays.




**Figure 6.** Analysis of RING domain for dimerization of RAD18. (A) Elution profile of HisRAD18<sup>ΔC2</sup>-FLAGRAD18<sup>ΔC2</sup> complexes from a Superdex 200 gel filtration column. The size markers, ferritin (440 kDa), aldolase (158 kDa), albumin (67 kDa), ovalbumin (43 kDa) and ribonuclease A (14 kDa) were eluted in 10.09, 12.26, 13.74, 14.98 and 17.56 ml, respectively. The complex was eluted in 15.35 ml, estimated the apparent molecular mass of the complex to be 37 kDa from a standard curve of the marker proteins. (B) Analysis of fractions eluted from a Superdex 200 gel filtration column (A). Fractions between 13 and 18.5 ml were analyzed by SDS-PAGE followed by staining with CBB and western blotting probed with the indicated antibodies. HisRAD18<sup>ΔC2</sup> (15.1 kDa) migrated slightly faster than FLAGRAD18<sup>ΔC2</sup> (14.2 kDa). (C) Elution profile of RAD6A-(HisRAD18-FLAGRAD18) and RAD6A-HisRAD18<sup>ΔN1</sup> complexes from a Superdex 200 gel filtration column. Respective complexes were eluted in 9.77 and 11.45 ml, corresponding to 490 and 220 kDa of the apparent molecular masses, and 62 and 52 Å of Stokes' radius, estimated from a standard curve of the marker proteins. (D) Peak fractions of gel filtration chromatography (C) were analyzed by SDS-PAGE followed by CBB staining and western blotting probed with the indicated antibodies. Relative chemiluminescence signals detected with a CCD camera are shown. Lane 1, RAD6A-(HisRAD18-FLAGRAD18) (3.7 pmol as a trimer); lane 2, RAD6A-HisRAD18<sup>ΔN1</sup> (3.7 pmol as a dimer).

as indicated in Figure 3A. We found that RING mutants with C25A, C25S, C25F, I27A, C28S, C27F or F53A/L54A substitutions eluted in void volumes on gel filtration chromatography, suggesting these mutants to form large aggregates due to highly disordered structures by misfolding. Therefore we did not further analyze them. However, we could successfully obtain one mutant

complex containing I50A/R51A substitutions in RAD18 (hereafter designated as RAD18<sup>RING</sup>, see Figures 3A and 7A, lane 2). A previous report suggested, based on the crystal structure of the c-Cbl-UbcH7 complex, that the amino acid residues Ile50 and Arg51 of RAD18 should be located in a predicted RAD6-interacting  $\alpha$ -helix (2). Thus, it was expected that I50A/R51A mutations would affect





**Figure 7.** Analysis of ligase activity and complex formation of a RING mutant of RAD18. (A) Purified complexes (3.7 pmol) were analyzed by SDS-PAGE followed by staining with CBB. I, RAD6A-(<sup>His</sup>RAD18-<sup>FLAG</sup>RAD18); II, RAD6A-(<sup>His</sup>RAD18<sup>RING</sup>-<sup>FLAG</sup>RAD18<sup>RING</sup>); III, RAD6A-(<sup>His</sup>RAD18<sup>RING</sup>-<sup>FLAG</sup>RAD18). Structures are represented schematically, RING domains with a mutation being shown in white boxes. (B) Ligase activities of the respective RAD6A-RAD18 complexes. Increasing amounts of the complexes (0.5, 1 and 2 pmol) shown in (A) were subjected to standard assays. (C) Purified complexes (3.7 pmol) were analyzed by SDS-PAGE followed by staining with CBB. I, RAD6A-(<sup>His</sup>RAD18<sup>AC1</sup>-<sup>FLAG</sup>RAD18); II, RAD6A-(<sup>His</sup>RAD18<sup>RING</sup>-<sup>FLAG</sup>RAD18<sup>AC1</sup>); III, RAD6A-(<sup>His</sup>RAD18<sup>RING/AC1</sup>-<sup>FLAG</sup>RAD18). (D) Ligase activities of the respective RAD6A-RAD18 complexes. Increasing amounts of the complexes (0.5, 1 and 2 pmol) shown in (C) were subjected to standard assays.

the ligase activity of RAD18. In fact, it was much reduced when compared to that of the wild-type complex (Figure 7B). Then, we reconstituted a hetero complex with the mutant and wild-type, RAD6A-(<sup>His</sup>RAD18<sup>RING</sup>-<sup>FLAG</sup>RAD18) (Figure 7A, lane 3). Surprisingly, its ligase activity was essentially identical to that of the wild-type (Figure 7B), indicating that an interaction between RAD6 and only one RING domain in the two RAD18 subunits is sufficient for ligase activity. In addition, I50A/R51A mutations were combined with the  $\Delta$ C1 mutation and two mutant complexes, RAD6A-(<sup>His</sup>RAD18<sup>RING</sup>-<sup>FLAG</sup>RAD18<sup>AC1</sup>) and RAD6A-(<sup>His</sup>RAD18<sup>RING/AC1</sup>-<sup>FLAG</sup>RAD18) were reconstituted (Figure 7C). Analysis of their ligase activities demonstrated these to be as active as the wild-type (Figure 7D), indicating that one RAD6 molecule in the complex has the potential to interact with either subunit of the RAD18 dimer.

#### Functional interaction between SAP and R6BD for ligase activity


It has been shown that the SAP domain of RAD18 has DNA-binding activity (15,38), which is separable from its essential function for ligase activity (38). When a mutant RAD18 containing L250A/L265A substitutions (hereafter designated as RAD18<sup>SAP</sup>, see Figure 3D) was successfully reconstituted into the ternary complex, RAD6A-(<sup>His</sup>RAD18<sup>SAP</sup>-<sup>FLAG</sup>RAD18<sup>SAP</sup>) (Figure 8A, lane 2), the

ligase activity of the mutant complex was reduced to an undetectable level (Figure 8B). However, activity was restored in a hetero complex with the wild-type, RAD6A-(<sup>His</sup>RAD18<sup>SAP</sup>-<sup>FLAG</sup>RAD18) (Figure 8B), demonstrating that only one of the two SAP domains in the complex is sufficient for the essential SAP function. Next, the SAP mutation was combined with the  $\Delta$ C1 mutation to reconstitute RAD6A-(<sup>His</sup>RAD18<sup>SAP</sup>-<sup>FLAG</sup>RAD18<sup>AC1</sup>) and RAD6A-(<sup>His</sup>RAD18<sup>SAP/AC1</sup>-<sup>FLAG</sup>RAD18) (Figure 8C). Interestingly, the former had quite reduced ligase activity, but the latter exhibited the wild-type level (Figure 8D), indicating that the active SAP domain should be present on the same RAD18 molecule to which RAD6 binds. In contrast, such functional interaction was not observed between RING and SAP mutants; ligase activities of RAD6A-(<sup>His</sup>RAD18<sup>RING</sup>-<sup>FLAG</sup>RAD18<sup>SAP</sup>) and RAD6A-(<sup>His</sup>RAD18<sup>RING</sup>-<sup>FLAG</sup>RAD18<sup>SAP</sup>) complexes (Figure 8E) were similar to that of wild-type (Figure 8F).

#### DISCUSSION

In this study, we established a method to distinguish between the two subunits of RAD18 in the RAD6-(RAD18)<sub>2</sub> complex by introducing different tags at the N-termini. This enabled us to purify RAD18 complexes composed of wild-wild, wild-mutant or mutant-mutant subunits and facilitated analysis of structure-function relationships in RAD18.





**Figure 8.** Analysis of ligase activity and complex formation of a SAP mutant of RAD18. (A) Purified complexes (3.7 pmol) were analyzed by SDS-PAGE followed by staining with CBB. I, RAD6A-(<sup>His</sup>RAD18-<sup>FLAG</sup>RAD18); II, RAD6A-(<sup>His</sup>RAD18<sup>SAP</sup>-<sup>FLAG</sup>RAD18<sup>SAP</sup>); III, RAD6A-(<sup>His</sup>RAD18<sup>SAP</sup>-<sup>FLAG</sup>RAD18). Structures were represented schematically, SAP domains with a mutation being shown in white boxes. (B) Ligase activities of the respective RAD6A-RAD18 complexes. Increasing amounts of the complexes (0.5, 1 and 2 pmol) shown in (A) were subjected to standard assays. (C) Purified complexes (3.7 pmol) were analyzed by SDS-PAGE followed by staining with CBB. I, RAD6A-(<sup>His</sup>RAD18<sup>ΔC1</sup>-<sup>FLAG</sup>RAD18); II, RAD6A-(<sup>His</sup>RAD18<sup>SAP</sup>-<sup>FLAG</sup>RAD18<sup>ΔC1</sup>); III, RAD6A-(<sup>His</sup>RAD18<sup>SAP/ΔC1</sup>-<sup>FLAG</sup>RAD18). (D) Ligase activities of the respective RAD6A-RAD18 complexes. Increasing amounts of the complexes (0.5, 1 and 2 pmol) shown in (C) were subjected to standard assays. (E) Purified complexes (3.7 pmol) were analyzed by SDS-PAGE followed by staining with CBB. I, RAD6A-(<sup>His</sup>RAD18-<sup>FLAG</sup>RAD18); II, RAD6A-(<sup>His</sup>RAD18<sup>RING/SAP</sup>-<sup>FLAG</sup>RAD18); III, RAD6A-(<sup>His</sup>RAD18<sup>RING</sup>-<sup>FLAG</sup>RAD18<sup>SAP</sup>). (F) Ligase activities of the respective RAD6A-RAD18 complexes. Increasing amounts of the complexes (0.5, 1 and 2 pmol) shown in (E) were subjected to standard assays.

Previously, it was considered that the RAD6-RAD18 complex could be composed of two RAD6 and two RAD18, since the observations that RAD18 binds to RAD6 and it forms a dimer led to the assumption that each RAD18 molecule in the dimer should bind to RAD6 (15,20,21). However, there is little evidence for such an assignment based on the stoichiometry of RAD6. In this study, we provided hard lines of evidence that the complex is composed of one subunit of RAD6 and two subunits of RAD18, so that the overall structure of the ternary complex should be asymmetric.

The RING domains in E3s have an essential function in ligase activity by mediating interactions with E2s. Zheng

*et al.* (2) have suggested, on the basis of the crystal structure of the c-Cbl-UbcH7 complex, that amino acid residues I50 and R51 in the RING domain of RAD18 are located in a predicted RAD6-interacting  $\alpha$ -helix. Consistent with this, a mutant complex containing I50A/R51A substitutions in both RAD18 subunits exhibited severely reduced ligase activity, probably due to an impaired interaction between the altered RING domain and RAD6A. When the mutant subunit was complexed with wild-type RAD18 subunit, ligase activity was restored to the wild-type level, indicating that only one of the two RING domains is necessary for enzyme function. Furthermore, even when the RING mutation

was combined with a deletion mutation of R6BD in the same or other RAD18 subunit, the complexes exhibited robust ligase activity. These results imply that RAD6 can bind in either one of the two R6BDs, then interacting with one of the two RING domains in the RAD18 dimer.

Why should the RING domain of RAD18 form a dimer? We found that inactivation of one RING domain does not affect the ligase activity, indicating that the close proximity of two active RING domains in RAD18 is not important for the enzyme function. From our observations that some RING mutants such as C25A, C25S, C25F, I27A, C28S, C28F and F53A/L54A, probably disrupting the RING structure (Figure 3D), appeared to form large aggregates, we suggest that the RING structure is required for stable dimerization, which is in turn necessary for sustaining the functionally active RING structure of RAD18. From the results, we cannot exclude the possibility of retention of some enzyme functions even in aggregates. Indeed, it has been reported that the C28F mutant can fully complement the homologous recombination defects of RAD18-null cells (39).

The SAP domain is a unique eukaryotic module involved in sequence- or structure-specific DNA binding (36,38,49–52). Additionally, in RAD18 it has an essential role in ligase activity, separable from its function in DNA binding (38). We further demonstrated that one of the two SAP domains in the RAD6–(RAD18)<sub>2</sub> complex is sufficient for ligase activity. Interestingly, experiments using hetero complexes, in which mutation in the SAP domain was combined with a deletion mutation of R6BD, revealed an interesting relationship between SAP and R6BD. Although one of the two SAP domains and one of the two R6BDs are sufficient for enzyme activity, the active SAP domain should be present on the same RAD18 molecule to which RAD6 binds in the complex. It makes a sharp contrast to the case of RING mutant as described above. A simple explanation is that the SAP domain acts as a hinge connecting the R6BD and RING domain to enable precise juxtaposition between RING and RAD6. This might be very important for ligase activity because a deletion mutant of R6BD hardly supported PCNA ubiquitination in the presence of an excess amount of RAD6A, which should be high enough to detect ligase activities for many other E2–E3 pairs (4–6,9–2). It seems likely that binding affinity between RAD6 and the RING domain of RAD18 is very low. That could be the reason why R6BD is essential for ligase activity of the RAD6–RAD18 complex. The tight interaction between RAD6 and R6BD confers on the complex the ability to monoubiquitinate PCNA, also inhibiting the activity of RAD6 catalyzing ubiquitin chain formation (19).

In the above, we have shown that RAD6 and RAD18 forms a ternary complex, RAD6–(RAD18)<sub>2</sub> and demonstrated that one R6BD site is sufficient for the ligase activity. Then, a question that immediately arises is why RAD6 can bind to one of the two R6BD sites, but not to both of them. Although we have no experimental data to answer the question at the moment, such a case is not unprecedented. CHIP (*C*-terminal of *Hsp70* interacting protein), a protein containing a *C*-terminal Ubox domain (similar to RING) and an *N*-terminal

TPR domain, forms an asymmetric dimer mediated by the Ubox domains, in which only one of the two Ubox domains is available for binding to E2 (Ubc13) because the other Ubox domain is blocked due to interaction with the TPR domain (53). It is tempting to speculate that a segment including R6BD located in a *C*-terminal portion of the two RAD18 subunits may interact with one of the two RING domains, consequently preventing the R6BD from interacting with RAD6. Thus, one RAD6 molecule can bind to the other R6BD and is positioned closely to another unoccupied RING domain for catalyzing the ligase function. Evidently, further experiments are required to elucidate this interesting possibility.

During the course of finally completing this manuscript, we have learned that Huang and co-workers have recently determined the structure of a RAD18 RING(1–99) dimer, also showing that this binds to RAD6 at a 2:2 ratio, whereas the full-length RAD18 dimer binds only to a single RAD6 molecule (54). Our study was conducted independently of their work.

## SUPPLEMENTARY DATA

Supplementary Data are available at NAR Online: Supplementary Materials and Methods, Supplementary References [26,45,55].

## ACKNOWLEDGEMENTS

The authors thank Dr Toshiki Tsurimoto (Kyushu University, Fukuoka, Japan) and Dr Tomohiko Ohta (St Marianna University School of Medicine, Kanagawa, Japan) for providing PCNA expression and ubiquitin-encoding plasmids, respectively. The authors would also like to express our appreciation to Dr Haruo Ohmori (Kyoto University, Kyoto, Japan) for his critical reading of manuscripts and valuable suggestions. The authors are grateful to Fumie Okubo, Kazumi Shimamoto and Mai Yoshida for their laboratory assistance.

## FUNDING

Grants-in-Aid from the Ministry of Education, Culture, Sports, Science and Technology of Japan (to Y.M. and K.K.); Grants-in-Aid for Cancer Research from the Ministry of Health, Labour and Welfare (to K.K.). Funding for open access charge: Grants-in-Aid from the Ministry of Education, Culture, Sports, Science and Technology of Japan.

*Conflict of interest statement.* None declared.

## REFERENCES

1. Deshaies, R.J. and Joazeiro, C.A. (2009) RING domain E3 ubiquitin ligases. *Annu. Rev. Biochem.*, **78**, 399–434.
2. Zheng, N., Wang, P., Jeffrey, P.D. and Pavletich, N.P. (2000) Structure of a c-Cbl-UbcH7 complex: RING domain function in ubiquitin-protein ligases. *Cell*, **102**, 533–539.

3. Brzovic, P.S., Rajagopal, P., Hoyt, D.W., King, M.C. and Kleit, R.E. (2001) Structure of a BRCA1-BARD1 heterodimeric RING-RING complex. *Nat. Struct. Biol.*, **8**, 833–837.
4. Hashizume, R., Fukuda, M., Maeda, I., Nishikawa, H., Oyake, D., Yabuki, Y., Ogata, H. and Ohta, T. (2001) The RING heterodimer BRCA1-BARD1 is a ubiquitin ligase inactivated by a breast cancer-derived mutation. *J. Biol. Chem.*, **276**, 14537–14540.
5. Xia, Y., Pao, G.M., Chen, H.W., Verma, I.M. and Hunter, T. (2003) Enhancement of BRCA1 E3 ubiquitin ligase activity through direct interaction with the BARD1 protein. *J. Biol. Chem.*, **278**, 5255–5263.
6. Buchwald, G., van der Stoop, P., Weichenrieder, O., Perrakis, A., van Lohuizen, M. and Sixma, T.K. (2006) Structure and E3-ligase activity of the Ring-Ring complex of polycomb proteins Bmi1 and Ring1b. *EMBO J.*, **25**, 2465–2474.
7. Satijn, D.P. and Otte, A.P. (1999) RING1 interacts with multiple Polycomb-group proteins and displays tumorigenic activity. *Mol. Cell. Biol.*, **19**, 57–68.
8. Kawai, H., Lopez-Pajares, V., Kim, M.M., Wiederschain, D. and Yuan, Z.M. (2007) RING domain-mediated interaction is a requirement for MDM2's E3 ligase activity. *Cancer Res.*, **67**, 6026–6030.
9. Linares, L.K., Hengstermann, A., Ciechanover, A., Muller, S. and Scheffner, M. (2003) HdmX stimulates Hdm2-mediated ubiquitination and degradation of p53. *Proc. Natl Acad. Sci. USA*, **100**, 12009–12014.
10. Linke, K., Mace, P.D., Smith, C.A., Vaux, D.L., Silke, J. and Day, C.L. (2008) Structure of the MDM2/MDMX RING domain heterodimer reveals dimerization is required for their ubiquitylation in trans. *Cell Death Differ.*, **15**, 841–848.
11. Mace, P.D., Linke, K., Feltham, R., Schumacher, F.R., Smith, C.A., Vaux, D.L., Silke, J. and Day, C.L. (2008) Structures of the cIAP2 RING domain reveal conformational changes associated with ubiquitin-conjugating enzyme (E2) recruitment. *J. Biol. Chem.*, **283**, 31633–31640.
12. Liew, C.W., Sun, H., Hunter, T. and Day, C.L. (2010) RING domain dimerization is essential for RNF4 function. *Biochem. J.*, **431**, 23–29.
13. Bailly, V., Lamb, J., Sung, P., Prakash, S. and Prakash, L. (1994) Specific complex formation between yeast RAD6 and RAD18 proteins: a potential mechanism for targeting RAD6 ubiquitin-conjugating activity to DNA damage sites. *Genes Dev.*, **8**, 811–820.
14. Bailly, V., Lauder, S., Prakash, S. and Prakash, L. (1997) Yeast DNA repair proteins Rad6 and Rad18 form a heterodimer that has ubiquitin conjugating, DNA binding, and ATP hydrolytic activities. *J. Biol. Chem.*, **272**, 23360–23365.
15. Notenboom, V., Hibbert, R.G., van Rossum-Fikkert, S.E., Olsen, J.V., Mann, M. and Sixma, T.K. (2007) Functional characterization of Rad18 domains for Rad6, ubiquitin, DNA binding and PCNA modification. *Nucleic Acids Res.*, **35**, 5819–5830.
16. Bailly, V., Prakash, S. and Prakash, L. (1997) Domains required for dimerization of yeast Rad6 ubiquitin-conjugating enzyme and Rad18 DNA binding protein. *Mol. Cell. Biol.*, **17**, 4536–4543.
17. Watanabe, K., Tateishi, S., Kawasuji, M., Tsurimoto, T., Inoue, H. and Yamaizumi, M. (2004) Rad18 guides pol $\eta$  to replication stalling sites through physical interaction and PCNA monoubiquitination. *EMBO J.*, **23**, 3886–3896.
18. Tateishi, S., Sakuraba, Y., Masuyama, S., Inoue, H. and Yamaizumi, M. (2000) Dysfunction of human Rad18 results in defective postreplication repair and hypersensitivity to multiple mutagens. *Proc. Natl Acad. Sci. USA*, **97**, 7927–7932.
19. Hibbert, R.G., Huang, A., Boelens, R. and Sixma, T.K. (2011) E3 ligase Rad18 promotes monoubiquitination rather than ubiquitin chain formation by E2 enzyme Rad6. *Proc. Natl Acad. Sci. USA*, **108**, 5590–5595.
20. Miyase, S., Tateishi, S., Watanabe, K., Tomita, K., Suzuki, K., Inoue, H. and Yamaizumi, M. (2005) Differential regulation of Rad18 through Rad6-dependent mono- and polyubiquitination. *J. Biol. Chem.*, **280**, 515–524.
21. Ulrich, H.D. and Jentsch, S. (2000) Two RING finger proteins mediate cooperation between ubiquitin-conjugating enzymes in DNA repair. *EMBO J.*, **19**, 3388–3397.
22. Friedberg, E.C., Walker, G.C., Siede, W., Wood, R.D., Schultz, R.A. and Ellenberger, T. (2006) DNA repair and mutagenesis. *2nd edn.* ASM, Washington, DC.
23. Hoege, C., Pfander, B., Moldovan, G.L., Pyrowolakis, G. and Jentsch, S. (2002) RAD6-dependent DNA repair is linked to modification of PCNA by ubiquitin and SUMO. *Nature*, **419**, 135–141.
24. Garg, P. and Burgers, P.M. (2005) Ubiquitinated proliferating cell nuclear antigen activates translesion DNA polymerases  $\eta$  and REV1. *Proc. Natl Acad. Sci. USA*, **102**, 18361–18366.
25. Haracska, L., Unk, I., Prakash, L. and Prakash, S. (2006) Ubiquitylation of yeast proliferating cell nuclear antigen and its implications for translesion DNA synthesis. *Proc. Natl Acad. Sci. USA*, **103**, 6477–6482.
26. Masuda, Y., Piao, J. and Kamiya, K. (2010) DNA replication-coupled PCNA mono-ubiquitination and polymerase switching in a human *in vitro* system. *J. Mol. Biol.*, **396**, 487–500.
27. Unk, I., Hajdu, I., Fatyol, K., Hurwitz, J., Yoon, J.H., Prakash, L., Prako, S. and Haracska, L. (2008) Human HLF functions as a ubiquitin ligase for proliferating cell nuclear antigen polyubiquitination. *Proc. Natl Acad. Sci. USA*, **105**, 3768–3773.
28. Unk, I., Hajdu, I., Fatyol, K., Szakal, B., Blastyak, A., Bermudez, V., Hurwitz, J., Prakash, L., Prako, S. and Haracska, L. (2006) Human SHPRH is a ubiquitin ligase for Mms2-Ubc13-dependent polyubiquitylation of proliferating cell nuclear antigen. *Proc. Natl Acad. Sci. USA*, **103**, 18107–18112.
29. Bienko, M., Green, C.M., Crossetto, N., Rudolf, F., Zapart, G., Coull, B., Kannouche, P., Wider, G., Peter, M., Lehmann, A.R. *et al.* (2005) Ubiquitin-binding domains in Y-family polymerases regulate translesion synthesis. *Science*, **310**, 1821–1824.
30. Wood, A., Garg, P. and Burgers, P.M. (2007) A ubiquitin-binding motif in the translesion DNA polymerase Rev1 mediates its essential functional interaction with ubiquitinated proliferating cell nuclear antigen in response to DNA damage. *J. Biol. Chem.*, **282**, 20256–20263.
31. Zhuang, Z., Johnson, R.E., Haracska, L., Prakash, L., Prakash, S. and Benkovic, S.J. (2008) Regulation of polymerase exchange between Pol $\eta$  and Pol $\delta$  by monoubiquitination of PCNA and the movement of DNA polymerase holoenzyme. *Proc. Natl Acad. Sci. USA*, **105**, 5361–5366.
32. Day, T.A., Palle, K., Barkley, L.R., Kakusho, N., Zou, Y., Tateishi, S., Verreault, A., Masai, H. and Vaziri, C. (2010) Phosphorylated Rad 18 directs DNA polymerase  $\eta$  to sites of stalled replication. *J. Cell. Biol.*, **191**, 953–966.
33. Bish, R.A. and Myers, M.P. (2007) Werner helicase-interacting protein 1 binds polyubiquitin via its zinc finger domain. *J. Biol. Chem.*, **282**, 23184–23193.
34. Crossetto, N., Bienko, M., Hibbert, R.G., Perica, T., Ambrogio, C., Kensche, T., Hofmann, K., Sixma, T.K. and Dikic, I. (2008) Human Wrn1p is localized in replication factories in a ubiquitin-binding zinc finger-dependent manner. *J. Biol. Chem.*, **283**, 35173–35185.
35. Hofmann, K. (2009) Ubiquitin-binding domains and their role in the DNA damage response. *DNA Repair*, **8**, 544–556.
36. Aravind, L. and Koonin, E.V. (2000) SAP - a putative DNA-binding motif involved in chromosomal organization. *Trends Biochem. Sci.*, **25**, 112–114.
37. Nakajima, S., Lan, L., Kanno, S., Usami, N., Kobayashi, K., Mori, M., Shiomi, T. and Yasui, A. (2006) Replication-dependent and -independent responses of RAD18 to DNA damage in human cells. *J. Biol. Chem.*, **281**, 34687–34695.
38. Tsuji, Y., Watanabe, K., Araki, K., Shinohara, M., Yamagata, Y., Tsurimoto, T., Hanaoka, F., Yamamura, K., Yamaizumi, M. and Tateishi, S. (2008) Recognition of forked and single-stranded DNA structures by human RAD18 complexed with RAD6B protein triggers its recruitment to stalled replication forks. *Genes Cells*, **13**, 343–354.
39. Huang, J., Huen, M.S., Kim, H., Leung, C.C., Glover, J.N., Yu, X. and Chen, J. (2009) RAD18 transmits DNA damage signalling to elicit homologous recombination repair. *Nat. Cell. Biol.*, **11**, 592–603.
40. Fukuda, K., Morioka, H., Imajou, S., Ikeda, S., Ohtsuka, E. and Tsurimoto, T. (1995) Structure-function relationship of the eukaryotic DNA replication factor, proliferating cell nuclear antigen. *J. Biol. Chem.*, **270**, 22527–22534.

41. Masuda, Y., Suzuki, M., Piao, J., Gu, Y., Tsurimoto, T. and Kamiya, K. (2007) Dynamics of human replication factors in the elongation phase of DNA replication. *Nucleic Acids Res.*, **35**, 6904–6916.
42. Tomida, J., Masuda, Y., Hiroaki, H., Ishikawa, T., Song, I., Tsurimoto, T., Tateishi, S., Shiomi, T., Kamei, Y., Kim, J. *et al.* (2008) DNA damage-induced ubiquitylation of RFC2 subunit of replication factor C complex. *J. Biol. Chem.*, **283**, 9071–9079.
43. Niwa, H., Yamamura, K. and Miyazaki, J. (1991) Efficient selection for high-expression transfectants with a novel eukaryotic vector. *Gene*, **108**, 193–199.
44. Maki, H. and Kornberg, A. (1987) Proofreading by DNA polymerase III of *Escherichia coli* depends on cooperative interaction of the polymerase and exonuclease subunits. *Proc. Natl Acad. Sci. USA*, **84**, 4389–4392.
45. Studier, F.W., Rosenberg, A.H., Dunn, J.J. and Dubendorff, J.W. (1990) Use of T7 RNA polymerase to direct expression of cloned genes. *Methods Enzymol.*, **185**, 60–89.
46. Siegel, L.M. and Monty, K.J. (1966) Determination of molecular weights and frictional ratios of proteins in impure systems by use of gel filtration and density gradient centrifugation. Application to crude preparations of sulfite and hydroxylamine reductases. *Biochim. Biophys. Acta*, **112**, 346–362.
47. Worthylake, D.K., Prakash, S., Prakash, L. and Hill, C.P. (1998) Crystal structure of the *Saccharomyces cerevisiae* ubiquitin-conjugating enzyme Rad6 at 2.6 Å resolution. *J. Biol. Chem.*, **273**, 6271–6276.
48. Yuasa, M.S., Masutani, C., Hirano, A., Cohn, M.A., Yamaizumi, M., Nakatani, Y. and Hanaoka, F. (2006) A human DNA polymerase  $\eta$  complex containing Rad18, Rad6 and Rev1; proteomic analysis and targeting of the complex to the chromatin-bound fraction of cells undergoing replication fork arrest. *Genes Cells*, **11**, 731–744.
49. Ahn, J.S. and Whitby, M.C. (2003) The role of the SAP motif in promoting Holliday junction binding and resolution by SpCCE1. *J. Biol. Chem.*, **278**, 29121–29129.
50. Chou, C.H., Wang, J., Knuth, M.W. and Reeves, W.H. (1992) Role of a major autoepitope in forming the DNA binding site of the p70 (Ku) antigen. *J. Exp. Med.*, **175**, 1677–1684.
51. Gohring, F., Schwab, B.L., Nicotera, P., Leist, M. and Fackelmayer, F.O. (1997) The novel SAR-binding domain of scaffold attachment factor A (SAF-A) is a target in apoptotic nuclear breakdown. *EMBO J.*, **16**, 7361–7371.
52. Kipp, M., Gohring, F., Ostendorp, T., van Drunen, C.M., van Driel, R., Przybylski, M. and Fackelmayer, F.O. (2000) SAF-Box, a conserved protein domain that specifically recognizes scaffold attachment region DNA. *Mol. Cell. Biol.*, **20**, 7480–7489.
53. Zhang, M., Windheim, M., Roe, S.M., Pegg, M., Cohen, P., Prodromou, C. and Pearl, L.H. (2005) Chaperoned ubiquitylation—crystal structures of the CHIP U box E3 ubiquitin ligase and a CHIP-Ubc13-Uev1a complex. *Mol. Cell*, **20**, 525–538.
54. Huang, A., Hibbert, R.G., de Jong, R.N., Das, D., Sixma, T.K. and Boelens, R. (2011) Symmetry and asymmetry of the RING-RING dimer of Rad18. *J. Mol. Biol.*, **410**, 424–435.
55. Gomes, X.V., Gary, S.L. and Burgers, P.M. (2000) Overproduction in *Escherichia coli* and characterization of yeast replication factor C lacking the ligase homology domain. *J. Biol. Chem.*, **275**, 14541–14549.