

広島大学学術情報リポジトリ
Hiroshima University Institutional Repository

Title	Nonneutral evolution of volume fluctuations in lysozymes revealed by normal-mode analysis of compressibility
Author(s)	Mimura, Shigefumi; Yamato, Takahisa; Kamiyama, Tadashi; Gekko, Kunihiko
Citation	Biophysical Chemistry , 161 : 39 - 45
Issue Date	2012
DOI	10.1016/j.bpc.2011.11.005
Self DOI	
URL	https://ir.lib.hiroshima-u.ac.jp/00034798
Right	(c) 2012 Elsevier B.V. All rights reserved.
Relation	



(Research Article)

Nonneutral evolution of volume fluctuations in lysozymes revealed by normal-mode analysis of compressibility

Shigefumi Mimura^a, Takahisa Yamato^b, Tadashi Kamiyama^c, Kunihiko Gekko^{a,*}

^a *Department of Mathematical and Life Sciences, Graduate School of Science, Hiroshima University, 1-3-1 Kagamiyama, Higashi-Hiroshima 739-8526, Japan*

^b *Graduate School of Science, Nagoya University, Chikusa-ku, Nagoya 464-8602, Japan*

^c *Department of Chemistry, School of Science and Engineering, Kinki University, 3-4-1 Kowakae, Higashi-Osaka 577-8502, Japan*

Abbreviations: TIS, wild type with Thr40/Ile55/Ser91 (chicken); SVT, wild type with Ser40/Val55/Thr91 (bobwhite); SIS, variant with Ser40/Ile55/Ser91; TIT, variant with Thr40/Ile55/Thr91; TVS, variant with Thr40/Val55/Ser91; SIT, variant with Ser40/Ile55/Thr91; SVS, variant with Ser40/Val55/Ser91; TVT, variant with Thr40/Val55/Thr91; PDB, Protein Data Bank.

* *Corresponding author (Present address):* Institute for Sustainable Sciences and Development, Hiroshima University, 1-3-1 Kagamiyama, Higashi-Hiroshima 739-8526, Japan. Phone, +81-82-424-6481; Fax, +81-82-424-7327; E-mail, gekko@hiroshima-u.ac.jp

ABSTRACT

The evolution of structural fluctuations of proteins was examined by calculating the isothermal compressibility (β_T) values of chicken lysozyme and its six evolutionary mutants at Thr40, Ile55, and Ser91 (a ternary mutant corresponding to bobwhite lysozyme) from their X-ray structures by normal-mode analysis at 300 K. The β_T values of the two extant lysozymes from chicken and bobwhite were 1.61 and 1.59 Mbar⁻¹, respectively, but five other evolutionary mutants showed larger β_T values of up to 2.17 Mbar⁻¹. These results suggest that ancestral lysozymes exhibit larger volume fluctuations than extant ones, and hence that the molecular evolution of lysozymes has followed a nonneutral evolutionary pathway. The evolutionary mutants contained large amount of cavities, although no change was visible in the X-ray structures. There was a linear correlation between β_T and total cavity volume, predicting that the cavity volume or atomic packing is an important factor regulating volume fluctuations during the molecular evolution of this protein.

Keywords: isothermal compressibility, lysozyme, molecular evolution, nonneutral pathway, normal-mode analysis, volume fluctuations

1. Introduction

There has been considerable controversy about whether protein evolution follows a neutral or nonneutral pathway [1–4], but this problem remains undefined at the molecular level because few proteins have a series of evolutionary variants. Game-bird lysozymes consist of 129 total amino-acid residues, 3 of which (residues 40, 55, and 91) are occupied either by the triplet Thr40/Ile55/Ser91 (TIS) for chicken or by the triplet Ser40/Val55/Thr91 (SVT) for bobwhite. These three residues are located together just beneath the active-site cleft in the three-dimensional structure [5,6], and they play an important role in the enzyme function. Lysozymes in game birds are encoded by a single gene [7] in which the possibility of simultaneous double or triple mutations is negligibly small. Hence, their ancestral lysozymes could be expected to evolve along the possible three-step pathways between these triplets shown in Fig. 1: TIS (chicken) \rightarrow SIS (variant with Ser40/Ile55/Ser91), TIT (variant with Thr40/Ile55/Thr91), TVS (variant with Thr40/Val55/Ser91) (single mutation) \rightarrow SIT (variant with Ser40/Ile55/Thr91), SVS (variant with Ser40/Val55/Ser91), and TVT (variant with Thr40/Val55/Thr91) (double mutation) \rightarrow SVT (bobwhite); however, none of these six intermediates (SIS, TIT, TVS, SIT, SVS, and TVT) is found in extant lysozymes. Malcolm *et al.* [5] constructed these six intermediates by site-directed mutagenesis of chicken lysozyme and found that the thermostabilities of the variants fell outside the range of the two extant lysozymes from chicken and bobwhite. This finding suggests that these mutations have followed a nonneutral evolutionary pathway. It is of interest how the structural fluctuations of these ancestral lysozymes have changed during the molecular evolution to gain functional adaptation.

(Fig. 1)

The structural fluctuation of proteins has been investigated by various techniques, including X-ray crystallography (B-factor), NMR (order parameter), hydrogen-deuterium exchange, compressibility, and computer simulation [8]. Compressibility is an important measure of volume fluctuation [9] because it is sensitive to packing defects or cavities in the interior of a protein molecule, and these are easily compressed by pressure and result in internal motions or flexibility in response to thermal or mechanical forces. Directly measuring the isothermal compressibility (β_T) of a protein is generally difficult, as reflected by the small number of related reports [10,11]. This has led to most studies of protein compressibility

considering the adiabatic compressibility, β_s , which can be determined more easily and accurately by sound-velocity measurements [12–21]. An important finding of these compressibility studies is that substituting a single amino acid can induce large changes in β_s , and that mutants having larger β_s values exhibit higher enzyme activity [18–21]. This means that the local structural changes associated with mutation significantly modify the overall dynamics of a protein so as to affect its function. Computer simulation of β_T is useful to obtain a detailed description of molecular movement at the atomic level, and can be implemented by the molecular dynamics and the normal-mode analysis [22–29]. The former is more common due to simulations now extending into the nanosecond range, while normal-mode analysis is advantageous for rapidly calculating β_T based on a harmonic approximation of the potential energy function around a minimum energy conformation.

To address how volume fluctuations vary during protein evolution, the present study calculated the β_T values of the above extant and evolutionary lysozymes (except for SVS, whose X-ray structure has not yet been determined) by normal-mode analysis because no experimental β_T values are known for these evolutionary lysozymes. The β_T values obtained are discussed in terms of the modified atomic packing and the molecular evolution of lysozymes.

2. Methods

2.1. Normal-mode analysis

The β_T values of lysozymes were calculated as a sum of the volume fluctuations from individual normal modes within the potential energy of the harmonic range, essentially according to the method of Yamato and colleagues developed for studying the dynamics of deoxymyoglobin and TIM-barrel protein [27,28]. The atomic coordinates of the X-ray crystal structures of lysozymes at 1.8-Å resolution were obtained from the Protein Data Bank (PDB) [14,15] and converted into dihedral-angle coordinates using the program FEDER/2 constructed by Wako *et al.* [30]. The following PDB IDs were used: 1HEL (chicken), 1HER (SIS), 1HEO (TVS), 1HEM (TIT), 1HEQ (SIT), 1HEN (TVT), and 1HEP (bobwhite). We defined the volume of a protein with the excluded volume calculated by the method of Higo and Go using a 1.4-Å probe for the radius of water [31].

The energy minimization of the protein structures was carried out in torsion angle space using the program FEDER/2 with the energy function ECEPP/2 and Newton's

method [30]. Conformational energy function E was defined as a sum of five contributions:

$$E = E_c + E_{\text{hyd}} + E_{\text{es}} + E_{\text{tor}} + E_{\text{ss}} \quad (1)$$

where E_c is the van der Waals force (Lennard-Jones type 6-12 potential), E_{hyd} is the hydrogen-bond energy (Lennard-Jones type 10-12 potential), E_{es} is the electrostatic energy, E_{tor} is the torsional energy associated with the barriers to rotation about a rotatable bond, and E_{ss} is the disulfide-bond (loop-closing) energy. Using normal-mode variable σ_i for the i -th harmonic oscillator with angular frequency ω_i , the conformational energy (E) and the mean-square amplitude ($\langle \sigma_i^2 \rangle$) of the equilibrium fluctuation of each normal-mode variable are given by

$$E = (1/2)(\sum \omega_i^2 \sigma_i^2) \quad (2)$$

$$\langle \sigma_i^2 \rangle = \kappa_B T / \omega_i^2 \quad (3)$$

where κ_B is the Boltzmann constant and T is the absolute temperature. In addition to the harmonic approximation of the conformational energy, we assumed a linear volume change (ΔV) due to deviation from the minimum energy conformation with volume V :

$$\Delta V = \sum v_i \sigma_i \quad (4)$$

where v_i is the gradient of the volume with respect to the i -th normal-mode variable at the minimum energy conformation. From Eqs. (3) and (4), the mean-square volume fluctuation $\langle (\Delta V)^2 \rangle$ is given by

$$\langle (\Delta V)^2 \rangle = \kappa_B T \sum (v_i^2 / \omega_i^2) \quad (5)$$

The volume fluctuation is directly related to β_T from statistical thermodynamics [9]:

$$\langle (\Delta V)^2 \rangle = \kappa_B T V \beta_T \quad (6)$$

Thus, isothermal compressibility β_T can be calculated using the following equation [27,28]:

$$\beta_T = \Sigma(v_i^2/\omega_i^2)/V \quad (7)$$

In the present study, β_T was calculated at 300 K.

2.2. Calculation of accessible surface area and cavity volume

The accessible surface area (A_s) and the internal cavities of lysozymes were evaluated from their X-ray structures using the program GRASP (Graphical Representation and Analysis of Structural Properties) [32]. The A_s value was calculated with a probe radius of 1.4 Å, which corresponds to the size of the water molecule. Cavities are analytically defined by the surface described by rolling a probe sphere around an internal atomic surface [33,34]. The total volume (V_{cav}) of cavities is sensitive to the probe size: a probe that is either too large or too small cannot detect small cavities. The cavity is often estimated using probes with radii (R_c) of around 1.2 Å [35,36]. In the present study, the mean values of V_{cav} estimated with four R_c values (1.05, 1.00, 1.15, and 1.20 Å) were used in correlation analysis with compressibility data. The number of internal water molecules in cavities was zero for SIS, TVS, and SIT, one for TIS, TVT, and SVT, and three for TIT. These water molecules were eliminated in the calculation of V_{cav} ; however, its correlation with compressibility was not affected by the presence or absence of internal water molecules in cavities.

3. Results and discussion

3.1. Volume fluctuation and isothermal compressibility

Fig. 2 compares the displacements of C_α atoms of amino-acid residues estimated by normal-mode analysis for two extant lysozymes (TIS and SVT) and two evolutionary intermediates (SIS and SIT) with those estimated from the B-factor of the X-ray structure. The overall features of the C_α displacements were similar in both estimations for all of the lysozymes. The C_α displacements were determined to be smaller by the normal-mode analysis than by the B-factor, which is as expected since normal-mode analysis is carried out for an energy-minimum structure whereas the X-ray structure involves the static disorder of the crystal. The RMSD (root-mean-square deviation) values of C_α displacements for all of the lysozymes was 0.12 Å, indicating that the energy-minimized structures of these intermediates were close to each other and to those of the two extant lysozymes.

(Fig. 2)

Fig. 3 shows the volume changes of the protein molecule from its minimum energy conformation as a function of wave number for extant (TIS and SVT) and two evolutionary (SIS and SIT) lysozymes. Other intermediates showed similar features (data not shown). It is evident from the figure that atom movements faster than 200 cm^{-1} induce no substantial changes in volume, but that slower movements produce larger volume changes of up to 15 \AA^3 . Thus, normal modes with low frequencies play a substantial role in pressure-induced deformations or volume fluctuations of these proteins.

(Fig. 3)

The distributions of volume fluctuations summed within 10-cm^{-1} ranges below 50 cm^{-1} and within the range of $50\text{--}1,000 \text{ cm}^{-1}$ are shown as histograms for all of the lysozymes in Fig. 4. Each lysozyme showed different fluctuation patterns that depended on the rates of movements: extant lysozymes (TIS and SVT) had the largest fluctuation within the range of $10\text{--}20 \text{ cm}^{-1}$, whereas most of the evolutionary intermediates showed enhanced fluctuations for slower movements. The following residues exhibited changes in their mean dihedral angles of over 10° in total mode: residue 129 for TIS (chicken); residues 1 and 127 for SVT (bobwhite); residue 117 for SIS; residues 17, 100, 102, 103, and 129 for SIT; residues 1, 102, 117, and 129 for TIT; residues 41, 67, 71, 102, 117, 120, 121, 126, 127, and 129 for TVT; and residues 1, 48, and 104 for TVS. These results indicate that the local structures are sensitively influenced by the substitution of a single amino acid, especially in evolutionary intermediates. The atomic movements were not analyzed in detail since the local compressibility changes fell outside the scope of the present study.

(Fig. 4)

The calculated values of β_T and $\langle(\Delta V)^2\rangle$ are listed in Table 1 together with V_{cav} and the thermal transition temperature (T_m) reported by Malcolm *et al.* [5]. The β_T values of chicken and bobwhite lysozymes were almost the same (1.61 and 1.59 Mbar^{-1}), but all of the evolutionary mutants showed larger β_T values of up to 2.17 Mbar^{-1} although the β_T value of SIS was similar to that of the extant lysozymes, suggesting that volume

fluctuations are larger for ancestral lysozymes than for extant ones. Volume fluctuations amount to only 0.17–0.20% of the excluded volume, but they induce large variations of up to 16% (39.3–45.7 Å³) among the mutants (Table 1).

(Table 1)

3.2. Comparison of calculated and experimental values of β_T

The β_T value of chicken lysozyme has been experimentally estimated to be 4.7 and 9.8 Mbar⁻¹ from the X-ray crystal structures under high pressures [37,38], and 7.5 Mbar⁻¹ by a high-pressure NMR study [39]. A similar value was estimated from the partial adiabatic compressibility β_s (4.7 Mbar⁻¹) determined from sound-velocity measurements in solution [12]. These experimentally observed β_T and β_s values are considerably larger than the value of 1.61 Mbar⁻¹ obtained in the present study, although direct comparison between the calculated and experimental values of compressibility is intrinsically difficult because the experimental values are partial quantities involving complicated contributions from hydration. Such a low β_T value obtained by normal-mode analysis is not improbable since lysozyme is classified as a rigid protein in the accumulated compressibility data, and a protein should be more rigid in the energy-minimum crystal structure due to suppressed large-scale motions of the molecule [23].

This present study employed a simple computational procedure for the systematic comparison of compressibility among different mutants, but accurate predictions of β_T fell outside the scope of this work. Future studies should address two issues: (1) accurate models of protein dynamics and (2) accurate definitions of protein volumes. Regarding the first of these issues, in the normal-mode analysis *in vacuo*, large-scale molecular movements are associated with low-frequency normal modes. In solution, such movements would be better described by multiple hierarchical modes. For instance, Kitao *et al.* proposed a JAM (jumping-among-minima) model for the energy landscape of human lysozyme in solution at room temperature [40]. Regarding the second issue, the computational estimate of the compressibility depends on the definition of protein volume. Imai *et al.* recently proposed a method to evaluate partial volumes based on a statistical mechanics approach, which provides a powerful computational tool to calculate molecular volumes [41]. In spite of these two issues, it is noteworthy that evolutionary mutants of lysozyme have different β_T values when using the same

calculation procedure, and it is likely that these differences will be magnified in solution.

3.3. Contribution of the cavity volume to compressibility

It is of interest that the evolutionary lysozymes have considerably different β_T values since only small changes are visible in their tertiary structures [6]. Such significant mutation effects have also been observed in the β_s values determined by sound-velocity measurements in solution [18–21]. Two main factors contribute to the partial compressibility of a protein in solution: cavities and hydration [12,13]. In general, cavities contribute positively and hydration contributes negatively to partial compressibility, and hence the experimentally observed partial compressibility is determined by the relative contributions of these two terms. However, the contribution of hydration is not included in normal-mode analysis, and so the observed difference in β_T values of evolutionary lysozymes could be dominantly ascribed to modifications to the cavities. Since the apparent compressibility of a cavity is about tenfold larger than that of water [12], even a small perturbation in the cavity would cause large changes in both β_T and β_s .

Fig. 5 shows typical distributions of the cavities in four lysozyme molecules, as evaluated with an R_c of 1.15 Å. The figure indicates that the number and positions of cavities vary among the mutants. The mean values of V_{cav} estimated with four R_c values (1.05, 1.00, 1.15, and 1.20 Å) are listed in Table 1. It is noteworthy that any mutation induces a large variation in V_{cav} and that all of the evolutionary intermediates have larger V_{cav} values than the extant lysozymes.

(Fig. 5)

Fig. 6 shows plots of β_T against V_{cav} for all of the lysozymes examined, from which it is evident that β_T increases with the total cavity volume (correlation coefficient $r = 0.82$). We observed a similar positive correlation between the β_s and V_{cav} values of 27 globular proteins (unpublished data). Therefore, the cavity volume or atomic packing would be an important factor determining fluctuations in the molecular evolution of lysozyme.

(Fig. 6)

3.4. Relationship of volume fluctuation to evolution and function

According to the neutral theory of molecular evolution [2], the overall properties of evolutionary intermediates should fall within the range exhibited by the extant species. As indicated in Table 1, the values of β_T and $\langle (\Delta V)^2 \rangle$ of the evolutionary mutants fall outside the range of the two wild types, which is negligibly narrow in the present case. This constitutes evidence that volume fluctuations of these lysozymes have followed a nonneutral evolutionary pathway. Such a lack of neutrality was also found for the thermal stability [5]: the T_m values of some intermediates fall outside the range of the two wild types (Table 1). We found a linear positive correlation ($r = 0.67$) between β_T and T_m except for the data for TVS in Table 1. A similar correlation ($r = 0.64$) was previously found between the β_s and T_m values of 14 globular proteins [13]. These results suggest that highly flexible intermediates are tolerant to large thermal fluctuations in the molecular evolution of lysozyme, although the thermodynamic mechanism for this correlation is unclear since the thermal stability of a protein is determined by the change in heat capacity between the native and denatured states.

It is presently unknown how the volume fluctuations are related to the function of these evolutionary lysozymes, because their enzyme activity has not been measured. However, the differences ($\sim 16\%$) in the volume fluctuations among these evolutionary lysozymes would be sufficient to affect the function because volume fluctuations of chicken lysozyme reduce by 3%, 7%, and 14% when it is bound to its inhibitors: monomer, dimer, and trimer of *N*-acetylglucosamine, respectively [42]. In fact, there is a considerable amount of experimental evidence that mutation induces comparable changes in compressibility and enzyme function. Single amino-acid substitutions in aspartate aminotransferase, dihydrofolate reductase, and cyclic AMP receptor protein induce significant changes in β_s , and mutants with larger β_s values exhibit higher enzyme activity [18–20]. Evolutionally designed hyperactive dihydrofolate reductase has a very large β_s value, as predicted from the results obtained for a single mutation [21]. The β_s values of lysozyme and ovalbumin from chicken are smaller than those of the respective proteins from turkey, which is more evolved [43]. These previous results together with those of the present study suggest that proteins become less compressible as they evolve. Ancestral proteins might be more flexible and functionally more active than the extant ones in order to allow them to adapt to diverse environments.

It is probably difficult to confirm the above hypothesis on the fluctuation-related stability and function experimentally because series of evolutionary mutants rarely exist in various proteins. However, the present study has demonstrated that normal-mode

analysis as well as other types of computer simulation can yield useful information on the compressibility and the role of structural fluctuations in the molecular evolution of proteins when evolutionary intermediates are constructed by site-directed mutagenesis and their tertiary structures are resolved. Further computational and experimental studies on the structural fluctuation of evolutionary intermediates would provide new insight into the structure–function relationship of extant proteins.

4. Conclusions

Normal-mode analysis was applied to calculate the β_T values of chicken lysozyme and its six evolutionary mutants. The evolutionary intermediates were more compressible than the extant lysozymes and fell outside the range of the neutral evolutionary pathway. The observed linear correlation between β_T and V_{cav} indicates that the cavity volume or atomic packing is an important factor regulating volume fluctuations in the molecular evolution of proteins. Computer simulations of compressibility would be useful for studying the role of structural fluctuations in the molecular evolution of proteins.

Acknowledgments

The authors thank Mr. S. Ohtao for his technical assistance in compressibility computations. This work was financially supported by a Grant-in-Aid for Scientific Research from the Ministry of Education, Culture, Sports, Science, and Technology of Japan (no. 16041231 to K.G.).

References

- [1] R. Holmquist, beta-Galactosidase and selective neutrality, *Science* 203 (1979) 1012–1014.
- [2] M. Kimura, *The neutral theory of molecular evolution* (Cambridge University Press, Cambridge, 1983).
- [3] M.W. Nachman, S.N. Boyer, C.F. Aquadro, Nonneutral evolution at the mitochondrial NADPH dehydrogenase subunit 3 gene in mice, *Proc. Natl. Acad. Sci. USA* 91 (1994) 6364–6368.
- [4] W.S. Smith, J.R. Hale, C. Neylon, Applying neutral drift to the directed molecular evolution of β -glucuronidase into a β -galactosidase: two different evolutionary pathways lead to the same variant, *BMC Res Notes* 4 (2011) 138.
- [5] B.A. Malcolm, K.P. Wilson, B.W. Matthews, J.F. Kirsch, A.C. Wilson, Ancestral lysozymes reconstructed, neutrality tested, and thermostability linked to hydrocarbon packing, *Nature* 345 (1990) 86–89.
- [6] K.P. Wilson, B.A. Malcolm, B.W. Matthews, Structural and thermodynamic analysis of compensating mutations within the core of chicken egg white lysozyme, *J. Biol. Chem.* 267 (1992) 10842–10849.
- [7] K.M. Helm-Bychowski, A.C. Wilson, Rates of nuclear DNA evolution in pheasant-like birds: evidence from restriction maps, *Proc. Natl. Acad. Sci. USA* 83 (1986) 688–692.
- [8] M. Karplus, J.A. MaCammon, The internal dynamics of globular proteins, *CRC Crit. Rev. Biochem.* 9 (1981) 293–349.
- [9] A. Cooper, Thermodynamic fluctuations in protein molecules, *Proc. Natl. Acad. Sci. USA* 73 (1976) 2740–2741.
- [10] H. Seemann, R. Winter, C.A. Royer, Volume, expansivity and isothermal compressibility changes associated with temperature and pressure unfolding of *Staphylococcal* nuclease, *J. Mol. Biol.* 307 (2001) 1091–1102.
- [11] K. Gekko, M. Araga, T. Kamiyama, E. Ohmae, K. Akasaka, Pressure dependence of the apparent specific volume of bovine serum albumin: insight into the difference between isothermal and adiabatic compressibilities, *Biophys. Chem.* 144 (2009) 67–71.
- [12] K. Gekko, H. Noguchi, Compressibility of globular proteins in water at 25°C, *J. Phys. Chem.* 83 (1979) 2706–2714.
- [13] K. Gekko, Y. Hasegawa, Compressibility-structure relationship of globular proteins, *Biochemistry* 25 (1986) 6563–6571.

- [14] B. Gavish, E. Gratton, C.J. Hardy, Adiabatic compressibility of globular proteins, *Proc. Natl. Acad. Sci. USA* 80 (1983) 750–754.
- [15] B. Nölting, S.G. Sligar, Adiabatic compressibility of molten globules, *Biochemistry* 32 (1993) 12319–12323.
- [16] D.P. Kharakoz, Protein compressibility, dynamics, and pressure, *Biophys. J.* 79 (2000) 511–525.
- [17] N. Taulier, T.V. Chalikian, Compressibilities of protein transitions, *Biochim. Biophys. Acta* 1595 (2002) 48–70.
- [18] K. Gekko, Y. Tamura, E. Ohmae, H. Hayashi, H. Kagamiyama, H. Ueno, A large compressibility change of protein induced by a single amino acid substitution, *Protein Sci.* 5 (1996) 542–545.
- [19] K. Gekko, T. Kamiyama, E. Ohmae, K. Katayanagi, Single amino acid substitutions in flexible loops can induce large compressibility changes in dihydrofolate reductase, *J. Biochem.* 128 (2000) 21–27.
- [20] K. Gekko, M. Obu, J. Li, J.C. Lee, A linear correlation between the energetics of allosteric communication and protein flexibility in the *Escherichia coli* cyclic AMP receptor protein revealed by mutation-induced changes in compressibility and amide hydrogen-deuterium exchange, *Biochemistry* 43 (2004) 3844–3852.
- [21] M. Iwakura, K. Maki, H. Takahashi, T. Takenawa, A. Yokota, K. Katayanagi, T. Kamiyama, K. Gekko, Evolutional design of a hyperactive cysteine- and methionine-free mutant of *Escherichia coli* dihydrofolate reductase, *J. Biol. Chem.* 281 (2006) 13234–13246.
- [22] E. Paci, M. Marchi, Intrinsic compressibility and volume compression in solvated proteins by molecular dynamic simulation at high pressure, *Proc. Natl. Acad. Sci. USA* 93 (1996) 11609–11614.
- [23] F. Tama, O. Miyashita, A. Kitao, N. Go, Molecular dynamics simulation shows large volume fluctuations of proteins, *Eur. Biophys. J.* 29 (2000) 472–480.
- [24] C. Scharnagl, M. Reif, J. Friedrich, Local compressibilities of proteins: comparison of optical experiments and simulations for horse heart cytochrome-c, *Biophys. J.* 89 (2005) 64–75.
- [25] K. Mori, Y. Seki, Y. Yamada, H. Matsumoto, K. Soda, Evaluation of intrinsic compressibility of proteins by molecular dynamics simulation, *J. Chem. Phys.* 125 (2006) 054903.
- [26] N. Smolin, R. Winter, A molecular dynamics simulation of SNase and its hydration shell at high temperature and high pressure, *Biochim. Biophys. Acta* 1764 (2006) 522–534.

- [27] T. Yamato, J. Higo, Y. Seno, N. Go, Conformational deformation in deoxymyoglobin by hydrostatic pressure, *Proteins* 16 (1993) 327–340.
- [28] N. Kobayashi, T. Yamato, N. Go, Mechanical property of a TIM-barrel protein, *Proteins* 28 (1997) 109–116.
- [29] T.J. Kamerzell, J.R. Unruh, C.K. Johnson, C.R. Middaugh, Conformational flexibility, hydration and state parameter fluctuations of fibroblast growth factor-10: effects of ligand binding, *Biochemistry* 45 (2006) 15288–15300.
- [30] H. Wako, S. Endo, K. Nagayama, N. Go, FEDER/2: program for static and dynamic conformational energy analysis of macro-molecules in dihedral angle space, *Comput. Phys. Commun.* 91 (1995) 233–251.
- [31] J. Higo, N. Go, Algorithm for rapid calculation of excluded volume of large molecules, *J. Comput. Chem.* 10 (1989) 376–379.
- [32] A. Nicholls, K.A. Sharp, B. Honig, Protein folding and association: insights from the interfacial and thermodynamic properties of hydrocarbons, *Proteins* 11 (1991) 281–296.
- [33] A. Shrake, J.A. Rupley, Environment and exposure to solvent of protein atoms. Lysozyme and insulin, *J. Mol. Biol.* 79 (1973) 351–371.
- [34] F.M. Richards, Calculation of molecular volumes and areas for structures of known geometry, *Methods Enzymol.* 115 (1985) 440–464.
- [35] A.A. Rashin, M. Iofin, B. Honig, Internal cavities and buried waters in globular proteins, *Biochemistry* 25 (1986) 3619–3625.
- [36] S.J. Hubbard, P. Argos, A functional role for protein cavities in domain:domain motions, *J. Mol. Biol.* 261 (1996) 289–300.
- [37] C.M. Kundrot, F.M. Richards, Crystal structure of hen egg-white lysozyme at a hydrostatic pressure of 1000 atmospheres, *J. Mol. Biol.* 193 (1987) 157–170.
- [38] R. Fourme, R. Kahn, M. Mezouar, E. Girard, C. Hoerentrup, T. Prangé, I. Ascone, High-pressure protein crystallography (HPPX): instrumentation, methodology and results on lysozyme crystals, *J. Synchrotron Radiat.* 8:5 (2001) 1149–1156.
- [39] M. Refaee, T. Tezuka, K. Akasaka, M.P. Williamson, Pressure-dependent changes in the solution structure of hen egg-white lysozyme, *J. Mol. Biol.* 327 (2003) 857–865.
- [40] A. Kitao, S. Hayward, N. Go, Energy landscape of a native protein: jumping-among-minima model, *Proteins* 33 (1998) 496–517.
- [41] T. Imai, A. Kovalenko, F. Hirata, Partial molar volume of proteins studied by the three-dimensional reference interaction site model theory, *J. Phys. Chem. B*

109 (2005) 6658–6665.

[42] K. Gekko, K. Yamagami, Compressibility and volume changes of lysozyme due to inhibitor binding, *Chem. Lett.* 27 (1998) 839–840.

[43] K. Gekko, K. Yamagami, Flexibility of food proteins as revealed by compressibility, *J. Agric. Food Chem.* 39 (1991) 57–62.

Figure Captions

Fig. 1. Assumed evolutionary pathways between two extant lysozymes (chicken and bobwhite): TIS (chicken) \rightarrow SIS, TIT, TVS (single mutation) \rightarrow SIT, SVS, TVT (double mutation) \rightarrow SVT (bobwhite).

Fig. 2. Comparisons of displacements of C_{α} atoms of amino-acid residues in extant (TIS and SVT) and evolutionary (SIS and SIT) lysozymes estimated by normal-mode analysis (\circ) and from the X-ray structure (B-factor, \blacksquare).

Fig. 3. Volume changes from the energy-minimum conformation of extant (TIS and SVT) and evolutionary (SIS and SIT) lysozymes as functions of the wave number for the normal mode.

Fig. 4. Distribution of volume fluctuations of lysozymes within 10-cm^{-1} ranges below 50 cm^{-1} and within the range of $50\text{--}1,000\text{ cm}^{-1}$.

Fig. 5. Distribution of cavities in extant (TIS and SVT) and evolutionary (SIS and SIT) lysozymes. Cavities were evaluated with a probe radius of 1.15 \AA and are indicated by gray balls.

Fig. 6. Plot of β_T against V_{cav} for lysozymes. Solid line represents the least-squares linear regression. V_{cav} is the mean value estimated with four probe radii (1.05 , 1.00 , 1.15 , and 1.20 \AA).




Fig. 1.




Fig. 2




Fig. 3







Fig. 4




TIS (Chicken)



SIS



SIT



SVT (Bobwhite)

Fig. 5




Fig. 6

Table 1

Compressibilities and volume fluctuations of extant and evolutionary lysozymes.

Lysozyme	PDB ID	β_T (Mbar ⁻¹)	$\langle(\Delta V)^2\rangle$ ^{a)} (Å ³)	V_{cav} ^{b)} (Å ³)	T_m ^{c)} (°C)
TIS (Chicken)	1HEL	1.61	39.3 (0.17)	65.8	73.9
SIS	1HER	1.61	39.3 (0.17)	100.8	73.0
TVS	1HEO	1.98	44.0 (0.19)	119.2	71.2
TIT	1HEM	2.08	44.8 (0.19)	211.2	77.5
SIT	1HEQ	1.75	41.1 (0.18)	143.0	75.5
TVT	1HEN	2.17	45.7 (0.20)	163.5	74.5
SVT (Bobwhite)	1HEP	1.59	39.4 (0.17)	95.0	73.4

a) Values within parentheses represent ratios (%) of $\langle(\Delta V)^2\rangle$ to the total molecular volume.

b) Mean total cavity volumes estimated with probe radii of 1.05–1.20 Å.

c) Taken from Ref. 5.