

広島大学学術情報リポジトリ
Hiroshima University Institutional Repository

Title	Replication study of the association of SNPs in the LHX3-QSOX2 and IGF1 loci with adult height in the Japanese population; wide-ranging comparison of each SNP genotype distribution
Author(s)	Fujihara, Junko; Takeshita, Haruo; Kimura, Kaori; Kataoka, Kaori; Soejima, Mikiko; Koda, Yoshiro; Yuasa, Isao; Iida, Reiko; Ueki, Misuzu; Nagao, Masataka; Kominato, Yoshihiko; Yasuda, Toshihiro
Citation	Legal Medicine , 14 (4) : 205 - 208
Issue Date	2012
DOI	10.1016/j.legalmed.2012.02.001
Self DOI	
URL	https://ir.lib.hiroshima-u.ac.jp/00034824
Right	(c) 2012 Elsevier Ireland Ltd. All rights reserved.
Relation	



Manuscript Number:

Title: Replication study of the association of SNPs in the LHX3-QSOX2 and IGF1 loci with adult height in the Japanese population; wide-ranging comparison of each SNP genotype distribution

Article Type: Brief Communication

Keywords: Ethnic difference; Height; IGF1; LHX3-QSOX2; Personal identification; SNPs

Corresponding Author: Prof. Haruo Takeshita, M.D. ph.D

Corresponding Author's Institution: Shmane University School of Medicine

First Author: Junko Fujihara

Order of Authors: Junko Fujihara; Haruo Takeshita, M.D. ph.D

Abstract: Adult height is a highly heritable trait involving multiple genes. Recent genome-wide association studies have identified that SNP rs12338076 in the LHX3-QSOX2 locus, and rs1457595 and rs17032362 in the IGF1 locus are associated with human height in the Japanese population (Okada et al. Hum Mol Genet 2010; 19:2303-12). We performed a replication study to examine the associations between these 3 SNPs and adult height in the Japanese population based on autopsy cases. However, it was not possible to confirm that all these SNPs influenced adult height in the study population. We first conducted a wide-ranging survey of these 3 SNPs in the above genes using 9 different populations including Asians, Africans and Caucasians, and demonstrated that the genotypes of rs12338076 and rs17032362 were distributed in an ethnicity-dependent manner; even within Asian populations, the genotype distributions of the SNPs differed widely. Although there are differences in height distribution between different populations, possibly due to genetic factors and/or gene-environmental interactions, the contradictory results of the association study and ethnic differences in genotype distribution allow us to assume that these height-related SNPs in the genes may contribute to adult height to a slight extent, at least in the Japanese population. It is anticipated that the present information will be useful for developing a reliable tool for personal identification through elucidation of the genetic basis of human height.

27 October 2011

To: Dr. N. Ikeda
Editor-in-Chief
Legal Medicine

Dear Dr. N. Ikeda

Please find the manuscript entitled “**Replication study of the association of SNPs in the LHX3-QSOX2 and IGF1 loci with adult height in the Japanese population; wide-ranging comparison of each SNP genotype distribution**”

We would be most grateful if the manuscript could be reviewed and considered for publication in *Legal Medicine* as Brief Communication.

Each of us has made a significant contribution to the design, execution, analysis and writing up of this study. All the authors and authorities in the institute have approved submission of this manuscript. We swear that materials have not been previously reported, and that it is not consideration for publication elsewhere.

We look forward to your favorable response.

Sincerely yours,

Haruo TAKESHITA, MD, PhD (corresponding author)

Department of Legal Medicine
Shimane University School of Medicine,
Enya 89-1 Izumo, 693-8501, Japan
Tel: +81-853-20-2156
Fax: +81-853-20-2155
E-mail: htakeshi@med.shimane-u.ac.jp

1 (Brief communication)

2

3 **Replication study of the association of SNPs in the *LHX3-QSOX2***
4 **and *IGF1* loci with adult height in the Japanese population;**
5 **wide-ranging comparison of each SNP genotype distribution**

6 Junko Fujihara^a, Haruo Takeshita^{a*}, Kaori Kimura-Kataoka^a, Mikiko Soejima^b, Yoshiro
7 Koda^b, Isao Yuasa^c, Reiko Iida^d, Misuzu Ueki^e, Masataka Nagao^f, Yoshihiko Kominato^g,
8 Toshihiro Yasuda^e

9

10 ^a *Department of Legal Medicine, Shimane University School of Medicine*

11 *89-1 Enya, Izumo, Shimane 693-8501, Japan*

12 ^b *Department of Forensic Medicine and Human Genetics, Kurume University School of*
13 *Medicine, Fukuoka, Japan*

14 ^c *Division of Legal Medicine, Faculty of Medicine, Tottori University, Tottori, Japan*

15 ^d *Division of Life Sciences and* ^e *Division of Medical Genetics and Biochemistry, Faculty of*
16 *Medical Sciences, University of Fukui, Fukui, Japan*

17 ^f *Department of Legal Medicine, Graduate School of Biomedical Sciences, Hiroshima*
18 *University, Hiroshima, Japan*

19 ^g *Department of Legal Medicine, Gunma University School of Medicine, Maebashi, Japan*

20

21

22 *Corresponding author: Haruo Takeshita, M.D., Ph.D.

23 E-mail: htakeshi@med.shimane-u.ac.jp (H. Takeshita)

24

(Abstract)

Adult height is a highly heritable trait involving multiple genes. Recent genome-wide association studies have identified that SNP rs12338076 in the *LHX3-QSOX2* locus, and rs1457595 and rs17032362 in the *IGF1* locus are associated with human height in the Japanese population (Okada et al. Hum Mol Genet 2010; 19:2303–12). We performed a replication study to examine the associations between these 3 SNPs and adult height in the Japanese population based on autopsy cases. However, it was not possible to confirm that all these SNPs influenced adult height in the study population. We first conducted a wide-ranging survey of these 3 SNPs in the above genes using 9 different populations including Asians, Africans and Caucasians, and demonstrated that the genotypes of rs12338076 and rs17032362 were distributed in an ethnicity-dependent manner; even within Asian populations, the genotype distributions of the SNPs differed widely. Although there are differences in height distribution between different populations, possibly due to genetic factors and/or gene-environmental interactions, the contradictory results of the association study and ethnic differences in genotype distribution allow us to assume that these height-related SNPs in the genes may contribute to adult height to a slight extent, at least in the Japanese population. It is anticipated that the present information will be useful for developing a reliable tool for personal identification through elucidation of the genetic basis of human height.

Key words: Ethnic difference/Height /IGF1/LHX3-QSOX2 / Personal identification/SNPs

1. Introduction

In crime cases with no known suspect and no DNA database match, information on externally visible traits such as gender, eye and hair color, and height provided by DNA-based investigations would be valuable for forensic personal identification [1]. Among them, height is under strong genetic influence, with an estimated heritability of up to 80-90% [2, 3]. Recent genome-wide association studies conducted using single nucleotide polymorphisms (SNPs) have revealed many genetic loci responsible for human height: approximately 50 genetic variants [4-6]. Among height-related genes, 3 SNPs (rs1047275, rs7868682, and rs7968902) in the high mobility group-A2 (HMGA2) gene have been shown to be associated with height in Caucasians [7]. Furthermore, we have confirmed the association between rs1042725 in *HMGA2* and adult height in Japanese autopsy cases [8]. Recently, Okada et al. have demonstrated that SNP rs12338076 in the LIM homeobox 3 (*LHX3*)-quiescin Q6 sulfhydryl oxidase 2 (*QSOX2*) locus, and rs1457595 and rs17032362 in the insulin-like growth factor 1 (*IGF1*) locus significantly influence adult height in Japanese subjects [9]. However, replication studies of genetic associations in independent populations, even those belonging to the same ethnic group, are indispensable for evaluating the role of related genes in complex traits such as height [7].

In this context, the first aim of the present study was to examine whether 3 SNPs, rs12338076 in *LHX3-QSOX2*, and rs1457595 and rs17032362 in *IGF1*, could be associated with adult height in Japanese autopsy samples as a replication study. As genetic and environmental backgrounds for height vary between different ethnic groups, our second aim was to clarify whether the distributions of the SNPs vary among different populations, and therefore we performed a wide-ranging population study of these 3 SNPs in 9 different populations, including Asians, Africans and Caucasians.

2. Materials and methods

2.1. Biological samples

1 Genomic DNA was extracted from blood or bloodstain using a QIAamp DNA mini kit
2 (QIAGEN Inc., Chatsworth, CA) according to the manufacture's instruction. Samples
3 collected randomly from healthy and unrelated subjects derived from 8 different populations:
4 251 Koreans (Seoul, Korea), 36 Mongolians (Ulaanbaatar, Mongolia), 70 Tibetans (Katmandu,
5 Nepal), 31 Sinhalese (Kandy, Sri Lanka), 71 Germans (Munchen, Germany), 210 Turks
6 (Adana area, Turk), 41 Ovambos (Bantus, Namibia), and 41 Xhosas (Cape Town of South
7 Africa). The study including usage of genomic DNA derived from autopsy cases was
8 reviewed and approved by the Human Ethics Committee of Shimane University School of
9 Medicine. This study was explained in writing to participants and written consent was
10 obtained; since a part of participants had been without informed consent, usage of them as
11 anonymous samples was approved by Ethics Committee of the Kurume University.

12 The height of 102 Japanese subjects (65 male and 35 female; mean age 57.5 years; mean
13 height 160.1 cm) autopsied in Shimane prefecture was measured. Subjects less than 18 year
14 old were excluded. Genomic DNA was extracted from blood samples of these subjects
15 collected at autopsy using a QIAamp DNA mini kit.

16 17 2.2. SNP typing

18 In this study, all 3 SNPs were analyzed by polymerase chain reaction (PCR) followed by
19 restriction fragment length polymorphism (RFLP) analysis [10]. Among them, we used a
20 mismatched PCR-amplification method for genotyping SNPs (rs1457595 and rs17032362)
21 [11]. Incorporation of a deliberate mismatch close to the 3'-terminus of a PCR primer
22 allowed the creation of each enzyme recognition site. Primers for the specific amplification
23 of the DNA fragments encompassing a substitution site corresponding to the SNPs were
24 newly designed on the basis of the nucleotide sequence of the LHX3-QSOX2 (NM_181701)
25 and IGF1 (AY260957) genes; the following primer sets were used: 12338076-F,
26 CTTCCCGTAGATCTTGGTACAAGGTGGC, and 12338076-R,
27 CCTCCCAAAGTGCTAGGATTACAGGCGTGA, for rs12338076; 1457595-R,

1 GGAGGTTGATAGAGTAGG, and 1457595-R,
2 GCTGTGTTGGTTGGCTGGCAAATGATCGT, for rs1457595; 17032362-F,
3 CCAAATACAGCCAACAGCAACTTTC, and 17032362-R,
4 TACGTGGCATTGGCTTAGGAGCCTG, for rs17032362. The underlined residues indicate
5 the mismatched nucleotide. Amplification was performed in a 25- μ l reaction mixture using
6 approximately 2 ng of DNA. The reaction mixture contained a buffer (15 mM Tris-HCl, pH
7 8.0, 50 mM KCl), 1.5 mM MgCl₂, 0.5 μ M of each primer, 200 μ M dNTPs, and 1.25 U of Taq
8 polymerase (AmpliTaq Gold; Applied Biosystems, Foster City, CA). Two μ l of the PCR
9 product obtained using each pair of primers was digested with each enzyme (New England
10 Biolabs, Ipswich, MA, USA): *Hha I* for rs12338076, *HpyI88III* for rs1457595 and
11 rs17032362 according to the manufacture's instruction. The digests were separated in an 8%
12 polyacrylamide gel, and the patterns on the gels were visualized by silver staining as
13 described previously [8]. Nucleotide sequences of the representative subjects were
14 confirmed by direct sequencing of the PCR products in which a substitution site
15 corresponding to each SNP was included; the dideoxy chain-terminating method with the
16 BigDye[®] Terminator Cycle Sequencing kit was employed using a Genetic Analyzer 310
17 (Applied Biosystems) according to the manufacture's instruction.

19 2.3. Statistical analysis

20 χ^2 -analysis was performed to evaluate the Hardy-Weinberg equilibrium. Single
21 regression analysis was used to compare adult height among different genotypes using the
22 program STATCEL2 (OMS Publishing, Inc.). Differences at $p < 0.01$ were considered to be
23 statistically significant.

25 3. Results and discussion

26 A DNA fragment containing a substitution site at each SNP position in *LHX3-QSOX2* and
27 *IGF1* was separately amplified using a set of primers and subjected to digestion with each

1 enzyme. In rs12338076, the product amplified from the A-allele was completely digested
2 with *Hha I* to yield fragments of 149 and 37 bp, whereas that from the G-allele yielded neither
3 of these fragments. The same procedure was employed for the other 2 SNPs. The validity
4 of the genotyping results obtained by these methods was confirmed by sequencing analyses of
5 genomic DNA derived from several representative subjects.

6 First, in order to evaluate the association of the SNPs, rs12338076 in *LHX3-QSOX2*, and
7 rs1457595 and rs17032362 in *IGF1*, with adult height variation, we examined both the
8 genotype of each SNP and height in autopsied Japanese subjects ($n = 102$). As shown in Fig.
9 1, no significant association between the genotypes in each SNP and height was found.
10 Furthermore, linear regression analysis assuming an additive inheritance model also
11 demonstrated no significant association of any of the 3 SNPs with height. Therefore, these
12 findings allow us to conclude that all of the SNPs, rs12338076 in *LHX3-QSOX2*, and
13 rs1457595 and rs17032362 in *IGF1*, did not influence adult height in the Japanese population
14 examined.

15 Next, worldwide distribution analysis of the 3 SNPs (rs12338076, rs1457595 and
16 rs17032362) was performed in 5 Asian, 2 African and 2 Caucasian populations. The
17 genotype distributions and the allele frequencies determined in this study are summarized in
18 Table 1. These genotype distributions were in Hardy-Weinberg equilibrium in all the
19 populations examined. In the SNP rs12338076 (*LHX3-QSOX2*), the C-allele was
20 predominant in African populations, and showed the highest frequency. In contrast, the
21 A-allele was predominant in Caucasian and Asian populations. The C-allele of the SNP
22 rs1457595 (*IGF1*) was predominant in all of the populations; the minor A-allele was not
23 found in Germans. In the SNP rs17032362 (*IGF1*), the G-allele was not observed in
24 Caucasian and African populations, being mono-allelic. Among Asian populations, the
25 G-allele was predominant. Therefore, among the 3 SNPs, rs12338076 and rs17032362 were
26 confirmed to show an ethnicity-dependent pattern in its genotype distribution. Furthermore,
27 these findings indicate that Asians exhibit wide genetic diversity with regard to the 3 SNPs

1 among different populations. In contrast, 3 SNPs (rs1042725, rs7968902 and rs7868682) in
2 the HMGA2 gene related to adult height showed a similar genotype distribution among Asian
3 populations, including the same ones examined in this study [8]. Also, the genotype
4 distributions of each of the 3 SNPs in this study were found to be consistent with those of
5 HapMap studies (data not shown).

6 Okada et al. have reported a novel association of *LHX3-QSOX2* loci with height in the
7 Japanese population [9]. They have also demonstrated an association of the *IGF1* locus with
8 height. The LHX3, together with LHX4, is a homeodomain transcription factor playing
9 essential roles in pituitary gland and nerve system development. Mutations in the LHX3 and
10 LHX4 genes have been identified in patients with complex and variable syndromes involving
11 short stature, and with combined pituitary hormone deficiency diseases [12]. IGF1 belongs
12 to a family of proteins involved in growth and development, and especially mediates several
13 growth-promoting effects of growth hormones [13, 14]. With regard to association of the
14 *IGF1* locus with height, similar results have been replicated in a Korean population [15]. In
15 contrast, no significant association between IGF1 polymorphism and adult height was found
16 in Caucasians [16-22]. In the present study, we were unable to confirm a significant
17 association of all of the 3 SNPs, rs12338076 in *LHX3-QSOX2*, and rs2457595 and
18 rs17032362 in *IGF1*, with adult height in our study population. Contradictory results
19 derived from different populations, even those belonging to the same ethnic group, might be
20 attributable to differences in the effect size of SNPs among populations, differing frequencies
21 of the causative alleles, study population sample size, or gene-gene or gene-environment
22 interactions. The present study is the first to have conducted a worldwide population survey
23 of height-related SNPs in the *LHX3-QSOX2* and *IGF1* loci. For rs17032362 in *IGF1* loci,
24 the G-allele was not observed in Caucasian or African populations. Even within Asian
25 populations, the genotype distributions of the 3 SNPs (rs12338076, rs1457595 and
26 rs17032362) were different. Although there are differences in the distribution of height
27 between different populations, possibly due to genetic factors and/or gene-environmental

1 interactions, these findings allow us to assume that these height-related SNPs in the above
2 genes may influence adult height to a slight extent, at least in the Japanese population.

3 From a forensic perspective, the effective use of short tandem repeats has made it possible
4 to perform personal identification with extremely high accuracy. However, in order to
5 perform this type of identification, control DNA samples from the individuals concerned are
6 indispensable. In this context, DNA markers associated with physical traits such as eye, hair
7 and skin color, or height, have a distinct advantage in that control DNA samples are not
8 required, making it possible to narrow down and group individuals with connections to
9 forensic scenarios. It is plausible that SNPs associated with height would be applicable as
10 one such type of DNA marker. In order to apply height-related SNPs for forensic personal
11 identification in a practical setting, cumulative investigations of the genetic background of
12 human height will be required. Therefore, it is anticipated that the present findings will
13 provide information useful for developing a reliable tool for personal identification through
14 elucidation of the genetic basis of human height.

15 16 **Acknowledgement**

17 *DNA samples of bloodstain samples of the Ovambo and Turkish populations were kindly provided*
18 *by Dr. B. Brinkmann. Blood samples of Korean and Mongolian populations were kindly provided by*
19 *Dr. K. Shiwaku. This study was supported in part by Grants-in-Aids for Scientific Research*
20 *(21659175 to HT and 22249023 to TY) from the Japan Society for the Promotion of Science.*

References

- [1] Kayser M, Schneider PM (2009) DNA-based prediction of human externally visible characteristics in forensics: Motivations, scientific challenges, and ethical considerations. *Forensic Sci Int Genet* 3: 154–161.
- [2] Silventoinen K, Kaprio J, Lahelma E, Koskenvuo M. Relative effect of genetic and environmental factors on body height: differences across birth cohorts among Finnish men and women. *Am J Public Health* 2000; 90: 627–30.
- [3] Macgregor S, Cornes BK, Martin NG, Visscher PM. Bias, precision and heritability of self-reported and clinically measured height in Australian twins. *Hum Genet* 2006; 120: 571–80.
- [4] Weedon MN, Frayling TM. Reaching new heights: insights into the genetics of human stature. *Trends in Genet* 2008; 24: 595–603.
- [5] Lettre G, Jackson AU, Gieger C, Schumacher F, Berndt, SI, Sanna S, et al. Identification of ten loci associated with height highlights new biological pathways in human growth. *Nat Genet* 2008; 40: 584–91.
- [6] Weedon MN, Lango H, Lindgren CM, Wallace C, Evans DM, Mangino M, et al. Genome-wide association analysis identifies 20 loci that influence adult height. *Nat Genet* 2008; 40: 575–83.
- [7] Yang, TL, Guo Y, Zhang LS, Tian Q, Yan H, Guo Y F, et al. *HMGA2* is confirmed to be associated with human adult height. *Ann Hum Genet* 2010; 74: 11–6.
- [8] Takeshita H, Fujihara J, Soejima M, Koda Y, Kimura-Kataoka K, Ono R, et al. Confirmation that SNPs in the high mobility group-A2 gene (*HMGA2*) are associated with adult height in the Japanese population; wide-ranging population survey of height-related SNPs in *HMGA2*. *Electrophoresis* 2011; 32:1844–51.
- [9] Okada Y, Kamatani Y, Takahashi A, Matsuda K, Hosono N, Ohmiya H, et al. A genome-wide association study in 19 633 Japanese subjects identified *LHX3-QSOX2* and *IGF1* as adult height loci. *Hum Mol Genet* 2010; 19:2303–12.

- 1 [10] Kumar R, Dunn LL. Designed diagnostic restriction fragment length polymorphisms for
2 the detection of point mutations in ras oncogenes. *Oncogene Res* 1989; 4: 235–41.
- 3 [11] Yasuda T, Nadano D, Tenjo E, Takeshita H, Sawazaki K, Nakanaga M, et al. Genotyping
4 of human deoxyribonuclease I polymorphism by the polymerase chain reaction.
5 *Electrophoresis* 1995; 16: 1889–93.
- 6 [12] Mullen RD, Colvin SC, Hunter CS, Savage JJ, Walvoord EC, Bhangoo APS, et al. Roles
7 of the LHX3 and LHX4 LIM-homeodomain factors in pituitary development. *Mol*
8 *Cell Endocrinol* 2007; 265-266: 190–5.
- 9 [13] Clemmons DR. Modifying IGF1 activity: an approach to treat endocrine disorders,
10 atherosclerosis and cancer. *Nat. Rev. Drug. Discov* 2007; 6: 821–33.
- 11 [14] Kemp SF. Insulin-like growth factor-I deficiency in children with growth hormone
12 insensitivity: current and future treatment options. *BioDrugs* 2009; 23: 155–63.
- 13 [15] Kim JJ, Lee HI, Park T, Kim K, Lee JE, Cho NH, et al. Identification of 15 loci
14 influencing height in a Korean population. *J Hum Genet* 2010; 55, 27–31.
- 15 [16] Weedon MN., Lettre G., Freathy, RM., Lindgren, CM., Voight, BF., Perry, JR, et al. A
16 common variant of HMGA2 is associated with adult and childhood height in the general
17 population. *Nat Genet* 2007; 39, 1245–50.
- 18 [17] Sanna S, Jackson AU, Nagaraja R, Willer C, Chen WM, Bonnycastle LL, Set al.
19 Common variants in the GDF5-UQCC region are associated with variation in human
20 height. *Nat Genet* 2008; 40: 198–203.
- 21 [18] Weedon MN, Lango H, Lindgren CM, Wallace C, Evans DM, Mangino M, et al. (2008)
22 Genome-wide association analysis identifies 20 loci that influence adult height. *Nat*
23 *Genet* 2008; 40, 575–83.
- 24 [19] Lettre G, Jackson AU, Gieger C, Schumacher FR, Berndt SI, Sanna S, et al.
25 Identification of ten loci associated with height highlights new biological pathways in
26 human growth. *Nat Genet* 2008; 40, 584–91.
- 27 [20] Gudbjartsson DF, Walters GB, Thorleifsson G, Stefansson H, Halldorsson BV,

1 Zusmanovich P, Sulem P, et al. Many sequence variants affecting diversity of adult
2 human height. *Nat Genet* 2008; 40, 609–15.

3 [21] Johansson A, Marroni F, Hayward C, Franklin, CS, Kirichenko AV, Jonasson I, et al.
4 Common variants in the JAZF1 gene associated with height identified by linkage and
5 genome-wide association analysis. *Hum Mol Genet* 2009; 18, 373–80.

6 [22] Soranzo N, Rivadeneira F, Chinappen-Horsley U, Malkina I, Richards JB, Hammond N,
7 et al. Meta-analysis of genome-wide scans for human adult stature identifies novel Loci
8 and associations with measures of skeletal frame size. *PLoS Genet* 2009; 5, e1000445.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 (Figure legend)

2

3 **Fig. 1.** Comparison of height among Japanese autopsied subjects with different genotypes
4 of 3 SNPs, rs12338076 in *LHX3-QSOX2*, and rs2457595 and rs17032362 in *IGF1*. The
5 height is expressed as mean \pm standard deviation; the bar represents the standard deviation.
6 P-values were calculated by single regression analysis.

7

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Table 1

Genotype distribution and allele frequencies (95% CI) of SNPs of *IGF1* and *LHX3-QSOX2*

Population	rs12338076				rs1457595				rs17032362			
	Genotype, <i>n</i> (%)		Allele frequency		Genotype, <i>n</i> (%)		Allele frequency		Genotype, <i>n</i> (%)		Allele frequency	
Japanese (<i>n</i> = 102)	CC	7 (6.9)	C	0.275	CC	77 (75.5)	C	0.853	AA	18 (17.6)	A	0.324
	CA	42 (41.1)		(0.214 - 0.336)	CA	20 (19.6)		(0.804 - 0.902)	AG	30 (29.4)		(0.260 - 0.388)
	AA	53 (52.0)	A	0.725	AA	5 (4.9)	A	0.147	GG	54 (53.0)	G	0.676
				(0.664-0.786)				(0.098 - 0.196)				(0.612 - 0.740)
Korean (<i>n</i> = 251)	CC	34 (13.6)	C	0.365	CC	182 (72.5)	C	0.837	AA	41 (16.3)	A	0.311
	CA	115 (45.8)		(0.323 - 0.407)	CA	56 (22.3)		(0.805 - 0.869)	AG	74 (29.5)		(0.271 - 0.351)
	AA	102 (40.6)	A	0.635	AA	13 (5.2)	A	0.163	GG	136 (54.2)	G	0.689
				(0.593 - 0.677)				(0.131 - 0.195)				(0.649 - 0.729)
Mongolians (<i>n</i> = 36)	CC	5 (13.9)	C	0.375	CC	20(55.6)	C	0.708	AA	1 (2.8)	A	0.097
	CA	17 (47.2)		(0.263 - 0.487)	CA	11 (30.6)		(0.603 - 0.813)	AG	5 (13.9)		(0.029 - 0.165)
	AA	14 (38.9)	A	0.625	AA	5 (13.9)	A	0.292	GG	30 (83.3)	G	0.903
				(0.513- 0.737)				(0.187 - 0.397)				(0.835 - 0.971)
Tibetans (<i>n</i> = 70)	CC	4 (5.7)	C	0.207	CC	47 (67.1)	C	0.779	AA	15 (21.4)	A	0.413
	CA	21 (30.0)		(0.113 - 0.301)	CA	15 (21.4)		(0.683 - 0.875)	AG	29 (37.2)		(0.299 - 0.527)
	AA	45 (54.3)	A	0.793	AA	8 (11.4)	A	0.221	GG	26 (41.4)	G	0.587
				(0.699 - 0.887)				(0.125 - 0.317)				(0.473 - 0.701)
SriLanka Sinhalese (<i>n</i> = 31)	CC	1 (3.2)	C	0.097	CC	21 (67.7)	C	0.742	AA	2 (6.5)	A	0.258
	CA	4 (12.9)		(0.023 - 0.171)	CA	4 (12.9)		(0.633 - 0.851)	AG	12 (38.7)		(0.149 - 0.367)
	AA	26 (83.9)	A	0.903	AA	6(19.4)	A	0.258	GG	17 (54.8)	G	0.742
				(0.829 - 0.977)				(0.149 - 0.367)				(0.633 - 0.851)
German (<i>n</i> = 71)	CC	5 (7.0)	C	0.239	CC	71(100)		1.000	AA	71(100)		1.000
	CA	24 (33.8)		(0.169-0.309)	CA	0 (0)			AG	0 (0)		
	AA	42 (59.2)	A	0.761	AA	0 (0)	A	0.000	GG	0 (0)	G	0.000
				(0.691-0.831)								
Turks (<i>n</i> = 210)	CC	26 (12.4)	C	0.350	CC	202 (96.2)	C	0.981	AA	210 (100)	A	1.000
	CA	95 (45.2)		(0.304 - 0.396)	CA	8 (3.8)		(0.968 - 0.994)	AG	0 (0)		
	AA	89 (42.4)	A	0.650	AA	0 (0.0)	A	0.019	GG	0 (0)	G	0.000
				(0.604 - 0.696)				(0.006 - 0.032)				
Ovambos (<i>n</i> = 41)	CC	29 (70.7)	C	0.817	CC	37 (90.2)	C	0.951	AA	41 (100)	A	1.000
	CA	9 (22.0)		(0.733-0.901)	CA	4 (9.8)		(0.905 - 0.997)	AG	0 (0)		
	AA	3 (7.3)	A	0.183	AA	0 (0.0)	A	0.048	GG	0 (0)	G	0.000
				(0.099-0.267)				(0.002 - 0.094)				
Xhosans (<i>n</i> = 41)	CC	32 (78.1)	C	0.878	CC	18 (43.9)	C	0.659	AA	41 (100)	A	1.000
	CA	8 (19.5)		(0.807-0.949)	CA	18 (43.9)		(0.556 - 0.762)	AG	0 (0)		
	AA	1 (2.4)	A	0.122	AA	5 (12.2)	A	0.341	GG	0 (0)	G	0.000
				(0.051 - 0.193)				(0.238 - 0.444)				

Figure(s)

