

生物圏科学  
Biosphere Sci.  
52 : 87–115 (2013)

**A synopsis of the parasites from cyprinid fishes of the genus *Tribolodon*  
in Japan (1908-2013)**

Kazuya Nagasawa and Hirotaka Katahira

Graduate School of Biosphere Science, Hiroshima University

Published by

The Graduate School of Biosphere Science

Hiroshima University

Higashi-Hiroshima 739-8528, Japan

December 2013

## REVIEW

### A synopsis of the parasites from cyprinid fishes of the genus *Tribolodon* in Japan (1908-2013)

Kazuya Nagasawa<sup>1)\*</sup> and Hiroataka Katahira<sup>1,2)</sup>

<sup>1)</sup> Graduate School of Biosphere Science, Hiroshima University,  
1-4-4 Kagamiyama, Higashi-Hiroshima, Hiroshima 739-8528, Japan

<sup>2)</sup> Present address: Graduate School of Environmental Science, Hokkaido University,  
N10 W5, Sapporo, Hokkaido 060-0810, Japan

**Abstract** Four species of the cyprinid genus *Tribolodon* occur in Japan: big-scaled redbfin *T. hakonensis*, Sakhalin redbfin *T. sachalinensis*, Pacific redbfin *T. brandtii*, and long-jawed redbfin *T. nakamuraii*. Of these species, *T. hakonensis* is widely distributed in Japan and is important in commercial and recreational fisheries. Two species, *T. hakonensis* and *T. brandtii*, exhibit anadromy. In this paper, information on the protistan and metazoan parasites of the four species of *Tribolodon* in Japan is compiled based on the literature published for 106 years between 1908 and 2013, and the parasites, including 44 named species and those not identified to species level, are listed by higher taxon as follows: Ciliophora (2 named species), Myxozoa (1), Trematoda (18), Monogenea (0), Cestoda (3), Nematoda (9), Acanthocephala (2), Hirudinida (1), Mollusca (1), Branchiura (0), Copepoda (6), and Isopoda (1). For each taxon of parasite, the following information is given: its currently recognized scientific name, previous identification used for the parasite occurring in or on *Tribolodon* spp.; habitat (freshwater, brackish, or marine); site(s) of infection within or on the host; known geographical distribution in Japan; and the published source of each locality record. The highest number of the named parasite species was recorded from *T. hakonensis* (43), followed by *T. sachalinensis* (7), *T. brandtii* (6), and *T. nakamuraii* (1).

**Key words:** checklist, parasites, *Tribolodon brandtii*, *Tribolodon hakonensis*, *Tribolodon nakamuraii*, *Tribolodon sachalinensis*

## INTRODUCTION

Four species of the genus *Tribolodon* Sauvage, 1883 (Cypriniformes: Cyprinidae: Leuciscinae) are found in Japan: big-scaled redbfin *T. hakonensis* (Günther, 1877), Sakhalin redbfin (new English name) *T. sachalinensis* (Nikolskii, 1889), Pacific redbfin *T. brandtii* (Dybowski, 1872), and long-jawed redbfin (new English name) *T. nakamuraii* Doi and Shinzawa, 2000 (Hosoya, 2013). *Tribolodon hakonensis* is widely distributed in Japan (Kurawaka, 1977; Sakai, 1989, 1995), where it occurs in various types of freshwater waters, including upper to lower reaches of rivers, brooks, ponds, lakes, and reservoirs. As this species is euhaline, some individuals are found even in brackish and coastal marine waters during their growing period and return to fresh waters for spawning (Sakai, 1995; Ishizaki *et al.*, 2009). The species is usually

abundant in inland waters and is targeted by commercial and recreational fishermen. *Tribolodon brandtii* is distributed north of central Japan. This species also exhibits anadromy: individuals migrate as juveniles from fresh waters to brackish or marine waters and then return as adults to spawn in fresh waters (Sakai, 1989, 1995). Two species, *T. sachalinensis* and *T. nakamuraii*, occur only in fresh waters and are found in northern Japan (Sakai, 1989).

In 1908, a tapeworm of *Ligula* was reported from Lake Onuma, Hokkaido (Anonymous, 1908). Since then, many studies have been conducted for more than one century on the parasites of *Tribolodon* spp. in Japan. In the present checklist, based on the literature published for 106 years between 1908 and 2013, information on the parasites of cyprinid fishes of the genus *Tribolodon* in Japan is compiled in two lists, Parasite-Host List and Host-Parasite List. No abstracts of the papers presented at scientific meetings are herein cited. In total, 44 named species of parasites are listed along with those not identified to species level.

In the **PARASITE-HOST LIST**, parasites are arranged by higher taxon in the following order: Ciliophora, Myxozoa, Trematoda, Monogenea, Cestoda, Nematoda, Acanthocephala, Hirudinida, Mollusca, Branchiura, and Copepoda. Within each higher taxon, genera and species are listed alphabetically. For each species of parasite, the following information is provided:

1) The current **scientific name**, including author(s) and date(s), followed by any original combination, recognized synonym(s), or other identifications(s) that have been used in establishing records from *Tribolodon* spp. in Japan. No attempt has been made to evaluate the taxonomic validity of the published reports, but the ciliate described as *Cyclochaeta leucisci* is herein treated as Trichodinidae gen. sp. because this parasite needs more morphological information for exact identification.

2) The **habitat** in which the parasite was acquired and normally completes its life cycle is given as FW for fresh waters, B for brackish waters, and M for marine waters.

3) The **Site(s) of infection** of the parasite in or on its host. If the site was not given in the original record, the likely site was determined from other records and is enclosed in square brackets.

4) The **Distribution** of the parasite is indicated by prefecture (boundaries shown in Fig. 1), in geographical order from northeast to southwest. For marine or brackish-water species, the name of the prefecture nearest the collection site is given.

5) The **Record(s)**. The authors responsible for the records are listed in chronological order. If a parasite has been reported more than once, the references are numbered, but not when there has been only one record of the parasite. Each reference is followed by the locality or localities given in two parts, first the prefecture(s) and then the detailed collection locality or localities from which the parasite was reported. If no locality record was given, the geographical locality is shown by a dash (—). When all records are from the same prefecture, only the detailed collection locality or localities are listed.

6) Under **Remarks**, explanatory comments are given on systematics, nomenclature, useful references, and notes on specific items such as tentative parasite identifications in the original reports.

7) The **References** section includes works directly cited in the Parasite-Host List. If only a Japanese title was given by the original author(s), our translation of the title into English is provided in square brackets.

In the **HOST-PARASITE LIST**, hosts are listed alphabetically. In each higher taxon, parasites are also listed in alphabetical order, and after the name of each parasite, its geographical distribution is given in parenthesis.




Fig. 1. Map of Japan showing the prefectural boundaries. The following prefectural names are arranged in alphabetical order: Aichi-24; Akita-7; Aomori-5; Chiba-15; Ehime-41; Etorofu Island-1; Fukui-26; Fukuoka-43; Fukushima-10; Gifu-23; Gunma-14; Hiroshima-37; Hokkaido-4; Hyogo-33; Ibaraki-12; Ishikawa-25; Iwate-6; Kagawa-39; Kagoshima-49; Kanagawa-18; Kochi-42; Kumamoto-47; Kunashiri Island-3; Kyoto-29; Mie-28; Miyagi-8; Miyazaki-48; Nagano-20; Nagasaki-45; Nara-30; Niigata-11; Oita-46; Okayama-35; Okinawa-50; Osaka-31; Saga-44; Saitama-16; Shiga-27; Shikotan Island-2; Shimane-36; Shizuoka-21; Tochigi-13; Tokushima-40; Tokyo-17; Tottori-34; Toyama-22; Wakayama-32; Yamagata-9; Yamaguchi-38; and Yamanashi-19.

This checklist is the tenth in the following series of published synopses of the parasites of fishes and shellfishes in Japan: Nagasawa *et al.* (1987) for the parasites of salmonids; Nagasawa *et al.* (1989) for the parasites of freshwater fishes in Hokkaido; Nagasawa (1993a) for the parasites of squids and cuttlefishes; Nagasawa (1993b) for the parasites of gadids; Nagasawa *et al.* (2007a) for the parasites of ayu (*Plecoglossus altivelis altivelis*); Nagasawa *et al.* (2007b) for the parasites of eels (*Anguilla* spp.); Nagasawa *et al.* (2012) for the parasites of medaka (*Oryzias latipes*); Nagasawa (2012) for the parasites of Manila clams (*Ruditapes philippinarum*); and Nagasawa and Nitta (2012) for the parasites of freshwater and brackish-water fishes in Hiroshima Prefecture.

## A PARASITE-HOST LIST

### Ciliophora

- Chilodonella piscicola* (Zacharias, 1894) (FW)  
 Sites of infection: gills, fins  
 Host: *Tribolodon hakonensis*  
 Distribution: Hokkaido  
 Records: 1. Nagasawa *et al.* 1989 (Chitose River); 2. Urawa and Awakura 1994 (—)
- Trichodina fujitai* (Suzuki, 1950) (FW)  
 Previous identification: *Cyclochaeta fujitai* of Suzuki (1950)  
 Site of infection: gills  
 Host: *Tribolodon hakonensis*  
 Distribution: Yamagata, Osaka  
 Record: Suzuki 1950 (Yamagata:—; Osaka:—)  
 Remarks: Recently, this parasite has been transferred from *Cyclochaeta* to *Trichodina* by Nagasawa *et al.* (2012).
- Trichodinidae gen. sp. (FW)  
 Previous identification: *Cyclochaeta leucisci* of Suzuki (1950)  
 Includes: Trichodinidae gen. sp. of Anonymous (2002)  
 Sites of infection: body surface, gills  
 Host: *Tribolodon hakonensis*  
 Distribution: Yamagata, Gifu  
 Records: 1. Suzuki 1950 (Yamagata: Nezugaseki); 2. Anonymous 2002 (Gifu: Kiso River)  
 Remarks: Since more information on the morphology of this parasite is necessary for exact identification, it is herein treated as Trichodinidae gen. sp.
- Ciliophora gen. sp. (FW)  
 Site of infection: fins  
 Host: *Tribolodon hakonensis*  
 Distribution: Nara  
 Record: Nakamura *et al.* 2005 (Takami River)

### Myxozoa

- Chloromyxum richardsonii* Fujita, 1925 (FW)  
 Site of infection: gall bladder  
 Host: *Tribolodon hakonensis*  
 Distribution: Hokkaido  
 Record: Fujita 1925 (Toyohira River)  
 Remarks: There has been no record of this myxozoan since its original description by Fujita (1925).

Myxozoa gen. sp (FW)

Sites of infection: gills, viscera

Host: *Tribolodon hakonensis*

Distribution: Gifu, Nara

Records: 1. Anonymous 2002 (Gifu: Shinsakai River); 2. Nakamura *et al.* 2005 (Nara: Takami River)

## Trematoda

*Urorchis* sp. was listed as a parasite of *T. hakonensis* from Lake Biwa by Shimazu *et al.* (2011: 101) but there has been no record of the trematode from this fish (see Shimazu *et al.*, 2011: 57). Thus, *Urorchis* sp. is not included in this checklist.

*Allocreadium japonicum* Ozaki, 1926 (FW)

Site of infection: [intestine]

Hosts: *Tribolodon hakonensis* ?

*Tribolodon brandtii* ?

Distribution: Tokyo

Record: Shimazu 1988 (Tokyo: Tama River;—: Shinkawa)

Remarks: This trematode was collected from “Haya” (= ? *T. hakonensis*) and “Maruta” (= ? *T. brandtii* [as *T. taczonowskii*]) (Shimazu, 1988).

*Allocreadium tosai* Shimazu, 1988 (FW)

Previous identification: *Allocreadium transversale* of Shimazu (1981)

Site of infection: intestine

Hosts: *Tribolodon hakonensis* (2)

*Tribolodon sachalinensis* (1-2)

Distribution: Hokkaido

Records: 1. Shimazu 1981 (Kushiro River); 2. Shimazu 1988 (Lake Toro [as Lake Tôro])

*Allocreadium tribolodontis* Shimazu and Hashimoto, 1999 (FW)

Previous identification: *Allocreadium isoporum* of Shimazu (1981, 1988)

Site of infection: intestine

Hosts: *Tribolodon hakonensis* (3-4)

*Tribolodon sachalinensis* (1-3)

Distribution: Hokkaido, Iwate

Records: 1. Shimazu 1981 (Hokkaido: Kushiro River); 2. Shimazu 1988 (Hokkaido: Kushiro River); 3. Shimazu and Hashimoto 1999 (Hokkaido: Kushiro River [as River Kushiro]; Iwate: Hei River [as River Hei]); 4. Hashimoto 2000 (Iwate: Hei River)

*Asymphylogora innominata* (Faust, 1924) (FW)

Previous identification: *Asymphylogora macrostoma* of Yamaguti (1934), Shimazu (1992), Nakamura *et al.* (2000), and Shimazu and Urabe (2005)

Site of infection: intestine

Host: *Tribolodon hakonensis*

Distribution: Saitama, Nagano, Toyama, Fukui, Shiga, Nara, Hiroshima

Records: 1. Yamaguti 1934 (Toyama: Namerikawa); 2. Shimazu 1992 (Saitama: Oppe River; Nagano: Nogu River, Torii River, Lake Suwa; Fukui: Obama; Shiga: Lake Biwa; Hiroshima: Eno River, Saijo River); 3. Nakamura *et al.* 2000 (Nara: Takami River); 4. Shimazu and Urabe 2005 (Nara: Takami River); 5. Shimazu *et al.* 2011 (Shiga: Lake Biwa)

Remarks: Although *Asymphyiodora macrostoma* had been used as the scientific name of this trematode in Japan, Shimazu *et al.* (2011) currently used *A. innominata* for the species.

*Centrocestus armatus* (Tanabe, 1922) (metacercaria) (FW)

Sites of infection: musculature, gills, fins

Host: *Tribolodon hakonensis*

Distribution: Shizuoka, Gifu, Oita

Records: 1. Okabe 1940 (Oita: Chikugo River); 2. Ito 1968 (Shizuoka:—); 3. Ito and Mochizuki 1968 (Shizuoka: Tenryu River, Abe River, Kano River); 4. Anonymous 2002 (Gifu: Shinsakai River, Kiso River)

*Clinostomum complanatum* Rudolphi, 1814 (metacercaria) (FW)

Sites of infection: musculature, tissues around gills, pharynx

Host: *Tribolodon hakonensis*

Distribution: Tottori

Record: Aohagi *et al.* 1995 (Koyama Pond)

*Clonorchis sinensis* Looss, 1907 (metacercaria) (FW)

Site of infection: [musculature, subcutaneous tissues, gills, scales, fins]

Hosts: *Tribolodon hakonensis* (1-4)

*Tribolodon brandtii* (4)

Distribution: Miyagi, Nagano, Toyama, Shiga

Record: 1. Ichioka 1930 (Toyama: Ishizutsumi); 2. Komiya *et al.* 1957 (Nagano: Lake Suwa); 3. Komiya *et al.* 1960 (Shiga: Lake Biwa); 4. Yuda 1969 (Miyagi: Abukuma River, Matsushima Bay)

*Echinochasmus milvi* Yamaguti, 1939 (metacercaria) (FW)

Site of infection: [gills]

Host: *Tribolodon hakonensis*

Distribution:—

Record: Koga 1952 (unknown locality in Kyushu)

*Exorchis oviformis* Kobayashi, 1915 (metacercaria) (FW)

Sites of infection: musculature, subcutaneous tissues

Hosts: *Tribolodon hakonensis* (1-5)

*Tribolodon brandtii* (6)

Distribution: Miyagi, Niigata, Tokyo

Records: 1. Kobayashi 1915 (—); 2. Kobayashi 1921 (—); 3. Nihei 1961 (Tokyo: Edo River); 4.

Nihei 1962 (Tokyo: Edo River); 5. Saito *et al.* 1964 (Niigata: near Lake Toyanogata); 6. Yuda 1969 (Miyagi: Matsushima Bay)

*Holostephanus nipponicus* Yamaguti, 1939 (metacercaria) (FW)

Site of infection: [musculature]

Host: *Tribolodon hakonensis*

Distribution: Yamagata

Record: Saito and Otsuru 1965a (Miyamatsu River)

*Isoparorchis hypselobagri* (Billet, 1898) (FW)

Previous identification: *Leptolecithum eurytremum* of Kobayashi (1915, 1921)

Site of infection: body cavity

Host: *Tribolodon hakonensis*

Distribution: Ibaraki

Records: 1. Kobayashi 1915 (—); 2. Kobayashi 1921 (—); 3. Yamaguti 1934 (Ibaraki: Lake Kasumigaura [as Kasumiga-ura])

Remarks: Only immature worms of this trematode occurs in *T. hakonensis* (Kobayashi, 1915, 1921; Yamaguti, 1934). Although Kobayashi (1915, 1921) gave no detailed localities of the species, its adults were collected in various localities in Okayama Prefecture, Sawara in Chiba Prefecture, Lake Kasumigaura (as Kasumiga-ura) in Chiba Prefecture, and Lake Biwa in Shiga Prefecture. Recently, Nagasawa *et al.* (2013) reviewed the biology of the species infecting Japanese freshwater fishes based on the literature published between 1915 and 2013.

*Metagonimus katsuradai* Izumi, 1935 (metacercaria) (FW)

Sites of infection: scales, fins, gills

Host: *Tribolodon hakonensis*

Distribution: Oita

Record: Okabe 1940 (Chikugo River)

*Metagonimus miyatai* Saito, Chai, Kim, Lee and Rim, 1997 (metacercaria) (FW)

Sites of infection: epidermis, scales

Hosts: *Tribolodon hakonensis* (1-2)

*Tribolodon sachalinensis* (2)

Distribution: Hokkaido, Hiroshima

Records: 1. Saito 1984 (Hiroshima: Ota River); 2. Shimazu 2002 (Hokkaido: Ishikari River and its tributaries)

Remarks: See the remarks on *M. yokogawai*.

*Metagonimus takahashii* Suzuki in Takahashi, 1929 (metacercaria) (FW)

Previous identification: *Metagonimus yokogawai* var. *takahashii* of Ochi (1957), Saito and Otsuru (1965b), and Takahashi (1967)

Includes: *Metagonimus yokogawai ovatus* of Koga (1938, 1939) and Kokame (1939); *Metagonimus* sp. Koga type of Saito (1984) and Shimazu (2002)



Sites of infection: epidermis, scales

Hosts: *Tribolodon hakonensis* (1-8)

*Tribolodon sachalinensis* (8)

Distribution: Hokkaido, Niigata, Toyama, Ishikawa, Okayama, Hiroshima, Yamaguchi, Oita

Records: 1. Koga 1938 (Oita: Chikugo River); 2. Koga 1939 (Oita: Chikugo River); 3. Kokame 1939 (Ishikawa: Daishoji River); 4. Ochi 1957 (Okayama: Asahi River; Hiroshima: Ashida River, Nuta River, Ota River, Gono River; Yamaguchi: Asa River); 5. Saito and Otsuru 1965b (Niigata: Lake Yoroigata); 6. Takahashi 1967 (Oita: Chikugo River); 7. Saito 1984 (Toyama:—); 8. Shimazu 2002 (Hokkaido: Ishikari River and its tributaries)

Remarks: Saito (1984) suggested that *Metagonimus* worms reported from the Sho River, Toyama Prefecture (Saito, 1968a) and Chōkai Village, Akita Prefecture (Yoshimura *et al.*, 1972) may be identifiable as *Metagonimus* sp. Koga type.

*Metagonimus yokogawai* (Katsurada, 1912) (metacercaria) (FW)

Previous identification: *Loxotrema ovatum* of Kobayashi (1912)

Sites of infection: scales, fins, epidermis

Hosts: *Tribolodon hakonensis* (1-16, 17-33)

*Tribolodon sachalinensis* (24, 26, 30)

*Tribolodon brandtii* (26)

*Tribolodon* sp. (17, 24)

Distribution: Hokkaido, Aomori, Akita, Miyagi, Yamagata, Niigata, Gunma, Kanagawa, Shizuoka, Toyama, Ishikawa, Mie, Shimane, Hiroshima, Yamaguchi, Tokushima, Oita, Miyazaki, Kumamoto

Records: 1. Kobayashi 1912 (Miyagi: Shimekiri Swamp); 2. Koga 1922 (Kumamoto: Suizenji, Lake Ezu); 3. Ochi 1928 (Miyagi: Kitagami River; Mie: Choshi River, Kumano River; Tokushima: Yoshino River; Miyazaki: Ichinose River); 4. Ichioka 1930 (Toyama: Ishizutsumi); 5. Taki 1935 (Oita: Banjo River, Kusu River); 6. Gushima 1939a (—: Chikugo River); 7. Gushima 1939b (—: Chikugo River); 8. Gushima 1939c (—: Chikugo River); 9. Okabe 1940 (Oita: Chikugo River); 10. Takabayshi 1953 (Yamaguchi: Koto River); 11. Yokogawa *et al.* 1962 (Shizuoka: Kiku River); 12. Nihei *et al.* 1964 (Gunma: Tone River [as River Tone]); 13. Kagei 1966 (Shimane: Takatsu River); 14. Takahashi 1967 (Shimane: Kando River; Oita: Chikugo River); 15. Ito 1968 (Shizuoka: Abe River); 16. Okabe *et al.* 1968 (Oita: Chikugo River); 17. Saito 1968a (Yamagata: No River; Niigata: Sukebuchi River, Agano River, Shinano River, Uono River, Lake Fukushima, Lake Yoroigata; Toyama: Sho River, Koyabe River); 18. Saito 1968b (Niigata: Lake Fukushima; Toyama: Sho River); 19. Yoshimura *et al.* 1972 (Akita: rivers in Chōkai Village); 18. Kobayashi 1972 (Shizuoka: Kano River); 21. Kobayashi *et al.* 1972 (Shizuoka: Kano River); 22. Nakade 1972 (Aomori: Iwaki River; Akita: Koyoshi River); 23. Tani *et al.* 1974 (Akita: Yoneshiro River, Lake Hachirogata, Omono River, Koyoshi River); 24. Miyamoto and Kutsumi 1978 (Hokkaido: Biei River, Ishikari River); 25. Yoshimura *et al.* 1978 (Ishikawa: Machino River, Hoshi River, Kawarada River, San-no River); 26. Miyamoto and Kutsumi 1980 (Hokkaido: Ishikari River and tributaries); 27. Ohnishi 1983 (Ishikawa: Kawarada River, Machino River); 28. Saito 1984 (Hiroshima: Ota River); 29. Saito *et al.* 1984 (Yamagata: Nikko River, Nyu River, Oguni River, Mogami River, Daimon River, Matsu River, Nezugaseki River,

Atsumi River; Shimane:—; Hiroshima:—); 30. Miyamoto 1985 (Hokkaido: Asahikawa); 31. Ohnishi 1987 (—); 32. Saito *et al.* 1997 (Hiroshima: Ota River [as Ohta River]); 33. Uchida *et al.* 1999 (Kanagawa: Haya River, Sakawa River; Shizuoka: Kano River, Raiko River [as Raikou River])

Remarks: According to Shimazu (2002), the worms reported as *Metagonimus yokogawai* by Miyamoto and Kutsumi (1980) from Hokkaido contained *M. miyatai*, *M. yokogawai* and the Koga type. Although Uchida *et al.* (1999) reported *M. yokogawai* from *T. hakonensis* from Kanagawa and Shizuoka prefectures, they recognized four types (Type I [= *M. yokogawai*], Type II [= *M. miyatai*], type III [= *M. takahashii*] and Type IV [= Koga type]) in adult specimens from cats experimentally infected with *Metagonimus* metacercariae from the fish.

*Metagonimus* spp. (metacercariae) (FW)

Sites of infection: scales, under scales

Hosts: *Tribolodon hakonensis* (1-2, 4-6)

*Tribolodon brandtii* (3)

Distribution: Hokkaido, Miyagi, Yamagata, Shizuoka, Gifu, Hiroshima

Records: 1. Ito 1968 (Shizuoka:—); 2. Ito and Mochizuki 1968 (Shizuoka: Abe River); 3. Yuda 1969 (Miyagi: Matsushima Bay); 4. Miyamoto and Kutsumi 1980 (Hokkaido: Mu River [as Mukawa River]); 5. Saito 1984 (Yamagata:—; Hiroshima:—); 6. Loganathan *et al.* 1989 (Gifu: Nagara River [as River Nagaragawa])

*Neoplagioporus elongatus* (Goto and Ozaki, 1930) (FW)

Site of infection: intestine

Host: *Tribolodon hakonensis*

Distribution: Shiga

Records: 1. Shimazu 1990 (Lake Biwa); 2. Shimazu *et al.* 2011 (Lake Biwa)

*Pseudexorchis major* (Hasegawa, 1935) (metacercaria) (FW)

Sites of infection: scales, fins, gills

Hosts: *Tribolodon hakonensis* (1-6)

*Tribolodon brandtii* (2-3)

Distribution: Tokyo, Shizuoka, Gifu, Oita

Records: 1. Okabe 1940 (Oita: Chikugo River); 2. Nihei 1961 (Tokyo: Edo River); 3. Nihei 1962 (Tokyo: Edo River); 4. Ito 1968 (Shizuoka:—); 5. Ito and Mochizuki 1968 (Shizuoka: Abe River); 6. Anonymous 2002 (Gifu: Kiso River)

*Pseudozoogonoides ugui* Shimazu, 1974 (M)

Includes: *Pseudozoogonoides* sp. of Machida *et al.* (1972)

Site of infection: intestine

Host: *Tribolodon hakonensis*

Distribution: Hokkaido, Iwate

Records: 1. Machida *et al.* 1972 (Hokkaido: North Pacific Ocean off the Hidaka District); 2. Shimazu 1974 (Hokkaido: Inshore water of Wakkanai, Nukui River [as River Nukui] near

Gabino; Iwate: North Pacific Ocean off Otsuchi [as Ootsuchi]); 3. Hashimoto 2000 (Iwate: Hei River)

*Digenea* gen. spp. (metacercaria) (FW)

Sites of infection: epidermis, muscle

Hosts: *Tribolodon hakonensis* (1-4)

*Tribolodon sachalinensis* (3)

Distribution: Hokkaido, Akita, Toyama, Nara

Records: 1. Imagawa 1934 (Akita: Lake Hachirogata); 2. Saito and Otsuru 1965a (Toyama: Oyabe River); 3. Miyamoto and Kutsumi 1980 (Hokkaido: Tokachi River, Kushiro River); 3. Nakamura *et al.* 2000 (Nara: Takami River)

### Monogenea

*Dactylogyrus* sp. (FW)

Site of infection: gills

Hosts: *Tribolodon hakonensis* (1-2)

*Tribolodon sachalinensis* ? (2)

*Tribolodon brandtii* (2)

Distribution: Hokkaido

Records: 1. Nagasawa *et al.* 1989 (Lake Toro, Kushiro River, Ishikari-Furukawa); 2. Ogawa 1994 (Lake Toro)

*Diplozoon* sp. (FW)

Site of infection: gills

Hosts: *Tribolodon hakonensis* (1-6, 8-9)

*Tribolodon sachalinensis* (5-6)

*Tribolodon brandtii* (6)

*Tribolodon nakamuraii* (7)

Distribution: Hokkaido, Saitama, Niigata, Gifu, Nara

Records: 1. Okura *et al.* 1985a (Saitama: Ara River, Iruma River, Toki River, Oppe River); 2. Okura *et al.* 1985b (Saitama: Ara River, Saitama Prefectural Fisheries Experimental Station); 3. Suzuki and Okura 1987 (Saitama: Ara river, Iruma River, Toki River, Oppe River, Tone River, Kanna River); 4. Suzuki and Okura 1988 (Saitama:—); 5. Nagasawa *et al.* 1989 (Hokkaido: Horobetsu River, Teshio River, Mena River, Ebetsu, Lake Toro, Lake Barato); 6. Ogawa 1994 (Hokkaido: Lake Toro, Mena River, Chitose River); 7. Shindo 1997 (Niigata: aquarium); 8. Nakamura *et al.* 2000 (Nara: Takami River); 9. Anonymous 2002 (Gifu: Kiso River)

*Gyrodactylus* sp. (FW)

Includes: *Gyrodactylus* sp. 1 of Nagasawa *et al.* (1989)

Sites of infection: fins, gills

Host: *Tribolodon hakonensis*

Distribution: Hokkaido, Nara

Records: 1. Nagasawa *et al.* 1989 (Hokkaido: Lake Toro, Ebetsu); 2. Ogawa 1994 (Hokkaido: Lake Toro, Chitose River); 3. Nakamura *et al.* 2000 (Nara: Takami River)

Monopisthocotylea gen. sp. (FW)

Site of infection:—

Host: *Tribolodon hakonensis*

Distribution: Gifu

Record: Anonymous 2002 (Shinsakai River)

### Cestoda

*Bothriocephalus acheilognathi* Yamaguti, 1934 (FW)

Previous identification: *Coelobothrium oitense* of Kugi and Matsuo (1990)

Site of infection: intestine

Host: *Tribolodon hakonensis*

Distribution: Oita

Record: Kugi and Matsuo 1990 (Chikugo River)

Remarks: *Coelobothrium oitense* was regarded as a junior synonym of *B. acheilognathi* by Kuchta and Scholz (2007).

*Caryophyllaeides ergensi* Scholz, 1990 (FW)

Site of infection: intestine

Hosts: *Tribolodon hakonensis* (1-3)

*Tribolodon sachalinensis* (1)

Distribution: Hokkaido, Aomori, Shizuoka, Nagano

Records: 1. Scholz *et al.* 2001 (Hokkaido: Onishibetsu River, Lake Abashiri, Lake Toro [as Lake Tôro], Shokanbetsu River, Chitose River, Oono River, Mogusa River, Barato River; Aomori: Lake Ogawara; Nagano: Lake Suwa, Hiroi River, Shizuoka: Okitsu River); 2. Olson *et al.* 2001 (Nagano: Hiroi River); 3. Oros *et al.* 2010 (Aomori: Lake Ogawara [as Ogawara Lake])

Caryophyllidea fam. gen. sp. (FW)

Includes: "A monozootic cestode" of Shimazu (1981)

Site of infection: intestine

Host: *Tribolodon sachalinensis*

Distribution: Hokkaido

Record: Shimazu 1981 (Kushiro River)

Caryophyllidae gen. sp. (FW)

Site of infection: intestine

Host: *Tribolodon hakonensis*

Distribution: Iwate

Record: Hashimoto 2000 (Hienuki River, Hei River)

*Ligula interrupta* Rudolphi, 1810 (plerocercoid)

(FW)

Previous identification: *Digramma alternans* of Awakura *et al.* (1976), Awakura and Kawamura (1977), Kawamura (1982), and Sato *et al.* (1991)

Includes: *Ligula* sp. of Anonymous (1908), Hirasaka (1914), Ishii (1914, 1931), and Handa and Araki (1930)

Site of infection: body cavity

Hosts: *Tribolodon hakonensis* (1-12)

*Tribolodon sachalinensis* (11)

Distribution: Hokkaido, Tochigi, Gunma, Kanagawa, Yamanashi

Records: 1. Anonymous 1908 (Hokkaido: Lake Onuma); 2. Hirasaka 1914 (Yamanashi: Lake Yamanaka); 3. Ishii 1914 (Tochigi: Lake Chuzenji); 4. Handa and Araki 1930 (Hokkaido: Lake Onuma, Lake Konuma, Lake Junsai); 5. Ishii 1931 (Tochigi: Lake Chuzenji; Gunma: Lake Oze; Kanagawa: Lake Ashi); 6. Miyaji 1963 (Tochigi: Lake Chuzenji); 7. Awakura *et al.* 1976 (Hokkaido: Lake Toro [as Lake Tôro]); 8. Awakura and Kawamura 1977 (Hokkaido: Hidaka-horobetsu River); 9. Kawamura 1982 (Hokkaido: Samani River, Utabetsu River); 10. Satoh *et al.* 1991 (Kanagawa: Lake Ashi); 11. Awakura 1994 (Hokkaido: Lake Akan, Lake Shikaribetsu, Hidaka-horobetsu River, Samani River); 12. This paper (Gunma: a brook flowing into Lake Okutone) (Fig. 2)

Remarks: Recently, *Digramma* has been regarded as a junior synonym of *Ligula* (Kuchta *et al.*, 2008). Awakura (1994) summarized the information on this cestode (as *D. interrupta*) from cyprinid fishes in Hokkaido.



Fig. 2. A plerocercoid of *Ligula interrupta* Rudolphi, 1810 from a Japanese dace (*Tribolodon hakonensis*), 189 mm in standard length, collected in a brook flowing into Lake Okutone, Gunma Prefecture, central Japan, on 7 September 2000. Photograph by Y. Saito.

## Nematoda

- Anisakis simplex* (Rudolphi, 1809) (larva) (M)  
 Includes: *Anisakis* sp. type I of Miyamoto and Kutsumi (1980)  
 Sites of infection: musculature, mesentery  
 Host: *Tribolodon hakonensis*  
 Distribution: Hokkaido, Iwate  
 Records: 1. Miyamoto and Kutsumi 1980 (Hokkaido: Kushiro River); 2. Moravec *et al.* 1985 (Hokkaido: Lake Toro); 3. Hashimoto 2000 (Iwate: Hei River)
- Camallanus cotti* Fujita, 1927 (FW)  
 Site of infection: intestine  
 Host: *Tribolodon hakonensis*  
 Distribution:—  
 Record: Yamaguti 1935 (—)
- Camallanidae gen. sp. (FW)  
 Previous identification: “camallanid nematodes” of Shimazu (1981)  
 Site of infection: digestive tract  
 Host: *Tribolodon sachalinensis*  
 Distribution: Hokkaido  
 Record: Shimazu 1981 (Kushiro River)
- Gnathostoma nipponicum* Yamaguti, 1941 (larva) (FW)  
 Sites of infection: [abdominal wall, head, viscera]  
 Host: *Tribolodon hakonensis*  
 Distribution: Aomori  
 Records: 1. Oyamada *et al.* 1996a (unspecified rivers); 2. Oyamada *et al.* 1996b (unspecified localities); 3. Oyamada *et al.* 1997 (experimental infection)  
 Remarks: Although Oyamada *et al.* (1996b) gave no detailed localities of the species, the fish examined were collected in Shichinohe, Kamikita, Tenmabayashi, Tohoku, and Rokkasho.
- Hysterothylacium aduncum* (Rudolphi, 1802) (larva) (FW)  
 Site of infection: intestine  
 Host: *Tribolodon sachalinensis*  
 Distribution: Hokkaido  
 Record: Moravec *et al.* 1985 (Lake Toro)
- Pseudocapillaria tomentosa* (Dujardin, 1843) (FW)  
 Previous identification: *Capillaria ugui* of Yamaguti (1941)  
 Site of infection: intestine  
 Host: *Tribolodon hakonensis*  
 Distribution: Tokyo, Shizuoka, Fukui, Nara

Records: 1. Yamaguti 1941 (Fukui [as Hukui]: Obama); 2. Moravec and Nagasawa 1989 (Tokyo: Tama River); 3. Moravec *et al.* 1998 (Shizuoka: Okitsu River); 4. Nakamura *et al.* 2000 (Nara: Takami River)

*Raphidascaris gigi* Fujita, 1928 (larva) (FW)

Previous identifications: *Raphidascaris biwakoensis* of Fujita (1928); *Raphidascaris plecoglossi* of Fujita (1928)

Site of infection: abdominal cavity

Host: *Tribolodon hakonensis*

Distribution: Shiga

Record: Fujita 1928 (Lake Biwa)

*Rhabdochona coronacauda* Belouss, 1965 (FW)

Site of infection: intestine

Host: *Tribolodon hakonensis*

Distribution: Nara

Records: 1. Nakamura *et al.* 2000 (Takami River); 2. Hirasawa *et al.* 2004 (Takami River)

*Rhabdochona denunata honshuensis* Moravec and Nagasawa, 1989 (FW)

Includes: *Rhabdochona denunata* of Mori *et al.* (1998) and Nakamura *et al.* (2000)

Site of infection: intestine

Host: *Tribolodon hakonensis*

Distribution: Nara

Records: 1. Mori *et al.* 1998 (Takami River); 2. Nakamura *et al.* 2000 (Takami River); 3. Hirasawa *et al.* 2004 (Takami River)

*Rhabdochona zacconis* Yamaguti, 1935 (FW)

Site of infection: intestine

Host: *Tribolodon hakonensis*

Distribution: Hokkaido, Aomori, Iwate, Tokyo, Nagano, Shizuoka, Shiga, Nara

Records: 1. Moravec *et al.* 1981 (Hokkaido: Lake Akan, Lake Shikotsu [as Lake Chitose], Chihase River; Shiga: Lake Biwa); 2. Moravec and Nagasawa 1989 (Hokkaido: Amano River; Aomori: Kanita River; Tokyo: Tama River; Nagano: Chikuma River); 3. Moravec *et al.* 1998 (Shizuoka: Okitsu River); 4. Nakamura *et al.* 2000 (Nara: Takami River); 5. Hashimoto 2000 (Iwate: Hienuki River, Hei River)

Remarks: Moravec *et al.* (1981) reported this nematode from *T. hakonensis* and *Zacco platypus* collected in the above four localities, but it is not clear whether the specimens from Lake Biwa were collected from *T. hakonensis* and/or *Z. platypus*.

*Rhabdochona* sp. (FW)

Site of infection: intestine

Host: *Tribolodon brandtii*

Distribution: Niigata

Record: Hokari *et al.* 1973 (Agano River)

Rhabdochonidae gen. sp. (FW)

Previous identification: "rhabdochonid nematodes" of Shimazu (1981)

Site of infection: digestive tract

Host: *Tribolodon sachalinensis*

Distribution: Hokkaido

Record: Shimazu 1981 (Kushiro River)

Nematoda gen. sp. (FW)

Site of infection: intestine

Hosts: *Tribolodon hakonensis* (1)

*Tribolodon sachalinensis* (1)

*Tribolodon* sp. (2)

Distribution: Hokkaido

Records: 1. Inukai 1949 (Bibi River); 2. Kanoh 1949 (—)

### Acanthocephala

*Acanthocephalus opsariichthydis* Yamaguti, 1935 (FW)

Site of infection: [intestine]

Host: *Tribolodon hakonensis*

Distribution: Nagano

Record: Yamaguti 1939 (Lake Suwa)

*Acanthocephalus* sp. (FW)

Site of infection: intestine

Host: *Tribolodon hakonensis*

Distribution: Iwate

Record: Hashimoto 2000 (Hienuki River, Hei River)

*Pseudorhadinorhynchus leuciscus* (Krotov and Petrochenko, 1956) (M)

Site of infection: intestine

Host: *Tribolodon hakonensis*

Distribution: Hokkaido, Iwate

Records: 1. Machida and Araki 1982 (Hokkaido: North Pacific Ocean off the Hidaka District, Wakkanai, Nukuigawa, Otoshibe; Iwate: North Pacific Ocean off the Sanriku District); 2. Araki and Machida 1988 (Hokkaido: Samani); 3. Nagasawa *et al.* 1989 (Hokkaido: North Pacific Ocean off Kushiro, Assabu River); 4. Hashimoto 2000 (Iwate: Hei River)

Acanthocephala gen. sp. (FW)

Site of infection:—

Host: *Tribolodon hakonensis*



Distribution: Gifu

Record: Anonymous 2002 (Shinsakai River)

### Hirudinida

*Limnotrachelobdella okae* (Moore, 1924)

(B or M)

Site of infection: body surface

Host: *Tribolodon brandtii*

Distribution: Niigata

Record: Nagasawa *et al.* 2008 (Iwafune Fishing Port)

### Mollusca

*Pronodularia japonensis* (Lea, 1959) (glochidium)

(FW)

Previous identification: *Inversidens japonensis* of Miyabe *et al.* (2007)

Site of infection: [fins]

Host: *Tribolodon hakonensis*

Distribution: Chiba

Record: Miyabe *et al.* 2007 (experimental infection)

### Branchiura

*Argulus* sp.

(FW)

Sites of infection: body surface, fins

Host: *Tribolodon hakonensis*

Distribution: Nara

Record: Nakamura *et al.* 2000 (Takami River)

### Copepoda

*Caligus orientalis* Gusev, 1951

(B)

Site of infection: body surface

Host: *Tribolodon hakonensis*

Distribution: Hokkaido

Record: Urawa and Kato 1991 (Lake Mokoto)

*Caligus punctatus* Shiino, 1955

(M)

Site of infection: body surface

Host: *Tribolodon hakonensis*

Distribution: Aomori, Miyagi

Records: 1. Shiino 1955 (Miyagi: Matsushima Bay); 2. Shiino 1959 (Aomori: Asamushi)

- Ergasilus hypomesi* Yamaguti, 1936 (FW)  
 Sites of infection: body surface, fins, gills  
 Host: *Tribolodon hakonensis*  
 Distribution: Hokkaido  
 Record: Nagasawa *et al.* 1989 (Lake Barato)
- Lepeophtheirus salmonis* (Krøyer, 1837) (M)  
 Site of infection: body surface  
 Host: *Tribolodon hakonensis*  
 Distribution: Hokkaido  
 Record: Nagasawa *et al.* 1994 (Furuu River)
- Lernaea cyprinacea* Linnaeus, 1758 (FW)  
 Site of infection: head embedded in musculature with body protruding externally  
 Host: *Tribolodon hakonensis*  
 Distribution: Hokkaido  
 Records: 1. Tsutsumi 1978 (—: aquarium); 2. Nagasawa *et al.* 1989 (Hokkaido: Kikonai River)
- Neoergasilus japonicus* (Harada, 1930) (FW)  
 Site of infection: fins  
 Host: *Tribolodon hakonensis*  
 Distribution: Hokkaido  
 Record: Nagasawa *et al.* 1989 (Lake Barato, Ishikari River)

### Isopoda

- Nerocila japonica* Shioedte and Meinert, 1881 (M or B)  
 Site of infection: body surface  
 Host: *Tribolodon hakonensis*  
 Distribution: Shimane  
 Record: Yamauchi and Nagasawa 2012 (Lake Nakaumi)

### A HOST-PARASITE LIST

***Tribolodon brandtii* (Dybowski, 1872) Pacific redfin, “maruta” (Japanese name)**

Trematoda

- Allocreadium japonicum* (Tokyo)  
*Clonorchis sinensis* (Miyagi)  
*Exorchis oviformis* (Miyagi)  
*Metagonimus yokogawai* (Hokkaido)  
*Metagonimus* sp. (Miyagi)  
*Pseudexorchis major* (Tokyo)

Monogenea

*Dactylogyrus* sp. (Hokkaido)

*Diplozoon* sp. (Hokkaido)

Nematoda

*Rhabdochona* sp. (Niigata)

Hirudinida

*Limnotrachelobdella okae* (Niigata)

***Tribolodon hakonensis* (Günther, 1877) Big-scaled redfin, “ugui” (Japanese name)**

Ciliophora

*Chilodonella piscicola* (Hokkaido)

*Trichodina fujitai* (Yamagata, Osaka)

Trichodinidae gen. sp. (Yamagata, Gifu)

Ciliophora gen. sp. (Nara)

Myxozoa

*Chloromyxum richardsonii* (Hokkaido)

Myxozoa gen. sp. (Gifu, Nara)

Trematoda

*Allocreadium japonicum* (Tokyo)

*Allocreadium tosai* (Hokkaido)

*Allocreadium Tribolodontis* (Iwate)

*Asymphyiodora innominata* (Saitama, Nagano, Toyama, Fukui, Shiga, Nara, Hiroshima)

*Centrocestus armatus* (Shizuoka, Gifu, Oita)

*Clinostomum complanatum* (Tottori)

*Clonorchis sinensis* (Miyagi, Toyama, Nagano, Shiga)

*Echinochasmus milvi* (—)

*Exorchis oviformis* (Niigata, Tokyo)

*Holostephanus nipponicus* (Yamagata)

*Isoparorchis hypselobagri* (Ibaraki)

*Metagonimus katuradai* (Oita)

*Metagonimus miyatai* (Hokkaido, Hiroshima)

*Metagonimus takahashii* (Niigata, Toyama, Ishikawa, Okayama, Hiroshima, Yamaguchi, Oita)

*Metagonimus yokogawai* (Hokkaido, Aomori, Akita, Miyagi, Yamagata, Niigata, Gunma, Kanagawa, Shizuoka, Toyama, Ishikawa, Mie, Shimane, Hiroshima, Yamaguchi, Tokushima, Oita, Miyazaki, Kumamoto)

*Metagonimus* spp. (Hokkaido, Yamagata, Shizuoka, Gifu, Hiroshima)

*Neoplagioporus elongatus* (Shiga)

*Pseudexorchis major* (Tokyo, Shizuoka, Gifu, Oita)

*Pseudozoogonoides ugui* (Hokkaido, Iwate)

Digenea gen. spp. (Hokkaido, Akita, Toyama, Nara)

Monogenea

*Dactylogyrus* sp. (Hokkaido)

*Diplozoon* sp. (Hokkaido, Saitama, Gifu, Nara)

*Gyrodactylus* sp. (Hokkaido)

Monopisthocotylea gen. sp. (Gifu)

Cestoda

*Bothriocephalus acheilognathi* (Oita)

*Caryophyllaeides ergensi* (Hokkaido, Aomori, Shizuoka, Nagano)

Caryophyllidea fam. gen. sp. (Hokkaido)

Caryophyllidae gen. sp. (Iwate)

*Ligula interrupta* (Hokkaido, Tochigi, Gunma, Kanagawa, Yamanashi)

Nematoda

*Anisakis simplex* (Hokkaido, Iwate)

*Camallanus cotti* (—)

*Gnathostoma nipponicum* (Aomori)

*Hysterothylacium aduncum* (Hokkaido)

*Pseudocapillaria tomentosa* (Tokyo, Shizuoka, Fukui, Nara)

*Raphidascaris gigi* (Shiga)

*Rhabdochona coronacauda* (Nara)

*Rhabdochona denunata honshuensis* (Nara)

*Rhabdochona zacconis* (Hokkaido, Aomori, Iwate, Tokyo, Nagano, Shizuoka, Shiga, Nara)

Nematoda gen. sp. (Hokkaido)

Acanthocephala

*Acanthocephalus opsariichthydis* (Nagano)

*Acanthocephalus* sp. (Iwate)

*Pseudorhadinorhynchus leuciscus* (Hokkaido, Iwate)

Acanthocephala gen. sp. (Gifu)

Mollusca

*Pronodularia japonensis* (Chiba)

Branchiura

*Argulus* sp. (Nara)

Copepoda

*Caligus orientalis* (Hokkaido)

*Caligus punctatus* (Aomori, Miyagi)

*Ergasilus hypomesi* (Hokkaido)

*Lepeophtheirus salmonis* (Hokkaido)

*Lernaea cyprinacea* (Hokkaido)

*Neoergasilus japonicus* (Hokkaido)

Isopoda

*Nerocila japonica* (Shimane)

***Tribolodon nakamuraii* Doi and Shinzawa, 2000      Long-jawed redbin (new English name),  
“ukekuchi-ugui” (Japanese name)**

Monogenea

*Diplozoon* sp. (Niigata)

***Tribolodon sachalinensis* (Nikolskii, 1889) Sakhalin redfin (new English name),  
“ezo-ugui” (Japanese name)**

Trematoda

- Allocreadium tosai* (Hokkaido)  
*Allocreadium Tribolodontis* (Hokkaido)  
*Metagonimus miyatai* (Hokkaido)  
*Metagonimus takahashii* (Hokkaido)  
*Metagonimus yokogawai* (Hokkaido)  
 Digenea gen. spp. (Hokkaido)

Monogenea

- Dactylogyrus* sp. (Hokkaido)  
*Diplozoon* sp. (Hokkaido)

Cestoda

- Caryophyllaeides ergensi* (Hokkaido)  
*Ligula interrupta* (Hokkaido)

Nematoda

- Camallanidae gen. sp. (Hokkaido)  
 Rhabdochonidae gen. sp. (Hokkaido)  
 Nematoda gen. sp. (Hokkaido)

***Tribolodon* sp.**

Trematoda

- Metagonimus yokogawai* (Hokkaido, Yamagata, Niigata, Toyama)

Nematoda

- Nematoda gen. sp. (Hokkaido)

## ACKNOWLEDGMENTS

We thank Mr. Yuya Saito for providing us with a picture of *Ligula interrupta* and *Tribolodon hakonensis*.

## REFERENCES

- Anonymous, 1908. [On the parasite of crucian carp in Lake Onuma]. *Hokkaido Suisan Zasshi (Hokkaido Fisheries Magazine)*, **8**(5): 5-6. [In Japanese].
- Anonymous, 2002. [The parasite fauna of fishes in an experimental river and its characteristics]. *Annual Report of the Aqua Restoration Research Center 2001*: 206-218. [In Japanese].
- Aohagi, Y., Shibahara, T., Kagota, K., 1995. Metacercariae of *Clinostomum complanatum* found from new fish hosts, *Lateolabrax japonicus* and *Leuciscus hakonensis*. *Japanese Journal of Parasitology*, **44**: 340-342.
- Araki, J., Machida, M., 1988. Some acanthocephalans from marine fishes of northern Japan, with descriptions of two new species, *Acanthocephaloides ichiharai* and *A. claviformis*. *Bulletin of the National Science Museum, Tokyo, Series A*, **13**: 1-11.
- Awakura, T., 1994. Cestodes of freshwater fishes of Hokkaido. *Scientific Reports of the Hokkaido Fish*

- Hatchery*, (48): 79-82.
- Awakura, T., Kawamura, H., 1977. Note on the *Digramma alternans* (Rud., 1810) found on [sic] the Japanese dace, *Tribolodon hakonensis* inhabited [sic] a river in Hokkaido, Japan. *Fish Pathology*, **12**: 205-207. [In Japanese with English abstract].
- Awakura, T., Tonosaki, H., Ito, T., 1976. Ligulosis of cyprinid in the lakes of Hokkaido, Japan. *Scientific Reports of the Hokkaido Fish Hatchery*, (31): 67-81.
- Fujita, T., 1925. On two species of *Chloromyxum* found in the fresh-water fishes in Sapporo. *Annotationes Zoologicae Japonenses*, **10**: 277-280.
- Fujita, T., 1928. Further studies on nematodes from fishes of Lake Biwa. *Dobutsugaku Zasshi (Zoological Magazine)*, **40**: 303-314. [In Japanese with English abstract].
- Gushima, M., 1939a. [Immunological studies on *Metagonimus yokogawai*]. *Igaku Kenkyu*, **13**: 593-636. [In Japanese].
- Gushima, M., 1939b. [On the development of *Metagonimus yokogawai* in mice and chicks]. *Igaku Kenkyu*, **13**: 637-655. [In Japanese].
- Gushima, M., 1939c. [Immunological studies on *Metagonimus yokogawai* (continued)]. *Igaku Kenkyu*, **13**: 1713-1796. [In Japanese].
- Handa, Y., Araki, K., 1930. [Surveys in Lakes and Ponds. Part 1. Surveys in Lake Onuma, Lake Konuma and Lake Junsai]. *Suisan Chosa Hokoku (Fisheries Research Reports)*, **21**: 1-66. [In Japanese].
- Hashimoto, K., 2000. Comparison between fluvial and anadromous types of Japanese dace based on *Pseudozoogonoides ugui* Shimazu, 1974 (Digenea, Zoogonidae). *Japanese Society for Systematic Parasitology Circular*, (18): 1-4. [In Japanese].
- Hirasaka, K., 1914. ["*Ligula*" in Lake Yamanaka]. *Dobutsugaku Zasshi (Zoological Magazine)*, **26**: 89. [In Japanese].
- Hirasawa, R., Urabe, M., Yuma, M., 2004. Relationship between intermediate host taxon and infection by nematodes of the genus *Rhabdochona*. *Parasitology International*, **53**: 89-97.
- Hokari, N., Honma, Y., Ito, S., 1973. Winter food of some freshwater fishes in the lower reaches of the River Agano, with reference to parasitic nematodes. *Journal of Water and Waste*, **15**: 1197-1201. [In Japanese].
- Hosoya, K., 2013. *Tribolodon brandtii* (Dybowski, 1872), *Tribolodon nakamuraii* Doi and Shinzawa, 2000, *Tribolodon sachalinensis* (Nikolsky, 1889), *Tribolodon hakonensis* (Günther, 1877). In: *Fishes of Japan with Pictorial Keys to the Species, Third Edition*, ed., Nakabo, T., Tokai University Press, Hadano: 321. [In Japanese].
- Ichioka, S., 1930. [On the distribution of human parasites using freshwater fishes as intermediate hosts and *Ligula*-like larval diphyllbothriids in frogs in the area of Ishizutsumi Village, Nishitonami County, Toyama Prefecture, recognized as a new endemic area of rickets]. *Byorigaku Kyo (Bulletin of Pathology)*, **6**(4): 2-14. [In Japanese].
- Imagawa, O., 1934. [On the distribution of the parasites in Funakoshi Town on the coast of Lake Hachirogata in Akita Prefecture]. *Tokyo Iji Shinshi (Tokyo Medical Journal)*, (2909): 2889-2893. [In Japanese].
- Inukai, T., 1949. [Feeding habits of Japanese dace in the Bibi River, Hokkaido]. *Scientific Reports of the Hokkaido Fish Hatchery*, **4**: 57-61. [In Japanese].
- Ishii, S., 1914. [Survey on the tapeworm parasitic in Japanese dace in Lake Chuzenji]. *Report of the Imperial Fisheries Institute*, **9**: 283-305, 2 pls. [In Japanese].

- Ishii, S., 1931. [Parasites of Japanese fishes. 4. *Ligula* of freshwater fishes]. In: *Biology*, **18** (supplement): 190-196. Iwanami Shoten, Tokyo. [In Japanese].
- Ishizaki, D., Otake, F., Sato, T., Yodo, T., Yoshioka, M., Kashiwagi, M., 2009. Use of otolith microchemistry to estimate the migratory history of Japanese dace *Tribolodon hakonensis* in the Kamo River, Mie Prefecture. *Nippon Suisan Gakkaishi*, **75**: 419-424. [In Japanese with English abstract].
- Ito, J., 1968. Epidemiological studies of *Metagonimus yokogawai* in Shizuoka Prefecture, Japan. *Bulletin of the Faculty of Education, Shizuoka University, Natural Sciences Series*, (19): 83-92. [In Japanese with English abstract].
- Ito, J., Mochizuki, H., 1968. An epidemiologic study of human helminths in Shizuoka Prefecture VI. The metacercarial fauna in fresh and brackish water fish. *Japanese Journal of Parasitology*, **17**: 69-74. [In Japanese with English abstract].
- Kagei, N., 1966. Epidemiological studies on metagonimiasis in Japan. III. Epidemiology of *Metagonimus yokogawai* infection of Ayu, *Plecoglossus altivelis* T. & S. in Takatsu River, Shimane Prefecture. *Bulletin of the Institute of Public Health*, **15**: 38-47. [In Japanese with English abstract].
- Kanoh, Y., 1949. Ueber die Oekologie und Morphologie des japanischen Alands *Tribolodon* auf Hokkaido. *Seibutu*, **4**: 81-89. [In Japanese with German title].
- Kawamura, H., 1982. Freshwater fishes in the eastern Hidaka coast and Cape Erimo in Hokkaido, Japan. *Scientific Reports of the Hokkaido Fish Hatchery*, (37): 1-12. [In Japanese with English abstract].
- Kobayashi, H., 1912. [On a new genus of trematode]. *Saikingaku Zasshi*, (204): 780-786, 1 pl. [In Japanese].
- Kobayashi, H., 1915. Studies of endoparasitic trematodes from Japan (2). *Dobutsugaku Zasshi (Zoological Magazine)*, **27**: 50-57, 1 pl. [In Japanese].
- Kobayashi, H., 1921. One some digenetic trematodes in Japan. *Parasitology*, **12**: 380-410, 3 pls.
- Kobayashi, Y., 1972. [A research for *Metagonimus yokogawai* in the Kano basin (Metacercaria found in Ayu, *Tribolodon hakonensis hakonensis* and *Zacco platypus*)]. *Eisei Kensa*, **21**: 1006-1009. [In Japanese].
- Kobayashi, Y., Egawa, Y., Tutikawa, K., 1972. A research for *Metagonimus yokogawai* in the Kano basin. I. Number of metacercariae measured on the scales of three kinds of fish; Ayu, *Tribolodon hakonensis hakonensis* and *Zacco platypus*. *Teishin Igaku*, **24**: 775-781. [In Japanese].
- Koga, I., 1922. [On the visceral parasites, especially trematodes, studied in Kumamoto Prefecture]. *Proceedings of the Japanese Society of Pathology*, **12**: 120-149. [In Japanese].
- Koga, G., 1938. [Studies on trematodes of the genus *Metagonimus*]. *Igaku Kenkyu*, **12**: 3471-3528, 4 pls. [In Japanese].
- Koga, G., 1939. [On the second intermediate hosts of *Metagonimus* in the Chikugo River, Kyushu]. *Kyudai Iho*, **13**: 141-145. [In Japanese].
- Koga, Y., 1952. Studies on the life histories of the trematodes belonging to the family Echinostomatidae. Part 2. On the life history of a new trematode belonging to the genus *Microparyphium*. *Journal of the Kurume Medical Association*, **15**: 680-692. [In Japanese with English abstract].
- Kokame, Y., 1939. [On *Metagonimus yokogawai ovatus* using Japanese dace as the intermediate host in the Daishoji River, Ishikawa Prefecture]. *Tokyo Iji Shinshi (Tokyo Medical Journal)*, (3127): 793-796. [In Japanese].
- Komiya, Y., Suzuki, R., Kumada, M., Shiga, M., Koo, E., 1957. Epidemiological survey of liver-trematodes in the district of Lake Suwa. *Japanese Journal of Public Health*, **4**: 200-202. [In Japanese].

- Komiya, Y., Suzuki, N., Kumada, M., Fukushima, T., Kozai, I., 1960. On the distribution of *Clonorchis sinensis* around the Lake Biwa areas, Shiga Prefecture. *Japanese Journal of Parasitology*, **9**: 162-166. [In Japanese with English abstract].
- Kuchta, R., Scholz, T., 2007. Diversity and distribution of fish tapeworms of the "Bothriocephalidea" (Eucestoda). *Parassitologia*, **49**: 129-146.
- Kuchta, R., Scholz, T., Brabec, J., Bray, R. A., 2008. Suppression of the tapeworm order Pseudophyllidea (Platyhelminthes: Eucestoda) and the proposal of two new orders, Bothriocephalidea and Diphyllbothriidea. *International Journal for Parasitology*, **38**: 49-55.
- Kugi, G., Matsuo, K., 1990. A new cestode, *Coelobothrium oitense* n. sp. (Pseudophyllidea: Ptychobothriidae) from a Japanese freshwater fish, *Tribolodon hakonensis*. *Japanese Journal of Parasitology*, **39**: 255-257.
- Kurawaka, K., 1977. Cephalic lateral-line systems and geographical distribution in the genus *Tribolodon* (Cyprinidae). *Japanese Journal of Ichthyology*, **24**: 167-175.
- Loganathan, B. G., Tanabe, S., Tatsukawa, R., Ogawa, K., Goto, M., 1989. Temporal changes of morphologic abnormalities and parasitic infestation in fishes from the River Nagaragawa, Japan. *Nippon Suisan Gakkaishi*, **55**: 769-774.
- Machida, M., Araki, J., 1982. Redescription of *Pseudorhadinorhynchus leuciscus* (Krotov et Petrotschenko, 1956). *Research Bulletin of the Meguro Parasitological Museum*, (8): 49-51.
- Machida, M., Araki, J., Kamiya, H., Ohbayashi, M., 1972. Trematodes collected from sea fishes of the Hidaka District, Hokkaido. *Memoirs of the National Science Museum*, (5): 1-9. [In Japanese with English abstract].
- Miyabe, T., Takahashi, K., Inoue, M., 2007. Basic research related to artificial multiplication of *Inversidens japonensis*. *Bulletin of the Chiba Prefectural Fisheries Research Center*, (2): 53-60. [In Japanese].
- Miyaji, D., 1963. [Natural History of Animals in Fresh Waters]. Asahi Shimbun, Tokyo: 228 pp. [In Japanese].
- Miyamoto, K., 1985. Studies on zoonoses in Hokkaido 7. Survey of natural definitive hosts of *Metagonimus yokogawai*. *Japanese Journal of Parasitology*, **34**: 371-376. [In Japanese with English abstract].
- Miyamoto, K., Kutsumi, H., 1978. Studies on zoonoses in Hokkaido, Japan 2 [sic]. On the second intermediate hosts of *Metagonimus yokogawai* in Asahikawa City, Kamikawa District. *Japanese Journal of Parasitology*, **27**: 445-452. [In Japanese with English abstract].
- Miyamoto, K., Kutsumi, H., 1980. Studies on zoonoses in Hokkaido III [sic]. Prevalence rate of *Metagonimus yokogawai* metacercariae on [sic] the daces over Hokkaido. *Japanese Journal of Parasitology*, **29**: 415-422. [In Japanese with English abstract].
- Moravec, F., Nagasawa, K., 1989. Observations on some nematodes parasitic in Japanese freshwater fishes. *Folia Parasitologica*, **36**: 127-141.
- Moravec, F., Margolis, L., Boyce, N. P., 1981. Some nematodes of the genus *Rhabdochona* (Spirurida) from fishes of Japan. *Věstník Československe Společnosti Zoologické*, **45**: 277-290.
- Moravec, F., Nagasawa, K., Urawa, S., 1985. Some fish nematodes from fresh waters in Hokkaido, Japan. *Folia Parasitologica*, **32**: 305-316.
- Moravec, F., Nagasawa, K., Urushibara, Y., 1998. Observations on the seasonal maturation of the nematode *Rhabdochona zacconis* in Japanese dace, *Tribolodon hakonensis*, of the Okitsu River, Japan. *Acta Societatis Zoologicae Bohemicae*, **62**: 45-50.
- Mori, T., Urabe, M., Nagoshi, M., 1998. Relationship between body size of dark chub, *Zacco temminckii*



- and number of parasitic nematode, *Rhabdochona denudata*. *Biology of Inland Waters*, (13): 67-70. [In Japanese with English abstract].
- Nagasawa, K., 1993a. Review of human pathogenic parasites in the Japanese common squid (*Todarodes pacificus*). In: *Advances in Cephalopod Fisheries Biology*, eds., Okutani, T., O' Dor, R. K., Kubodera, T., Tokai University Press, Tokyo: 293-312.
- Nagasawa, K., 1993b. Parasites of gadid fishes in Japanese waters (Review and bibliography). *Scientific Reports of the Hokkaido Fisheries Experimental Station*, (42): 69-89. [In Japanese with English abstract].
- Nagasawa, K., 2012. A checklist and bibliography of the parasites of Manila clams (*Ruditapes philippinarum*) of Japan (1906-2012). *Bulletin of the Biogeographical Society of Japan*, **67**: 25-40. [In Japanese with English abstract].
- Nagasawa, K., Nitta, M., 2012. A checklist of the parasites of freshwater and brackish-water fishes of Hiroshima Prefecture, Japan (1925-2012). *Bulletin of Hiroshima University Museum*, **4**: 53-71. [In Japanese with English abstract].
- Nagasawa, K., Urawa, S., Awakura, T., 1987. A checklist and bibliography of parasites of salmonids of Japan. *Scientific Reports of the Hokkaido Salmon Hatchery*, (41): 1-75.
- Nagasawa, K., Urawa, S., Awakura, T., 1989. A checklist and bibliography of parasites of freshwater fishes of Hokkaido. *Scientific Reports of the Hokkaido Fish Hatchery*, (44): 1-49.
- Nagasawa, K., Takami, T., Murakami, Y., 1994. *Lepeophtheirus salmonis* (Copepoda: Caligidae) from white-spotted charr (*Salvelinus leucomaenis*), juvenile chum salmon (*Oncorhynchus keta*), and Japanese dace (*Tribolodon hakonensis*) from northern Japan. *Scientific Reports of the Hokkaido Fish Hatchery*, (48): 95-97.
- Nagasawa, K., Umino, T., Grygier, M. J., 2007a. A checklist of the parasites of ayu (*Plecoglossus altivelis altivelis*) (Salmoniformes: Plecoglossidae) in Japan (1912-2007). *Journal of the Graduate School of Biosphere Science, Hiroshima University*, **46**: 59-89.
- Nagasawa, K., Umino, T., Mizuno, K., 2007b. A checklist of the parasites of eels (*Anguilla* spp.) (Anguilliformes: Anguillidae) in Japan (1915-2007). *Journal of the Graduate School of Biosphere Science, Hiroshima University*, **46**: 91-121.
- Nagasawa, K., Yamauchi, T., Umino, T., 2008. Synopsis of leeches of the families Piscicolidae and Ozobranchidae (Annelida, Rhynchobdellida) in Japan (1895-2008). *Bulletin of the Biogeographical Society of Japan*, **63**: 151-171. [In Japanese with English abstract].
- Nagasawa, K., Morimoto, S., Asai, T., Kitagawa, T., Hosoya, K., 2012. A checklist of the parasites of medaka (*Oryzias latipes*) of Japan (1929-2012), with new records of *Lernaea cyprinacea* (Copepoda: Lernaeidae) in wild populations of medaka in Japan. *Bulletin of the Biogeographical Society of Japan*, **67**: 1-13. [In Japanese with English abstract].
- Nagasawa, K., Katahira, H., Nitta, M., 2013. *Isoparorchis hypselobagri* (Trematoda: Isoparorchidae) from freshwater fishes in western Japan, with a review of its host-parasite relationships in Japan (1915-2013). *Biogeography*, **15**: 11-20.
- Nakade, Y., 1972. Studies on trematode larvae found from fresh water snails, *Semisulcospira* spp., in Tohoku district. *Hirosaki Medical Journal*, **23**: 525-554. [In Japanese with English abstract].
- Nakamura, S., Urabe, M., Nagoshi, M., 2000. Seasonal change of prevalence and distribution of parasites in freshwater fishes at Higashi-yoshino, Nara Prefecture. *Biology of Inland Waters*, (15): 12-19. [In Japanese with English abstract].

- Nihei, E., 1961. Epidemiological studies on larval trematodes infecting the fresh-water fish in Tokyo. *National Defense Medical Journal*, **8**: 377-380. [In Japanese with English abstract].
- Nihei, E., 1962. Research on the flukes, especially *Clonorchis sinensis*, of fishes collected from main rivers in Tokyo. *Monthly Report of the Meguro Parasitological Museum*, (44): 2-4. [In Japanese].
- Nihei, E., Ichihara, A., Kamegai, Sh., 1964. On some injurious trematodes obtained from fresh-water fishes collected in the River Tone and its tributary waters. *Monthly Report of the Meguro Parasitological Museum*, (60): 2-5.
- Ochi, S., 1928. [Supplementary study on trematodes using freshwater fishes as second intermediate hosts: a species of larvae encysted in *Leuciscus hakonensis*]. *Tokyo Iji Shinshi (Tokyo Medical Journal)*, **52**: 1890-1905. [In Japanese].
- Ochi, G., 1957. [Studies on trematodes of *Metagonimus* in Japan]. *Tokyo Iji Shinshi (Tokyo Medical Journal)*, **74**: 591-599, 3 pls. [In Japanese].
- Ogawa, K., 1994. Monogenean parasites of freshwater fishes of Hokkaido, Japan. *Scientific Reports of the Hokkaido Fish Hatchery*, (48): 59-67.
- Ohnishi, Y., 1983. Experimental studies on *Metagonimus yokogawai* infection in the mouse. *Journal of the Juzen Medical Society*, **92**: 585-597. [In Japanese with English abstract].
- Ohnishi, Y., 1987. Eosinophil response in the mice infected with *Metagonimus yokogawai*. *Japanese Journal of Parasitology*, **36**: 271-275.
- Okabe, K., 1940. A synopsis of trematod [sic] cysts in fresh water fishes from Hukuoka Prefecture. *Hukuoka Acta Medica*, **33**: 81-107, 2 pls. [In Japanese with English abstract].
- Okabe, K., Kifune, T., Shiraiishi, S., 1968. Studies on the metacercaria of *Metagonimus yokogawai* obtained from some fresh-water fishes collected in the Chikugo River basin, Kyushu. *Journal of the Kurume Medical Association*, **31**: 295-304. [In Japanese with English abstract].
- Okura, T., Suzuki, S., Ootomo, Y., Tazaki, S., 1985a. [Distribution of *Diplozoon* sp. in the Arakawa River system and seasonal changes in prevalence on *Tribolodon hakonensis*]. *Bulletin of the Saitama Prefectural Fisheries Experimental Station*, (44): 82-85. [In Japanese].
- Okura, T., Suzuki, S., Ootomo, Y., 1985b. [Treatment method of *Diplozoon* sp. infecting *Tribolodon hakonensis* and changes in hematological features after treatment]. *Bulletin of the Saitama Prefectural Fisheries Experimental Station*, (44): 86-93. [In Japanese].
- Olson, P. D., Littlewood, D. T. J., Bray, R. A., Mariaux, J., 2001. Interrelationships and evolution of the tapeworms (Platyhelminthes: Cestoda). *Molecular Phylogenetics and Evolution*, **19**: 443-467.
- Oros, M., Scholz, T., Hanzelová, V., Mackiewicz, J. S., 2010. Scolex morphology of monozoic cestodes (Caryophyllidea) from the Palaearctic Region: a useful tool for species identification. *Folia Parasitologica*, **57**: 37-46.
- Oyamada, T., Esaka, Y., Kudo, N., Oyamada, T., Yoshikawa, T., Kamiya, H., 1996a. Prevalence of *Gnathostoma nipponicum* larvae in *Oncorhynchus masou* (Salmonidae) and *Tribolodon hakonensis* (Cyprinidae) collected from eastern Aomori Prefecture, Japan. *Japanese Journal of Parasitology*, **45**: 201-206.
- Oyamada, T., Esaka, Y., Kudo, N., Oyamada, T., Yoshikawa, T., 1996b. Epidemiological survey of *Gnathostoma nipponicum* larvae in fishes as the source of human infection in northern Japan. *Journal of the Japan Veterinary and Medical Association*, **49**: 574-578. [In Japanese with English abstract].
- Oyamada, T., Ikadai, H., Kudo, N., Yoshikawa, H., Oyamada, T., Yoshikawa, T., Suzuki, N., 1997. Susceptibility of several species of Cyprinidae and Salmonidae freshwater fish to larval *Gnathostoma*

- nipponicum* infection. *Journal of Veterinary Medical Science*, **59**: 1035-1037.
- Saito, S., 1968a. Studies on genus *Metagonimus* (Trematoda: Heterophyidae) I. Observations on the metacercariae of *Metagonimus* encysted in fresh water fishes in Niigata and neighbouring prefectures, especially on specific identification of them. *Niigata Igakkai Zasshi (Niigata Medical Journal)*, **82**: 679-693, 2 pls. [In Japanese].
- Saito, S., 1968b. Studies on genus *Metagonimus* (Trematoda: Heterophyidae) II. Experimental infections with the metacercariae of *Metagonimus* in mice and dogs. *Niigata Igakkai Zasshi (Niigata Medical Journal)*, **82**: 694-706, 3 pls. [In Japanese].
- Saito, S., 1984. [On the differences between the species of the genus *Metagonimus*]. *Japanese Society for Systematic Parasitology Circular*, (2): 1-4. [In Japanese].
- Saito, S., Otsuru, M., 1965a. Epidemiological surveys on the intermediate hosts of *Clonorchis sinensis* in the Hokuriku District and Yamagata Prefecture. *Medicine and Biology*, **71**: 180-185. [In Japanese].
- Saito, S., Otsuru, M., 1965b. Metacercaria of *Metagonimus yokogawai* var. *takahashii* encysted in the fishes in Niigata Prefecture. *Medicine and Biology*, **71**: 251-255. [In Japanese].
- Saito, S., Otsuru, M., Hasegawa, K., Hori, M., 1964. Some observations on metacercariae of trematodes encysted in fresh-water fishes in Niigata Prefecture, Japan. *Niigata Igakkai Zasshi (Niigata Medical Journal)*, **78**: 376-386, 2 pls. [In Japanese].
- Saito, S., Watanabe, T., Yamashita, T., 1984. Epidemiological survey of freshwater fish-borne helminths in Yamagata Prefecture. *Yamagata Medical Journal*, **2**: 71-79. [In Japanese with English abstract].
- Saito, S., Chai, J.-Y., Kim, K.-H., Lee, S.-H., Rim, H.-J., 1997. *Metagonimus miyatai* sp. nov. (Digenea: Heterophyidae), a new intestinal trematode transmitted by freshwater fishes in Japan and Korea. *Korean Journal of Parasitology*, **35**: 223-232.
- Sakai, H., 1989. *Leuciscus (Tribolodon) hakonensis*, *Leuciscus (Tribolodon)* sp., *Leuciscus (Tribolodon) brandti*, *Leuciscus (Tribolodon) ezoe*. In: *Freshwater Fishes of Japan*, eds., Kawanabe, H. and Mizuno, N., Yama-kei Publishers, Tokyo: 259-269. [In Japanese].
- Sakai, H., 1995. Life-histories and genetic divergence in three species of *Tribolodon* (Cyprinidae). *Memoirs of the Graduate School of Fisheries Sciences, Hokkaido University*, **42**: 1-98.
- Satoh, S., Komatsu, S., Tsuchiya, H., 1991. Note on the [sic] *Digramma alternans* (Rud., 1810) found on [sic] the common minnow, *Zacco platypus* in Lake Ashi-II. Infection of fishes and ecology. *Kanagawa-ken Tansuigyoryu Zosyokushiken Hokoku (Reports of the Kanagawa Prefectural Freshwater Fish Propagation Experimental Station)*, (27): 75-81. [In Japanese].
- Scholz, T., Shimazu, T., Olson, P. D., Nagasawa, K., 2001. Caryophyllidean tapeworms (Platyhelminthes: Eucestoda) from freshwater fishes in Japan. *Folia Parasitologica*, **48**: 275-288.
- Shiino, S. M., 1955. A new piscicola copepod belonging to the genus *Caligus* from Matsushima Bay. *Bulletin of the Biogeographical Society of Japan*, **16-19**: 135-140.
- Shiino, S. M., 1959. Sammlung der parasitischen Copepoden in der Präfecturuniversität von Mie. *Report of Faculty of Fisheries, Prefectural University of Mie*, **3**: 334-374.
- Shimazu, T., 1974. *Pseudozoogonoides ugui* sp. nov., a new digenetic trematode from the dace, *Tribolodon hakonensis*, from Hokkaido, Japan (Trematoda: Zoogonidae). *Bulletin of the Japanese Society of Scientific Fisheries*, **40**: 433-438.
- Shimazu, T., 1981. Some digenetic trematodes of freshwater fishes from eastern Hokkaido, Japan. *Journal of Nagano-ken Junior College*, (36): 13-26.
- Shimazu, T., 1988. Trematodes of the genus *Allocreadium* (Allocreadiidae) from freshwater fishes of Japan.

- Bulletin of the National Science Museum, Series A*, **14**: 1-21.
- Shimazu, T., 1990. Trematodes of a new genus, *Neoplagioporus* gen. sp. (Digenea: Opecoelidae: Plagioporinae), and an unidentified opecoelid from freshwater fishes of Japan. *Japanese Journal of Parasitology*, **39**: 384-396.
- Shimazu, T., 1992. Trematodes of the genera *Asymphylodora*, *Anapalaeorchis* and *Palaeorchis* (Digenea: Lissorchiidae) from freshwater fishes of Japan. *Journal of Nagano-ken Junior College*, (47): 1-19.
- Shimazu, T., 2002. Life cycle and morphology of *Metagonimus miyatai* (Digenea: Heterophyidae) from Nagano, Japan. *Parasitology International*, **51**: 271-280.
- Shimazu, T., Hashimoto, K.-I., 1999. A new species of the genus *Allocreadium* (Digenea, Allocreadiidae) from freshwater fishes of Japan. *Bulletin of the National Science Museum, Series A*, **25**: 27-31.
- Shimazu, T., Urabe, M., 2005. Digeneans found in freshwater fishes of the Uji River at Uji, Kyoto Prefecture, and the Takami River at Higashiyoshino, Nara Prefecture, Japan. *Journal of Nagano Prefectural College*, (60): 1-14.
- Shimazu, T., Urabe, M., Grygier, M. J., 2011. Digeneans (Trematoda) parasitic in freshwater fishes (Osteichthyes) of the Lake Biwa basin in Shiga Prefecture, central Honshu, Japan. *National Museum of Nature and Science Monographs*, (43): 1-105.
- Shindo, J., 1997. Parasitism of trematodes, *Diplozoon* sp., on the gills of captive cyprinid fish, *Tribolodon* sp. *Journal of Japanese Association of Zoos and Aquariums*, **38**: 88-92. [In Japanese].
- Suzuki, S., 1950. Studies on the urceolarid ciliates of Japan. *Bulletin of Yamagata University, Natural Science*, **1**: 181-218, 6 pls.
- Suzuki, S., Okura, T., 1987. [Distribution of *Diplozoon* sp. in rivers of Saitama Prefecture in 1984-1986]. *Bulletin of the Saitama Prefectural Fisheries Experimental Station*, (46): 76-87. [In Japanese].
- Suzuki, S., Okura, T., 1988. [Effects to kill the eggs of *Diplozoon* sp., a parasite of *Tribolodon hakonensis*]. *Bulletin of the Saitama Prefectural Fisheries Experimental Station*, (47): 88-90. [In Japanese].
- Takabayashi, Y., 1953. [Studies on trematodes using fishes as intermediate hosts, with special reference to a survey in Yamaguchi Prefecture]. *Acta Scholae Medicinalis in Gifu*, **1**: 219-226. [In Japanese].
- Takahashi, S., 1967. Studies on genus *Metagonimus*. *Journal of Okayama Medical Association*, **79**: 43-49. [In Japanese with English abstract].
- Taki, S., 1935. [On the pathogenicity of *Metagonimus yokogawai*]. *Acta Medica*, **9**: 1695-1704, 3 pls. [In Japanese].
- Tani, S., Ishida, K., Yoshimura, H., 1974. Epidemiological studies on metagonimiasis in Akita Prefecture. 1. The prevalence of metacercariae of *Metagonimus yokogawai* in fresh water fishes. *Akita Journal of Rural Medicine*, **21**: 25-30. [In Japanese with English abstract].
- Tsutsumi, T., 1978. [Treatment of fish diseases at aquaria. Part 7. The anchor worm, a parasite of freshwater fishes, and its control]. *Doyaku Kenkyu*, **12**: 21-22. [In Japanese].
- Uchida, A., Kawakami, Y., Kato, S., Murata, Y., 1999. Epidemiology of *Metagonimus yokogawai* metacercariae infection in natural and cultivated fresh water fishes of Kanagawa and Shizuoka Prefectures. *Journal of the Japan Veterinary Medical Association*, **52**: 115-119. [In Japanese with English abstract].
- Urawa, S., Awakura, T., 1994. Protozoan diseases of freshwater fishes in Hokkaido. *Scientific Reports of the Hokkaido Fish Hatchery*, (48): 47-58.
- Urawa, S., Kato, T., 1991. Heavy infections of *Caligus orientalis* Gussev (Copepoda: Caligidae) on caged rainbow trout *Oncorhynchus mykiss* in brackish water. *Gyobyō Kenkyū (Fish Pathology)*, **26**: 161-162.

- Yamaguti, S., 1934. Studies on the helminth fauna of Japan. Part 2. Trematodes of fishes, I. *Japanese Journal of Zoology*, **5**: 249-541.
- Yamaguti, S., 1935. Studies on the helminth fauna of Japan. Part 9. Nematodes of fishes, I. *Japanese Journal of Zoology*, **6**: 337-386.
- Yamaguti, S., 1939. Studies on the helminth fauna of Japan. Part 29. Acanthocephala, II. *Japanese Journal of Zoology*, **8**: 317-351, 9 pls.
- Yamaguti, S., 1941. Studies on the helminth fauna of Japan. Part 33. Nematodes of fishes, II. *Japanese Journal of Zoology*, **9**: 343-396, 3 pls.
- Yamauchi, T., Nagasawa, K., 2012. Redescription of the fish parasite *Nerocila japonica* Shioedte & Meinert, 1881 (Crustacea: Isopoda: Cymothoidae), with discussion on the previous records of *Nerocila acuminata* in Japan. *Systematic Parasitology*, **81**: 147-157.
- Yokogawa, M., Sano, M., Takahashi, T., Noguchi, M., Mochizuki, H., 1962. Studies on *Metagonimus yokogawai* Katsurada, 1913 in Ohama-machi, Shizuoka Prefecture. *Japanese Journal of Parasitology*, **11**: 157-164. [In Japanese with English abstract].
- Yoshimura, H., Ohmori, Y., Tani, S., Ishida, K., 1972. Epidemiological studies on metagonimiasis in Chōkai Village, Akita Prefecture. *Japanese Journal of Parasitology*, **21**: 400-407. [In Japanese with English abstract].
- Yoshimura, H., Kondo, H., Ohnishi, Y., Nishida, K., Akao, N., Okamoto, T., Ngrov, G., Minowa, M., Yamagishi, M., Tachikawa, Y., Yamada, O., Ashihara, Y., 1978. [Metagonimiasis in the Oku-noto region of Ishikawa Prefecture]. *Nippon Iji Sinpo (Japanese Medical Journal)*, (2822): 31-34. [In Japanese].
- Yuda, K., 1969. Epidemiological studies on clonorchiasis in Miyagi Prefecture, with special reference to its human infection source. *Japanese Journal of Public Health*, **16**: 611-621. [In Japanese with English abstract].

## 日本産ウグイ属魚類の寄生虫目録（1908-2013年）

長澤和也<sup>1)</sup>・片平浩孝<sup>1,2)</sup>

<sup>1)</sup> 広島大学大学院生物圏科学研究科, 〒739-8528 広島県東広島市鏡山1-4-4

<sup>2)</sup> 現住所：北海道大学地球環境科学研究院, 〒060-0810 北海道札幌市北区北10西5

**要 旨** 1908～2013年の106年間に出版された文献に基づき、日本産ウグイ属魚類4種（ウグイ *Tribolodon hakonensis*, エゾウグイ *Tribolodon sachalinensis*, マルタ *Tribolodon brandtii*, ウケクチウグイ *Tribolodon nakamuraii*）の寄生虫に関する情報を2つのリスト（寄生虫－宿主リスト, 宿主－寄生虫リスト）に整理して目録を作成した。寄生虫－宿主リストでは、44種の寄生虫と学名がまだ決定していない寄生虫の情報を、下記の高位分類群ごとに配列し、最新の学名、シノニム、寄生部位、地理的分布および報告者の情報を示した：絨毛虫類（2種：種小名まで決定している種数）、ミクソゾア類（1種）、吸虫類（18種）、単生類（0種）、条虫類（3種）、線虫類（9種）、鉤頭動物（2種）、ヒル類（1種）、軟体動物（1種）、エラオ類（0種）、カイアシ類（6種）、ワラジムシ類（1種）。宿主－寄生虫リストでは、ウグイ属魚類4種の種ごとに、各寄生虫の学名と地理的分布を示した。魚種ごとに寄生虫の種数を示すと、ウグイ43種、エゾウグイ7種、マルタ6種、ウケクチウグイ1種で、ウグイから報告されている寄生虫数が最も多かった。

**キーワード**：ウケクチウグイ、ウグイ、エゾウグイ、寄生虫、マルタ、目録