

Doctoral Thesis

Spatial distribution of radiocesium contamination
using a specific moss, *Hyophila propagulifera*
(Pottiaceae. Bryophyta) after Fukushima Daiichi
Nuclear Power Plant accident

Sharifah Nabihah Binti Syed Othman

Division of Integrated Arts and Sciences
Graduate School of Integrated Arts and Sciences
Hiroshima University

September 2018

ABSTRACT

Extensive studies were conducted on plants and soils to understand the mechanism of radionuclides uptake from the environment after the Fukushima Daiichi Nuclear Power Plant (FDNPP) accident. It is essential to measure and monitor the magnitude of radiation contamination on the environment to effectively and efficiently plan for radiation recovery. Bryophyte can be very precise and sensitive bio-indicators for heavy metal contaminations, while it can be a long-term monitoring tool if they are tolerant to the contamination. Therefore, it is important to investigate and understand the natural potential of bryophyte to control this radiation disruption. In my research, I studied the trend of radiocesium accumulation in a bryophyte, *Hyophila Propagulifera* Musci, as a bio-measuring tool focusing on the area of northeastern region of Fukushima Prefecture. *Hyophila propagulifera* was selected because of their wide distribution ranges, their ability to accumulate the radiocesium, their tendency to grow around all seasons of year and their ability to survive in dry weather or highly polluted area. Since they are commonly found on the surface of rock and roadside wall, we can minimize the influence of the soil factor, which enhance the variation of radiocesium contamination. The study may help in the monitoring field and decontamination practice effort as it can provide quick estimation of contamination by considering its relationship between the accumulation in the plant bodies and environmental factors of the surrounding. Such data can extensively help in determining the level of contamination of restricted area and understand other environmental factors that involve in radiocesium accumulation in the plant bodies.

This study will also investigate on how radiocesium contamination level detected by *H. propagulifera* correspond to air-dose rate measured by the expert group such as research group by JAEA (Japan Atomic Energy Agency). The methodology included comparing and analyzing the spatial distribution between air-dose rate and radiocesium contamination in the *H. propagulifera*, considering the different of environmental factors. The result of this study has shown the difference in spatial distribution and identified the significant influence of environmental factors between two different measurement approaches. A soil removal study (removal of soil from plant body) was conducted to determine the accuracy of the radiocesium accumulation in the plant bodies and to ensure the efficiency of washing procedure in producing consistent and reliable result in the study. All the results have presented the positive potential of *Hyophila propagulifera*, Musci as a monitoring tool to represent the contamination level in the environment.

TABLE OF CONTENTS

ABSTRACT	1
ACKNOWLEDGEMENT	3
TABLE OF CONTENTS	5
LIST OF TABLES.....	7
LIST OF FIGURES	8
LIST OF ABBREVIATIONS.....	11
CHAPTER 1: GENERAL INTRODUCTION	12
1-1 FDNPP ACCIDENT	14
1-1-1 Radionuclides emission and distribution	14
1-1-2 Environmental and radiocaesium contamination.....	15
1-2 MONITORING EFFORT IN FUKUSHIMA.....	16
1-2-1 Air dose contamination monitoring	17
1-2-2 Radiocaesium concentration in plant measurement.....	18
1-3 FOCAL SPECIES: HYOPHILA PROPAGULIFERA.....	20
1-3-1 Unique features & Previous studies.....	20
1-4 HYPOTHEIS & PROBLEM STATEMENT	21
CHAPTER 2: SPATIAL ANALYSIS OF RADIOCESIUM CONCENTRATIONS	
2-1 INTRODUCTION	25
2-2 METHOD.....	27
2-3 RESULT	30
2-4 DISCUSSION	36
2-5 SUMMARY.....	39

**CHAPTER 3: ESTIMATION OF SPATIAL DISTRIBUTION PATTERN OF AIR
DOSE RATE LEVEL AFTER FUKUSHIMA DAIICHI NUCLEAR POWER PLANT
EXPLOSION**

3-1 INTRODUCTION	40
3-2 METHOD.....	43
3-3 RESULT.....	46
3-4 DISCUSSION	53
3-5 SUMMARY.....	56

CHAPTER 4: THE EFFECT OF SOIL REMOVAL ON RADIOCESIUM

MEASUREMENTS

4-1 INTRODUCTION	57
4-2 METHOD.....	59
4-3 RESULT.....	62
4-4 DISCUSSION	65
4-5 SUMMARY.....	67

CHAPTER 5 : GENERAL DISCUSSION..... 68

REFERENCES..... 74

APPENDICES

APPENDIX A: SAMPLING ACTIVITEIS	88
APPENDIX B: WASHING PROCEDURES.....	91

論文目録

氏名 Sharifah Nabihah Binti Syed Othman 印

学位論文

論文題目 Spatial distribution of radiocesium contamination using a specific moss, *Hyophila propagulifera* (Pottiaceae, Bryophyta) after Fukushima Daiichi Nuclear Power Plant accident

公表の方法

公表年月日及び冊数 30年 6月 30日 3冊

Chapter 1	General Introduction
Chapter 2	Spatial Analysis of Radiocesium concentration (Publication 1)
Chapter 3	Estimation of spatial distribution pattern of air dose rate level after Fukushima Daiichi Nuclear Power Plant explosion (Publication 2)
Chapter 4	The effect of soil removal on radiocesium measurements (Publication 3)
Chapter 5	General Discussion References Appendices

参考論文

論文題目

Publication 1

Author :Nabihah, S., Okuda, T., Yamada, T. & Endo, S.

Title :Spatial analysis of radiocesium concentrations in *Hyophila propagulifera* (Pottiaceae, Bryophyta) within 50 km of Fukushima Daiichi Nuclear Power Plant.

Journal : Hikobia (Hikobia botanical society)

Volume :17, No.2. 101-107. 2016

Publication 2

Author : Nabihah, S. and Okuda, T.

Title : Estimation of spatial distribution pattern of air dose rate level after Fukushima Daiichi Nuclear Power Plant explosion using moss *Hyophila propagulifera*, (Pottiaceae, Bryophyta).

Journal : Hikobia (Hikobia botanical society)

Volume : Vol.17, No.4 (2018) (accepted).

Publication 3

Author : Nabihah, S. and Okuda, T.

Title : The effect of soil removal on radiocesium measurements of the moss samples from Fukushima prefecture, Japan.

Journal : Hikobia (Hikobia botanical society)

Volume : 17, No.3.201-205. 2017

備考

- (1) 論文題目が外国語の場合は、和訳を付けて、外国語、日本語の順序で列記すること。
- (2) 参考論文が 2 編以上ある場合は、列記すること。
- (3) 論文をまだ公表していないときは、公表の方法及び時期の予定を記載すること。
- (4) 用紙の規格は、A4 とし、縦にして、左横書きとすること。