

広島大学学術情報リポジトリ

Hiroshima University Institutional Repository

Title	Spatial Distribution of Radiocesium Countamination Using a Specific moss, <i>Hyophila Propagulifera</i> (Pottiaceae, Bryophyta) After Fukushima Daiichi Nuclear Power Plant Accident
Author(s)	Sharifah Nabihah Binti Syed Othman,
Citation	Bulletin of the Graduate School of Integrated Arts and Sciences, Hiroshima University. II, Studies in environmental sciences , 14 : 105 - 105
Issue Date	2019-12-31
DOI	
Self DOI	
URL	https://ir.lib.hiroshima-u.ac.jp/00048896
Right	掲載された論文，研究ノート，要旨などの出版権・著作権は広島大学大学院総合科学研究科に帰属する。 Copyright (c) 2019 Graduate School of Integrated Arts and Sciences, Hiroshima University, All rights reserved.
Relation	



Spatial Distribution of Radiocesium Countamination Using a Specific moss, *Hyophila Propagulifera* (Pottiaceae, Bryophyta) After Fukushima Daiichi Nuclear Power Plant Accident

Sharifah Nabihah Binti Syed Othman

Graduate School of Integrated Arts and Sciences, Hiroshima University

Extensive studies were conducted on plants and soils to understand the mechanism of radionuclides uptake from the environment after the Fukushima Daiichi Nuclear Power Plant (FDNPP) accident. It is essential to measure and monitor the magnitude of radiation contamination on the environment to effectively and efficiently plan for radiation recovery. Bryophyte can be very precise and sensitive bio-indicators for heavy metal contaminations, while it can be a long-term monitoring tool if they are tolerant to the contamination. Therefore, it is important to investigate and understand the natural potential of bryophyte to control this radiation disruption. In my research, I studied the trend of radiocesium accumulation in a bryophyte, *Hyophila Propagulifera* Musci, as a bio-measuring tool focusing on the area of northeastern region of Fukushima Prefecture. *Hyophila propagulifera* was selected because of their wide distribution ranges, their ability to accumulate the radiocesium, their tendency to grow around all seasons of year and their ability to survive in dry weather or highly polluted area. Since they are commonly found on the surface of rock and roadside wall, we can minimize the influence of the soil factor, which enhance the variation of radiocesium contamination. The study may help in the monitoring field and decontamination practice effort as it can provide

quick estimation of contamination by considering its relationship between the accumulation in the plant bodies and environmental factors of the surrounding. Such data can extensively help in determining the level of contamination of restricted area and understand other environmental factors that involve in radiocesium accumulation in the plant bodies.

This study will also investigate on how radiocesium contamination level detected by *H. propagulifera* correspond to air-dose rate measured by the expert group such as research group by JAEA (Japan Atomic Energy Agency). The methodology included comparing and analyzing the spatial distribution between air-dose rate and radiocesium contamination in the *H. propagulifera*, considering the different environmental factors. The result of this study has shown the difference in spatial distribution and identified the significant influence of environmental factors between two different measurement approaches. A soil removal study (removal of soil from plant body) was conducted to determine the accuracy of the radiocesium accumulation in the plant bodies and to ensure the efficiency of washing procedure in producing consistent and reliable result in the study. All the results have presented the positive potential of *Hyophila propagulifera*, Musci as a monitoring tool to represent the contamination level in the environment.