

広島大学学位請求論文

Ridge Parameters Optimization
based on Minimizing Model Selection Criterion
in Multivariate Generalized Ridge Regression

(モデル選択規準最小化に基づく多変量一般化
リッジ回帰のリッジパラメータ最適化)

2021年

広島大学大学院理学研究科

数学専攻

大石 峰暉

(広島大学 AI・データイノベーション教育研究センター 特任助教)

目 次

1. 主論文

Ridge Parameters Optimization based on Minimizing Model Selection Criterion in Multivariate Generalized Ridge Regression

(モデル選択規準最小化に基づく多変量一般化リッジ回帰のリッジパラメータ最適化)

大石 峰暉

Hiroshima Mathematical Journal (出版予定).

2. 参考論文

(1) A fast algorithm for optimizing ridge parameters in a generalized ridge regression by minimizing a model selection criterion.

Ohishi, M., Yanagihara, H. & Fujikoshi, Y.

Journal of Statistical Planning and Inference, 204 (2020) 187-205.

(2) Optimization of generalized C_p criterion for selecting ridge parameters in generalized ridge regression.

Ohishi, M., Yanagihara, H. & Wakaki, H.

Smart Innovation, Systems and Technologies, 193 (2020) 267-278.

(3) A fast optimization method for additive model via partial generalized ridge regression.

Fukui, K., Ohishi, M., Yamamura, M. & Yanagihara, H.

Smart Innovation, Systems and Technologies, 193 (2020) 279-290.

(4) Equivalence between adaptive-Lasso and generalized ridge estimators in linear regression with orthogonal explanatory variables after optimizing regularization parameters.

Ohishi, M., Yanagihara, H. & Kawano, S.

Annals of the Institute of Statistical Mathematics, 72 (6) (2020) 1501-1516.

主論文

Ridge Parameters Optimization based on Minimizing Model Selection Criterion in Multivariate Generalized Ridge Regression

Mineaki OHISHI

(Received Xxx 00, 0000)

ABSTRACT. A multivariate generalized ridge (MGR) regression provides a shrinkage estimator of the multivariate linear regression by multiple ridge parameters. Since the ridge parameters which adjust the amount of shrinkage of the estimator are unknown, their optimization is an important task to obtain a better estimator. For the univariate case, a fast algorithm has been proposed for optimizing ridge parameters based on minimizing a model selection criterion (MSC) and the algorithm can be applied to various MSCs. In this paper, we extend this algorithm to MGR regression. We also describe the relationship between the MGR estimator which is not sparse and a multivariate adaptive group Lasso estimator which is sparse, under orthogonal explanatory variables.

1. Introduction

We consider n pairs of data $\{\mathbf{y}_i, \mathbf{x}_i\}$ ($i = 1, \dots, n$), where \mathbf{y}_i is a p -dimensional vector of response variables, \mathbf{x}_i is a k -dimensional vector of explanatory variables, and n satisfies $n > \max\{p, k + 1\}$. A multivariate linear regression model is a statistical model for multiple response variables (e.g., Srivastava [19], Chap. 9; Timm [22], Chap. 4). Let $\mathbf{Y} = (\mathbf{y}_1, \dots, \mathbf{y}_n)'$ be an $n \times p$ matrix of response variables, $\mathbf{X} = (\mathbf{x}_1, \dots, \mathbf{x}_n)'$ be an $n \times k$ matrix of explanatory variables, and $\mathcal{E} = (\varepsilon_1, \dots, \varepsilon_n)'$ be an $n \times p$ matrix of error variables. Then, the multivariate linear regression model is given by

$$\mathbf{Y} = \mathbf{1}_n \boldsymbol{\mu}' + \mathbf{X} \boldsymbol{\Xi} + \mathcal{E}, \quad (1)$$

where $\mathbf{1}_n$ is an n -dimensional vector of ones, $\boldsymbol{\mu}$ is a p -dimensional vector of location parameters, and $\boldsymbol{\Xi} = (\boldsymbol{\xi}_1, \dots, \boldsymbol{\xi}_k)'$ is a $k \times p$ matrix of regression coefficients. We assume that \mathbf{X} is centralized and has full column rank, i.e., $\mathbf{X}'\mathbf{1}_n = \mathbf{0}_k$ and $\text{rank}(\mathbf{X}) = k$, and that $\varepsilon_1, \dots, \varepsilon_n$ are independently and identically distributed according to mean vector $\mathbf{0}_p$ and covariance matrix $\boldsymbol{\Sigma}$, where $\mathbf{0}_k$ is a k -dimensional vector of zeros. One of the most basic methods for estimating the unknown parameters $\boldsymbol{\mu}$ and $\boldsymbol{\Xi}$ in (1) is the least squares (LS) method. The LS

2010 *Mathematics Subject Classification.* Primary 62J07; Secondary 62H12.

Key words and phrases. Adaptive group Lasso regression, Generalized ridge regression, Model selection criterion, Multivariate analysis, Ridge parameters optimization, Sparsity.

estimators of $\boldsymbol{\mu}$ and $\boldsymbol{\Xi}$ are given by

$$\hat{\boldsymbol{\mu}} = \bar{\boldsymbol{y}} = \frac{1}{n} \boldsymbol{Y}' \mathbf{1}_n, \quad \hat{\boldsymbol{\Xi}} = \boldsymbol{M}^{-1} \boldsymbol{X}' \boldsymbol{Y} \quad (\boldsymbol{M} = \boldsymbol{X}' \boldsymbol{X}). \quad (2)$$

These estimators are equal to the maximum likelihood estimators (MLEs) of $\boldsymbol{\mu}$ and $\boldsymbol{\Xi}$ under normality, i.e., the assumption that

$$\varepsilon_1, \dots, \varepsilon_n \sim i.i.d. N_p(\mathbf{0}_p, \boldsymbol{\Sigma}).$$

The LS estimators can be obtained as simple forms as per (2) regardless of having good theoretical properties, e.g., unbiasedness and asymptotic normality. Unfortunately, it cannot be said that $\hat{\boldsymbol{\Xi}}$ is a good estimator, in the sense that the variance of the estimator becomes large when multicollinearity occurs.

For the univariate case, i.e., when $p = 1$, a generalized ridge (GR) regression was proposed by Hoerl & Kennard [10] to avoid the problem posed by multicollinearity. The GR regression can be expected to overcome this problem by shrinking an estimator of regression coefficients. The GR estimator can be obtained as closed form and the amount of shrinkage of the estimator is adjusted by k regularization parameters called ridge parameters. However, since the ridge parameters are unknown, to obtain a better estimator, we have a new problem to address, namely ridge parameters optimization. A model selection criterion (MSC) minimization method is one approach to solve the problem of ridge parameters optimization, which selects ridge parameters minimizing the MSC as the optimal ridge parameters. Most MSCs consist of a residual sum of squares (RSS) and generalized degrees of freedom (GDF). In other words, they account for model fit and model complexity. Salient examples include the C_p criterion (Mallows [12]), Akaike's information criterion (AIC; Akaike [1]) under normality, and the generalized cross-validation (GCV) criterion (Craven & Wahba [5]). Usually, the optimal parameters selected by an MSC minimization method cannot be obtained as closed forms and iterative calculation is often required. This presents difficulties in terms of the validity and applicability of such methods. Fortunately, Nagai *et al.* [14] showed that the optimal ridge parameters based on minimizing a generalized C_p (GC_p) criterion (Atkinson [3]) which is a generalization of the C_p criterion can be obtained as closed forms and Yanagihara [24] showed that the optimal ridge parameters based on minimizing the GCV criterion can be obtained as closed forms. There are various MSCs having a wide class like the GC_p criterion; for example, there are the generalized information criterion (GIC; Nishii [15]), which includes AIC, and the extended GCV (EGCV) criterion (Ohishi *et al.* [16]), which includes the GCV criterion. All these criteria can be regarded as bivariate functions of the RSS and GDF. Ohishi *et al.* [16] defined an MSC having a wider class as the bivariate function and proposed an algorithm to minimize it rapidly. Since the ridge parameters can be easily optimized by using various MSCs, the GR regression is a useful method to avoid problems arising from multicollinearity.

Ohishi *et al.* [16] also clarified a class of ridge parameters optimized by the MSC minimization method. From the results, under orthogonal explanatory

variables, the GR estimator which was previously non-sparse is now characterized by sparsity, i.e., includes 0, after the ridge parameters are optimized. On the other hand, Lasso regression (Tibshirani [20]) and adaptive Lasso (AL) regression (Zou [26]) which is an extension of the Lasso regression are well-known methods for providing a sparse estimator. They also give shrinkage estimators like the GR regression. Although the amount of shrinkage and extent of sparsity of the AL estimator (including the Lasso estimator) are adjusted by a regularization parameter called a tuning parameter, since this parameter is unknown, its optimization is required. Moreover, the AL estimator cannot usually be obtained without iterative calculation. However, Ohishi *et al.* [17] showed that the AL estimator can be obtained as closed form under orthogonal explanatory variables and the GR and AL estimators are equivalent after regularization parameters are optimized by the MSC minimization method.

Yanagihara *et al.* [25] and Nagai *et al.* [14] naturally extended the GR regression to a multivariate GR (MGR) regression. The MGR estimator is also a shrinkage estimator by k ridge parameters like the GR estimator and we have to consider the ridge parameters optimization. In the MSC minimization method for the MGR regression, although the ridge parameters optimized by the GC_p criterion minimization method can be obtained as closed forms (Nagai *et al.* [14]), whether this is the case for other criteria is unclear. Recently, Mori & Suzuki [13] proposed the ZMC_p criterion and the $ZKLIC$ which are modified versions of the modified C_p (MC_p) criterion (Fujikoshi & Satoh [8]) and the bias-corrected AIC (AIC_C ; Hurvich & Tsai [11]) for MGR regression. However, these MSCs are designed for selecting explanatory variables, not for optimizing ridge parameters. In this paper, we extend the algorithm proposed by Ohishi *et al.* [16] to MGR regression. Furthermore, we describe the relationship between MGR regression and multivariate adaptive group Lasso (MAGL) regression under orthogonal explanatory variables.

The remainder of the paper is organized as follows. In Section 2, we describe the MGR estimator and MSCs for optimizing ridge parameters, and define an MSC class. In Section 3, we extend the algorithm proposed by Ohishi *et al.* [16] to optimize ridge parameters in MGR regression by the MSC minimization method. In Section 4, the MSC class defined in Section 2 is extended, corresponding to various distances. Moreover, we propose an algorithm for minimizing the extended MSC. In Section 5, we propose a new method for optimizing ridge parameters by using MSCs. In Section 6, we describe the MAGL estimator and equivalence between the MGR and MAGL estimators under the regularization parameters optimized by the MSC minimization method. In Section 7, the performance of the ridge parameters optimized by the MSC minimization methods is compared by simulation. Technical details are provided in the Appendix.

2. Preliminaries

By a singular value decomposition, $n \times n$ and $k \times k$ orthogonal matrices \mathbf{P} and \mathbf{Q} and a $k \times k$ diagonal matrix $\mathbf{D} = \text{diag}(d_1, \dots, d_k)$ express \mathbf{X} as

$$\mathbf{X} = \mathbf{P} \begin{pmatrix} \mathbf{D}^{1/2} \\ \mathbf{O}_{n-k,k} \end{pmatrix} \mathbf{Q}' = \mathbf{P}_1 \mathbf{D}^{1/2} \mathbf{Q}', \quad (3)$$

where $\mathbf{O}_{n,k}$ is an $n \times k$ matrix of zeros, \mathbf{P}_1 is an $n \times k$ matrix obtained from the partition $\mathbf{P} = (\mathbf{P}_1, \mathbf{P}_2)$, which satisfies $\mathbf{P}_1' \mathbf{1}_n = \mathbf{0}_k$ and $\mathbf{P}_1' \mathbf{P}_1 = \mathbf{I}_k$, and d_1, \dots, d_k are eigenvalues of $\mathbf{M} (= \mathbf{X}'\mathbf{X})$ satisfying $d_1 \geq \dots \geq d_k > 0$. Then, the MGR estimators of $\boldsymbol{\mu}$ and $\boldsymbol{\Xi}$ are given by

$$\hat{\boldsymbol{\mu}} = \bar{\mathbf{y}}, \quad \hat{\boldsymbol{\Xi}}_{\boldsymbol{\theta}} = \mathbf{M}_{\boldsymbol{\theta}}^{-1} \mathbf{X}'\mathbf{Y} \quad (\mathbf{M}_{\boldsymbol{\theta}} = \mathbf{M} + \mathbf{Q}\boldsymbol{\Theta}\mathbf{Q}'), \quad (4)$$

where $\boldsymbol{\theta} = (\theta_1, \dots, \theta_k)'$, $\boldsymbol{\Theta} = \text{diag}(\theta_1, \dots, \theta_k)$ and $\theta_j \in \mathbb{R}_+ = \{\theta \in \mathbb{R} \mid \theta \geq 0\}$ ($j = 1, \dots, k$) is a regularization parameter called a ridge parameter. Since $\mathbf{M}_{\boldsymbol{\theta}} = \mathbf{M}$ when $\boldsymbol{\theta} = \mathbf{0}_k$, $\hat{\boldsymbol{\Xi}}_{\boldsymbol{\theta}}$ coincides with $\hat{\boldsymbol{\Xi}}$ in (2) when $\boldsymbol{\theta} = \mathbf{0}_k$ and the MGR estimators coincide with the GR estimators when $p = 1$. The MGR estimators in (4) denote the minimizers of the following penalized RSS (PRSS):

$$\text{tr} \{ (\mathbf{Y} - \mathbf{1}_n \boldsymbol{\mu}' - \mathbf{X}\boldsymbol{\Xi})' (\mathbf{Y} - \mathbf{1}_n \boldsymbol{\mu}' - \mathbf{X}\boldsymbol{\Xi}) + \boldsymbol{\Xi}' \mathbf{Q} \boldsymbol{\Theta} \mathbf{Q}' \boldsymbol{\Xi} \}. \quad (5)$$

Although the ridge parameters adjust the amount of shrinkage of the MGR estimator of $\boldsymbol{\Xi}$, since they are unknown, their optimization is an important task to obtain a better estimator. To simplify calculation, following Yanagihara [24] and Ohishi *et al.* [16], we transform the ridge parameters as

$$\delta_j = \frac{\theta_j}{d_j + \theta_j} \in [0, 1] \quad (j = 1, \dots, k).$$

Since this transformation is a one-to-one correspondence, the optimization of θ_j is equal to that of δ_j . Hence, we optimize δ_j instead of θ_j and we also call δ_j a ridge parameter in this paper. Let $\boldsymbol{\delta}$ and $\boldsymbol{\Delta}$ be a k -dimensional vector and a $k \times k$ diagonal matrix of the ridge parameters defined by $\boldsymbol{\delta} = (\delta_1, \dots, \delta_k)'$ and $\boldsymbol{\Delta} = \text{diag}(\delta_1, \dots, \delta_k)$, respectively, and let \mathbf{Z} be a $k \times p$ matrix defined by

$$\mathbf{Z} = (\mathbf{z}_1, \dots, \mathbf{z}_k)' = \mathbf{P}_1' \mathbf{Y}. \quad (6)$$

Then, the MGR estimator of $\boldsymbol{\Xi}$ in (4) can be rewritten as

$$\hat{\boldsymbol{\Xi}}_{\boldsymbol{\delta}} = \mathbf{Q}(\mathbf{I}_k - \boldsymbol{\Delta})\mathbf{D}^{-1/2}\mathbf{Z} = \hat{\boldsymbol{\Xi}} - \mathbf{Q}\boldsymbol{\Delta}\mathbf{D}^{-1/2}\mathbf{Z}. \quad (7)$$

In this paper, we optimize the ridge parameter $\boldsymbol{\delta}$ by using the MSC minimization method.

The MGR estimator in (7) gives a predictive matrix of \mathbf{Y} as

$$\hat{\mathbf{Y}}_{\boldsymbol{\delta}} = \mathbf{1}_n \hat{\boldsymbol{\mu}}' + \mathbf{X} \hat{\boldsymbol{\Xi}}_{\boldsymbol{\delta}} = \mathbf{H}_{\boldsymbol{\delta}} \mathbf{Y}, \quad \mathbf{H}_{\boldsymbol{\delta}} = \mathbf{J}_n + \mathbf{P}_1(\mathbf{I}_k - \boldsymbol{\Delta})\mathbf{P}_1',$$

where $\mathbf{J}_n = \mathbf{1}_n \mathbf{1}_n' / n$ and $\mathbf{H}_{\boldsymbol{\delta}}$ is an $n \times n$ matrix called a hat matrix. Most MSCs consist of the predictive matrix and the hat matrix. The predictive matrix is

used to evaluate model fit. We define an estimator and an unbiased estimator of the covariance matrix Σ as

$$\begin{aligned}\hat{\Sigma}(\delta) &= \frac{1}{n}(\mathbf{Y} - \hat{\mathbf{Y}}_\delta)'(\mathbf{Y} - \hat{\mathbf{Y}}_\delta), \\ \mathbf{S} &= \frac{1}{b}\hat{\Sigma}_0 \quad \left(\hat{\Sigma}_0 = \hat{\Sigma}(\mathbf{0}_k), b = 1 - (k+1)/n\right).\end{aligned}\tag{8}$$

Under normality, $\hat{\Sigma}(\delta)$ is a penalized MLE of Σ and $\hat{\Sigma}_0$ is an MLE of Σ . Then, model fit, i.e., the distance between \mathbf{Y} and $\hat{\mathbf{Y}}_\delta$ is defined by

$$\text{tr} \left\{ \hat{\Sigma}(\delta) \mathbf{S}^{-1} \right\}.$$

On the other hand, the hat matrix is used to evaluate model complexity and it is defined by the following GDF:

$$\text{df}(\delta) = p \text{tr}(\mathbf{H}_\delta).\tag{9}$$

The GC_p and EGCV criteria for optimizing ridge parameters consist of $\text{tr} \left\{ \hat{\Sigma}(\delta) \mathbf{S}^{-1} \right\}$ and $\text{df}(\delta)$. Similar to Yanagihara [24], we have the following lemma about $\hat{\Sigma}(\delta)$ and $\text{df}(\delta)$.

LEMMA 1. Let \mathbf{B}_δ and \mathbf{W} be $p \times p$ matrices defined by

$$\mathbf{B}_\delta = \mathbf{Z}' \Delta^2 \mathbf{Z}, \quad \mathbf{W} = n \hat{\Sigma}_0.$$

Then, $\hat{\Sigma}(\delta)$ and $\text{df}(\delta)$ can be partitioned into terms which do and do not include δ as follows:

$$\begin{aligned}\hat{\Sigma}(\delta) &= \frac{1}{n}(\mathbf{W} + \mathbf{B}_\delta) = \hat{\Sigma}_0 + \frac{1}{n} \sum_{j=1}^k \mathbf{z}_j \mathbf{z}_j' \delta_j^2, \\ \text{df}(\delta) &= p(1+k) - p \text{tr} \Delta = p \left\{ (1+k) - \sum_{j=1}^k \delta_j \right\}.\end{aligned}$$

From Lemma 1, we have

$$\text{tr} \left\{ \hat{\Sigma}(\delta) \mathbf{S}^{-1} \right\} = b \text{tr}(\mathbf{B}_\delta^*) + bp, \quad \mathbf{B}_\delta^* = \mathbf{W}^{-1/2} \mathbf{B}_\delta \mathbf{W}^{-1/2}.$$

Then, the GC_p and EGCV criteria for optimizing ridge parameters are defined by

$$\begin{aligned}GC_p(\delta) &= nb \text{tr}(\mathbf{B}_\delta^*) + nbp + \alpha \text{df}(\delta), \\ \text{EGCV}(\delta) &= \frac{b \text{tr}(\mathbf{B}_\delta^*) + bp}{\{1 - \text{df}(\delta)/np\}^\alpha},\end{aligned}$$

where α is a positive value adjusting the strength of the penalty for model complexity. Existing criteria are expressed by changing the value of α , for example, the GC_p and EGCV criteria coincide with the C_p and GCV criteria, respectively, when $\alpha = 2$ and the GC_p criterion coincides with the MC_p criterion (Yanagihara *et al.* [25]) when $\alpha = 2\{1 + (p+1)/(n-k-p-2)\}$. From the above, MSCs

for optimizing ridge parameters can be regarded as bivariate functions of $\text{tr}(\mathbf{B}_\delta^*)$ and $\text{df}(\delta)$. From Lemma 1, ranges of $\text{tr}(\mathbf{B}_\delta^*)$ and $\text{df}(\delta)$ are given as the following lemma.

LEMMA 2. *The $\text{tr}(\mathbf{B}_\delta^*)$ and $\text{df}(\delta)$ are included in the following ranges:*

$$\text{tr}(\mathbf{B}_\delta^*) \in [0, \text{tr}(\mathbf{Z}^* \mathbf{Z}^{*\prime})], \quad \text{df}(\delta) \in [p, p(1+k)],$$

where $\mathbf{Z}^* = \mathbf{Z}\mathbf{W}^{-1/2}$.

Moreover, let f be a bivariate function defined by the following class.

DEFINITION 1 (**Class of the bivariate function f**). For a positive value r_+ , f satisfies the following conditions:

- (A1) For any $(r, u) \in [0, r_+] \times [p, np]$, $f(r, u)$ is continuous.
- (A2) For any $(r, u) \in [0, r_+] \times [p, np]$, $f(r, u)$ is first order partially differentiable and its partial derivatives are positive.

We define an MSC for optimizing ridge parameters by using f in Definition 1 as

$$\text{MSC}(\delta) = f(\text{tr}(\mathbf{B}_\delta^*), \text{df}(\delta)). \quad (10)$$

For the GC_p and EGCV criteria, f is given by

$$f(r, u) = \begin{cases} f_{GC_p}(r, u) = nb(r+p) + \alpha u & (GC_p \text{ criterion}) \\ f_{EGCV}(r, u) = b(r+p)/(1-u/np)^\alpha & (EGCV \text{ criterion}) \end{cases},$$

and r_+ is given by

$$r_+ = \text{tr}(\mathbf{Z}^* \mathbf{Z}^{*\prime}).$$

Then, the optimal ridge parameters based on minimizing the MSC in (10) are given by

$$\hat{\delta} = (\hat{\delta}_1, \dots, \hat{\delta}_k)' = \arg \min_{\delta \in [0,1]^k} \text{MSC}(\delta).$$

3. Fast Optimization of Ridge Parameters

In this section, to obtain δ minimizing the MSC in (10), we extend the algorithm for optimizing ridge parameters in the GR regression proposed by Ohishi [16]. First, we define the following class of ridge parameters.

DEFINITION 2 (**Class of ridge parameters**). For $h \in \mathbb{R}_+$, a class of ridge parameters is defined by

$$\hat{\delta}(h) = (\hat{\delta}_1(h), \dots, \hat{\delta}_k(h))', \quad \hat{\delta}_j(h) = 1 - \text{soft}(1, h/\mathbf{z}'_j \mathbf{S}^{-1} \mathbf{z}_j),$$

where \mathbf{z}_j is the p -dimensional vector defined by (6). Furthermore, $\text{soft}(x, a)$ is a soft-thresholding operator (e.g., Donoho & Johnstone [7]), i.e., $\text{soft}(x, a) = \text{sign}(x)(|x| - a)_+$, and $(x)_+ = \max\{x, 0\}$.

When $\mathbf{S} = \mathbf{I}_p$ and $p = 1$, the class of ridge parameters in Definition 2 corresponds to that for the GR regression defined by Ohishi *et al.* [16]. Using this class, the MGR estimator in (7) is given as a function of h :

$$\hat{\boldsymbol{\Xi}}_{\hat{\delta}(h)} = \mathbf{Q}\mathbf{V}(h)\mathbf{Q}'\hat{\boldsymbol{\Xi}},$$

where \mathbf{Q} is the $k \times k$ orthogonal matrix defined by (3) and $\mathbf{V}(h)$ is a $k \times k$ diagonal matrix which has the following diagonal elements:

$$v_j(h) = 1 - \hat{\delta}_j(h) = \text{soft}(1, h/\mathbf{z}'_j\mathbf{S}^{-1}\mathbf{z}_j) \quad (j = 1, \dots, k).$$

The $\mathbf{V}(h)$ rewrites the predictive matrix of \mathbf{Y} as

$$\hat{\mathbf{Y}}_{\hat{\delta}(h)} = \{\mathbf{J}_n + \mathbf{P}_1\mathbf{V}(h)\mathbf{P}'_1\}\mathbf{Y},$$

where \mathbf{P}_1 is the $n \times k$ matrix defined by (3). Then, the ridge parameters optimized by the MSC minimization method are given by the following theorem (the proof is given in Appendix A.1).

THEOREM 1. *We define r_+ as*

$$r_+ = \text{tr}(\mathbf{Z}^*\mathbf{Z}'^*).$$

For f with the class in Definition 1, let $\phi(h)$ ($h \in \mathbb{R}_+ \setminus \{0\}$) be a function defined by

$$\phi(h) = \text{MSC}(\hat{\boldsymbol{\delta}}(h)),$$

and suppose that $\exists \nu > 0$ s.t. $\phi(\nu) < \lim_{h \rightarrow 0} \phi(h)$. Then, the ridge parameters optimized by the MSC minimization method are given by $\hat{\boldsymbol{\delta}}(\hat{h})$ and \hat{h} is given by

$$\hat{h} = \arg \min_{h \in \mathbb{R}_+ \setminus \{0\}} \phi(h).$$

From this theorem, the class of ridge parameters in Definition 2 is the class of the “optimal” ridge parameters.

Let t_j ($j = 1, \dots, k$) be the j th order statistic of $\mathbf{z}'_1\mathbf{S}^{-1}\mathbf{z}_1, \dots, \mathbf{z}'_k\mathbf{S}^{-1}\mathbf{z}_k$ and R_j ($j = 0, 1, \dots, k$) be a range defined by

$$R_j = \begin{cases} (0, t_1] & (j = 0) \\ (t_j, t_{j+1}] & (j = 1, \dots, k-1) \\ (t_k, \infty] & (j = k) \end{cases}. \quad (11)$$

Then, similar to Ohishi *et al.* [16], we have the following proposition.

PROPOSITION 1. *The $\phi(h)$ in Theorem 1 satisfies the following properties:*

- (P1) *For all $h \in \mathbb{R}_+ \setminus \{0\}$, $\phi(h)$ is continuous.*
- (P2) *For all $h \geq t_k$, $\phi(h) = f(r_+, p)$.*
- (P3) *The $\phi(h)$ can be expressed as the following piecewise function:*

$$\begin{aligned} \phi(h) &= \phi_a(h) \quad (h \in R_a; a = 0, 1, \dots, k) \\ &= f((c_{1,a} + c_{2,a}h^2)/nb, p(1 + k - a - c_{2,a}h)), \end{aligned}$$

where $c_{1,a}$ and $c_{2,a}$ are nonnegative constants given by

$$c_{1,a} = \begin{cases} 0 & (a = 0) \\ \sum_{j=1}^a t_j & (a = 1, \dots, k) \end{cases}, \quad c_{2,a} = \begin{cases} \sum_{j=a+1}^k \frac{1}{t_j} & (a = 0, 1, \dots, k-1) \\ 0 & (a = k) \end{cases}.$$

From the results, the MSC minimization problem for optimizing ridge parameters in the MGR regression can be solved by applying the fast algorithm for the GR regression proposed by Ohishi *et al.* [16]. That is, we have the following theorem.

THEOREM 2. *Suppose that the derivative of $\phi_a(h)$ in Proposition 1 is expressed as*

$$\frac{d}{dh}\phi_a(h) = \chi_a(h)\psi_a(h) \quad (h \in R_a; a = 0, 1, \dots, k-1),$$

and $\psi(h) = \psi_a(h)$ ($h \in R_a$) is continuous for all $h \in \mathbb{R}_+ \setminus \{0\}$, where $\chi_a(h)$ is a positive function and $\psi_a(h)$ is a polynomial. Moreover, suppose that $\exists \nu > 0$ s.t. $\phi(\nu) < \lim_{h \rightarrow 0} \phi(h)$ and let h_a be a root of $\psi_a(h) = 0$ satisfying

$$\exists \epsilon_a > 0 \text{ s.t. } \forall \epsilon \in (0, \epsilon_a), \psi_a(h_a - \epsilon) < 0. \quad (12)$$

Then, minimizer candidates of $\phi(h)$ are given by

$$\mathcal{S} = \left\{ \bigcup_{a \in \mathcal{A}} \{h_a\} \right\} \cup \mathcal{T},$$

$$\mathcal{A} = \{a \in \{0, 1, \dots, k-1\} \mid h_a \in R_a\}, \quad \mathcal{T} = \begin{cases} \{t_k\} & (\psi_{k-1}(t_k) < 0) \\ \emptyset & (\psi_{k-1}(t_k) \geq 0) \end{cases}.$$

Hence, the ridge parameters optimized by the MSC minimization method are given by $\hat{\delta}(\hat{h})$ and \hat{h} is given by

$$\hat{h} = \arg \min_{h \in \mathcal{S}} \phi(h).$$

Although the range of h is a set of positive values, Theorem 2 can reduce a search range of h to \mathcal{S} which is a set of discrete points. Furthermore, each element of \mathcal{S} is given as closed form and $\#(\mathcal{S}) \leq k+1$; hence we can quickly optimize the ridge parameters. In the theorem, although $\psi_a(h)$ is implicitly supposed as a linear or quadratic function, the theorem can naturally be extended to higher order polynomial functions. In particular, roots of $\psi_a(h) = 0$ can be obtained as closed forms when $\psi_a(h)$ is a cubic or a quartic function, by using Cardano's formula (e.g., David [6], Chap. 1) or Ferrari's method (e.g., Tignol [21], Chap. 3). Hence, if the degree of $\psi_a(h)$ is four or less, we can quickly optimize the MSC.

3.1. Examples. In this subsection, we provide specific examples of the MSC minimization methods for optimizing ridge parameters in the MGR regression. To emphasize that the optimal ridge parameters depend on α , we specify that α is given.

3.1.1. The GC_p criterion. Although the ridge parameters optimized by the GC_p criterion minimization method have already been given by Nagai *et al.* [14], here we show how to derive them by applying Theorem 2. The GC_p criterion for optimizing ridge parameters is given by

$$GC_p(\boldsymbol{\delta} \mid \alpha) = f_{GC_p}(\text{tr}(\mathbf{B}_{\boldsymbol{\delta}}^*), \text{df}(\boldsymbol{\delta}) \mid \alpha).$$

When $h \in R_a$ ($a = 0, 1, \dots, k$), ϕ and its derivative are given by

$$\begin{aligned} \phi(h \mid \alpha) &= \phi_a(h \mid \alpha) = c_{2,a}h^2 - \alpha pc_{2,a}h + nbp + c_{1,a} + \alpha p(1 + k - a), \\ \frac{d}{dh}\phi_a(h \mid \alpha) &= c_{2,a}(2h - \alpha p). \end{aligned}$$

Hence, the ridge parameters optimized by the GC_p criterion minimization method are given as the following closed form:

$$\hat{\boldsymbol{\delta}} = \hat{\boldsymbol{\delta}}(\hat{h}_\alpha), \quad \hat{h}_\alpha = \frac{\alpha p}{2}.$$

3.1.2. The EGCV criterion. The EGCV criterion for optimizing ridge parameters is given by

$$\text{EGCV}(\boldsymbol{\delta} \mid \alpha) = f_{\text{EGCV}}(\text{tr}(\mathbf{B}_{\boldsymbol{\delta}}^*), \text{df}(\boldsymbol{\delta}) \mid \alpha).$$

When $h \in R_a$ ($a = 0, 1, \dots, k$), ϕ and its derivative are given by

$$\begin{aligned} \phi(h \mid \alpha) &= \phi_a(h \mid \alpha) = \frac{bp + (c_{1,a} + c_{2,a}h^2)/n}{\{b + (a + c_{2,a}h)/n\}^\alpha}, \\ \frac{d}{dh}\phi_a(h \mid \alpha) &= \frac{c_{2,a}}{n^2\{b + (a + c_{2,a}h)/n\}^{\alpha+1}}\psi_a(h \mid \alpha), \\ \psi_a(h \mid \alpha) &= -(\alpha - 2)c_{2,a}h^2 + 2(a + nb)h - \alpha(nbp + c_{1,a}). \end{aligned}$$

When $\alpha = 2$, i.e., using the GCV criterion minimization method, we have

$$\psi_a(h \mid 2) = 2\{(a + nb)h - nbp - c_{1,a}\},$$

and a root of $\psi_a(h \mid 2) = 0$ is

$$h_a = \frac{nbp + c_{1,a}}{a + nb}.$$

Moreover, similar to Yanagihara [24], the following statement is true:

$$\exists! a^* \in \{0, 1, \dots, k - 1\} \text{ s.t. } h_{a^*} \in R_{a^*}.$$

Hence, the ridge parameters optimized by the GCV criterion minimization method are given by the following closed forms: $\hat{\boldsymbol{\delta}} = \hat{\boldsymbol{\delta}}(h_{a^*})$.

When $\alpha > 2$, since $\psi_a(h \mid \alpha)$ is a concave quadratic function, a root of $\psi_a(h \mid \alpha) = 0$ satisfying the condition (12) is given by

$$h_{\alpha,a} = \frac{(a + nb) - \sqrt{(a + nb)^2 - \alpha(\alpha - 2)c_{2,a}(nbp + c_{1,a})}}{(\alpha - 2)c_{2,a}}.$$

Therefore, candidates of \hat{h}_α are given by

$$\mathcal{S}_\alpha = \left\{ \bigcup_{a \in \mathcal{A}_\alpha} \{h_{\alpha,a}\} \right\} \cup \mathcal{T}_\alpha,$$

where \mathcal{A}_α and \mathcal{T}_α are sets given by

$$\begin{aligned} \mathcal{A}_\alpha &= \{a \in \{0, 1, \dots, k-1\} \mid h_{\alpha,a} \in R_a\}, \\ \mathcal{T}_\alpha &= \begin{cases} \{t_k\} & (r_+ > 2(1 - n^{-1})t_k/\alpha b - p) \\ \emptyset & (r_+ \leq 2(1 - n^{-1})t_k/\alpha b - p) \end{cases}. \end{aligned}$$

Hence, the ridge parameters optimized by the EGCV criterion minimization method are given by

$$\hat{\boldsymbol{\delta}} = \hat{\boldsymbol{\delta}}(\hat{h}_\alpha), \quad \hat{h}_\alpha = \arg \min_{h \in \mathcal{S}_\alpha} \phi(h \mid \alpha).$$

In the EGCV criterion minimization method, the number of minimizer candidates is only $k + 1$ at most.

3.2. Relationships between the Optimal Ridge Parameters. This subsection provides some theoretical properties concerning the relationships between the optimal ridge parameters. The class of the optimal ridge parameters satisfies

$$\forall h_1, h_2 \in \mathbb{R}_+, h_1 < h_2 \Rightarrow \hat{\delta}_j(h_1) \leq \hat{\delta}_j(h_2) \quad (j = 1, \dots, k),$$

with equality only when $h_1 \geq t_k$. This fact yields some relationships concerning the ridge parameters optimized by the GC_p and EGCV criteria minimization methods. Immediately, we have the following result which is similar to Nagai *et al.* [14].

PROPOSITION 2. *For positive values α_1 and α_2 , we define the ridge parameters optimized by the GC_p criterion minimization method as*

$$\hat{\delta}_{1,j} = \hat{\delta}_j(\hat{h}_{\alpha_1}), \quad \hat{\delta}_{2,j} = \hat{\delta}_j(\hat{h}_{\alpha_2}) \quad (j = 1, \dots, k),$$

where $\hat{h}_\alpha = \alpha p/2$. Then, we have

$$\alpha_1 < \alpha_2 \Rightarrow \hat{\delta}_{1,j} \leq \hat{\delta}_{2,j}.$$

This proposition states that the stronger the penalty for model complexity, the larger the amount of shrinkage of the estimator, when using the GC_p criterion minimization method. Next, we consider the ridge parameters optimized by the GC_p and the GCV criteria minimization methods. Similar to Yanagihara [24], we have the following lemma.

LEMMA 3. *The h_{α^*} obtained by the GCV criterion minimization method satisfies $h_{\alpha^*} \leq p$.*

This lemma leads to the following result which is similar to the case when $p = 1$ (Yanagihara [24]).

PROPOSITION 3. *Let $\hat{\delta}_{\alpha,j}^{GC_p}$ and $\hat{\delta}_j^{GCV}$ ($j = 1, \dots, k$) be the ridge parameters optimized by the GC_p and GCV criteria minimization methods, respectively. Then, we have*

$$\alpha \geq 2 \Rightarrow \hat{\delta}_j^{GCV} \leq \hat{\delta}_{\alpha,j}^{GC_p}.$$

The value of α in the MSC is often 2 or more. This means that the ridge parameters optimized by the GC_p criterion minimization method shrink the estimator more than the GCV criterion minimization method in most cases. Finally, we consider the ridge parameters optimized by the EGCV criterion minimization method. We express $\phi(h | \alpha) = \text{EGCV}(\hat{\boldsymbol{\delta}}(h) | \alpha)$ as

$$\phi(h | \alpha) = \hat{\sigma}^2(h)\eta(h | \alpha),$$

where

$$\begin{aligned} \hat{\sigma}^2(h) &= bp + b \text{tr}(\mathbf{B}_{\hat{\boldsymbol{\delta}}}^*), \\ \eta(h | \alpha) &= \frac{1}{\{1 - \text{df}(h)/np\}^\alpha}, \quad \text{df}(h) = \text{df}(\hat{\boldsymbol{\delta}}(h)), \end{aligned}$$

and let \hat{h}_α be the minimizer of $\phi(h | \alpha)$. Then, $\eta(h | \alpha)$ has the following property (the proof is given in Appendix A.2).

LEMMA 4. *Suppose that $0 < h_1 < h_2$. Then, we have*

$$\eta(h_2 | \alpha) \leq \eta(h_1 | \alpha).$$

This lemma leads to the following proposition (the proof is given in Appendix A.3).

PROPOSITION 4. *The EGCV criterion minimization method has the following properties:*

(1) *Suppose that $\alpha_1 < \alpha_2$. Then, we have*

$$\hat{h}_{\alpha_1} = t_k \Rightarrow \hat{h}_{\alpha_2} = t_k.$$

(2) *For positive values α_1 and α_2 , we define the ridge parameters optimized by the EGCV criterion minimization method as*

$$\hat{\delta}_{1,j} = \hat{\delta}_j(\hat{h}_{\alpha_1}), \quad \hat{\delta}_{2,j} = \hat{\delta}_j(\hat{h}_{\alpha_2}) \quad (j = 1, \dots, k),$$

and suppose that $\hat{h}_{\alpha_2} \neq t_k$. Then, we have

$$\alpha_1 < \alpha_2 \Rightarrow \hat{\delta}_{1,j} \leq \hat{\delta}_{2,j},$$

with equality only when $\hat{h}_{\alpha_1} \geq \mathbf{z}'_j \mathbf{S}^{-1} \mathbf{z}_j$.

This proposition states that the stronger the penalty for model complexity, the larger the amount of shrinkage of the estimator, when using the EGCV criterion minimization method.

4. Extending the MSC Class

In the previous section, we showed that the algorithm for the GR regression can be applied to minimize the MSC in (10), where the distance between \mathbf{Y} and $\hat{\mathbf{Y}}_{\delta}$ is defined by $\text{tr}\{\hat{\Sigma}(\delta)\mathbf{S}^{-1}\}$ and the MSC is defined by using $\text{tr}(\mathbf{B}_{\delta}^*)$ obtained from the distance. In this section, we focus on how to measure the distance.

Let g be a real-valued function defined by the following class.

DEFINITION 3 (Class of the function g). For any $p \times p$ positive definite matrix \mathbf{A} , the g satisfies the following conditions:

- (A1) The $g(\mathbf{A})$ is positive.
- (A2) The $\partial g(\mathbf{A})/\partial \mathbf{A}$ is a positive definite.

Using the function g , we extend the MSC in (10) to

$$\text{MSC}(\delta | g) = f(g(\mathbf{B}_{\delta}^*), \text{df}(\delta)), \quad (13)$$

where f is the bivariate function given by Definition 1. For example, g includes the following functions:

$$g(\mathbf{A}) = \begin{cases} g_{\text{LH}}(\mathbf{A}) = \text{tr}(\mathbf{A}) & \text{(LH-distance)} \\ g_{\text{LR}}(\mathbf{A}) = \log |\mathbf{I}_p + \mathbf{A}| & \text{(LR-distance)} \\ g_{\text{BNP}}(\mathbf{A}) = \text{tr} \{ \mathbf{A}(\mathbf{I}_p + \mathbf{A})^{-1} \} & \text{(BNP-distance)} \\ g_{\text{ML}}(\mathbf{A}) = \text{tr} \{ (\mathbf{I}_p + \mathbf{A})^{-1} \} + \log |\mathbf{I}_p + \mathbf{A}| - p & \text{(ML-distance)} \\ g_{\text{GLS}}(\mathbf{A}) = \text{tr}(\mathbf{A}^2)/2 & \text{(GLS-distance)} \end{cases} .$$

The MSC in (13) is equal to that in (10) when $g(\mathbf{A}) = g_{\text{LH}}(\mathbf{A})$ and the following equation holds:

$$g_{\text{LH}}(\mathbf{B}_{\delta}^*) = \text{tr}(\mathbf{B}_{\delta}\mathbf{W}^{-1}).$$

Since we can regard \mathbf{B}_{δ} as a between-group variation matrix and \mathbf{W} as a within-group variation matrix, $g_{\text{LH}}(\mathbf{B}_{\delta}^*)$ is a Lawley-Hotelling trace criterion (LH-statistic; e.g., Anderson [2], Chap. 8) which is a well-known statistic in multivariate analysis. That is, the MSC in (10) measures the distance between \mathbf{Y} and $\hat{\mathbf{Y}}_{\delta}$ based on the LH-statistic. Similarly, regarding the LR-distance and the BNP-distance, the following equations hold:

$$\begin{aligned} g_{\text{LR}}(\mathbf{B}_{\delta}^*) &= \log |(\mathbf{W} + \mathbf{B}_{\delta})\mathbf{W}^{-1}|, \\ g_{\text{BNP}}(\mathbf{B}_{\delta}^*) &= \text{tr} \{ \mathbf{B}_{\delta}(\mathbf{W} + \mathbf{B}_{\delta})^{-1} \}. \end{aligned}$$

They are a Likelihood-Ratio criterion and a Bartlett-Nanda-Pillai trace criterion, respectively, which are also well-known statistics (e.g., Anderson [2], Chap. 8). MSC based on the LR-distance includes the GIC and the AIC_C under normality. The above three distances based on the three statistics pertain to the

mean structure of a model. In contrast, there are distances with respect to the covariance structure of a model, e.g., the ML-distance and the GLS-distance. Regarding these distances, the following equations hold:

$$g_{\text{ML}}(\mathbf{B}_{\delta}^*) = \log \left| \hat{\Sigma}(\delta) \right| + \text{tr} \left\{ \hat{\Sigma}(\delta)^{-1} \hat{\Sigma}_0 \right\} - \log \left| \hat{\Sigma}_0 \right| - p,$$

$$g_{\text{GLS}}(\mathbf{B}_{\delta}^*) = \frac{1}{2} \text{tr} \left[\left\{ \left(\hat{\Sigma}_0 - \hat{\Sigma}(\delta) \right) \hat{\Sigma}_0^{-1} \right\}^2 \right].$$

They are distances between $\hat{\Sigma}(\delta)$ and $\hat{\Sigma}_0$ called a maximum likelihood fitting function and a generalized least square fitting function, respectively (e.g., Bollen [4], Chap. 4). Using $g(\mathbf{A})$, the GC_p and EGCV criteria, and the GIC and the AIC_C under normality are given by

$$GC_p(\delta) = nb_{\text{GLH}}(\mathbf{B}_{\delta}^*) + nbp + \alpha \text{df}(\delta),$$

$$\text{EGCV}(\delta) = \frac{bg_{\text{LH}}(\mathbf{B}_{\delta}^*) + bp}{\{1 - \text{df}(\delta)/np\}^\alpha},$$

$$\text{GIC}(\delta) = ng_{\text{LR}}(\mathbf{B}_{\delta}^*) + np \log b + \alpha \text{df}(\delta),$$

$$\text{AIC}_C(\delta) = ng_{\text{LR}}(\mathbf{B}_{\delta}^*) + np \log b + \frac{np\{n + \text{df}(\delta)\}}{n - p - 1 - \text{df}(\delta)}.$$

Using the GIC, it is also possible to adjust the strength of the penalty for model complexity, and for example, the GIC coincides with the AIC when $\alpha = 2$, the HQC (Hannan & Quinn [9]) when $\alpha = 2 \log \log n$, and the BIC (Schwarz [18]) when $\alpha = \log n$. For the GIC and AIC_C , the bivariate function $f(r, u)$ is given by

$$f(r, u) = \begin{cases} f_{\text{GIC}}(r, u) = n(r + p \log b) + \alpha u & (\text{GIC}) \\ f_{\text{AIC}_C}(r, u) = n(r + p \log b) + \frac{np(n + u)}{n - p - 1 - u} & (\text{AIC}_C) \end{cases}.$$

The following subsections describe two algorithms to minimize the MSC in (13).

4.1. Minimizing MSC via Iterative Method. This subsection describes an algorithm for solving the MSC minimization method via an iterative method with an iterative function. That is, we derive the iterative function. Notice that

$$\mathbf{B}_{\delta}^* = \sum_{j=1}^k \mathbf{z}_j^* \mathbf{z}_j^{*'} \delta_j^2, \quad \mathbf{z}_j^* = \mathbf{W}^{-1/2} \mathbf{z}_j.$$

Therefore, the following partial derivatives can be obtained:

$$\frac{\partial}{\partial \delta_j} \mathbf{B}_{\delta}^* = 2 \mathbf{z}_j^* \mathbf{z}_j^{*'} \delta_j, \quad \frac{\partial}{\partial \delta_j} \text{df}(\delta) = -p.$$

We express the (i, ℓ) element of a matrix \mathbf{A} as $a_{i\ell} = (\mathbf{A})_{i\ell}$ and define

$$\dot{g}_{i\ell}(\mathbf{B}) = \frac{\partial}{\partial a_{i\ell}} g(\mathbf{A}) \Big|_{\mathbf{A}=\mathbf{B}}, \quad \dot{\mathbf{G}}(\mathbf{B}) = \frac{\partial}{\partial \mathbf{A}} g(\mathbf{A}) \Big|_{\mathbf{A}=\mathbf{B}}. \quad (14)$$

\mathbf{B}_δ^* is a symmetric matrix, thus we have

$$\frac{\partial}{\partial \delta_j} g(\mathbf{B}_\delta^*) = \sum_{i=1}^p \sum_{\ell=i}^p \frac{\partial}{\partial \delta_j} (\mathbf{B}_\delta^*)_{i\ell} \cdot \dot{g}_{i\ell}(\mathbf{B}_\delta^*) = 2\mathbf{z}_j^{*\prime} \dot{\mathbf{G}}(\mathbf{B}_\delta^*) \mathbf{z}_j^* \delta_j.$$

Hence, a partial derivative of the MSC is given by

$$\begin{aligned} \frac{\partial}{\partial \delta_j} \text{MSC}(\boldsymbol{\delta} \mid g) &= 2\mathbf{z}_j^{*\prime} \dot{\mathbf{G}}(\mathbf{B}_\delta^*) \mathbf{z}_j^* \dot{f}_r(g(\mathbf{B}_\delta^*), \text{df}(\boldsymbol{\delta})) \delta_j \\ &\quad - p \dot{f}_u(g(\mathbf{B}_\delta^*), \text{df}(\boldsymbol{\delta})), \end{aligned}$$

where

$$\dot{f}_r(x, y) = \left. \frac{\partial}{\partial r} f(r, u) \right|_{(r,u)=(x,y)}, \quad \dot{f}_u(x, y) = \left. \frac{\partial}{\partial u} f(r, u) \right|_{(r,u)=(x,y)}.$$

By solving $\partial \text{MSC}(\boldsymbol{\delta} \mid g) / \partial \boldsymbol{\delta} = \mathbf{0}_k$, we can obtain the following iterative method:

$$\begin{aligned} \boldsymbol{\delta}^{(i+1)} &= \boldsymbol{\zeta}(\boldsymbol{\delta}^{(i)}) = \left(\zeta_1(\boldsymbol{\delta}^{(i)}), \dots, \zeta_k(\boldsymbol{\delta}^{(i)}) \right)' \quad (i = 0, 1, \dots), \\ \zeta_j(\boldsymbol{\delta}) &= 1 - \text{soft} \left(1, \tau(\boldsymbol{\delta}) / \mathbf{z}_j^{*\prime} \dot{\mathbf{G}}(\mathbf{B}_\delta^*) \mathbf{z}_j^* \right), \end{aligned} \quad (15)$$

where (i) is the iteration number, $\boldsymbol{\delta}^{(0)}$ is a given initial vector, and $\tau(\boldsymbol{\delta})$ is given by

$$\tau(\boldsymbol{\delta}) = \frac{p \dot{f}_u(g(\mathbf{B}_\delta^*), \text{df}(\boldsymbol{\delta}))}{2 \dot{f}_r(g(\mathbf{B}_\delta^*), \text{df}(\boldsymbol{\delta}))} > 0.$$

By repeating the update of $\boldsymbol{\delta}^{(i)}$ with the iterative function $\boldsymbol{\zeta}$, we can obtain the optimal $\boldsymbol{\delta}$. This iterative method has the following property (the proof is given in Appendix A.4).

PROPOSITION 5. *For a k -dimensional vector $\boldsymbol{\epsilon}$ wherein all elements are nonnegative, suppose that*

$$\tau(\boldsymbol{\delta}) \leq \tau(\boldsymbol{\delta} + \boldsymbol{\epsilon}), \quad \mathbf{z}_j^{*\prime} \dot{\mathbf{G}}(\mathbf{B}_\delta^*) \mathbf{z}_j^* \geq \mathbf{z}_j^{*\prime} \dot{\mathbf{G}}(\mathbf{B}_{\delta+\boldsymbol{\epsilon}}^*) \mathbf{z}_j^*. \quad (16)$$

Then, the iterative method with iterative function (15) converges if $\forall j \in \{1, \dots, k\}$, $\delta_j^{(1)} \geq \delta_j^{(0)}$ or $\forall j \in \{1, \dots, k\}$, $\delta_j^{(1)} \leq \delta_j^{(0)}$.

From this proposition, when assumption (16) holds, the iterative method with iterative function (15) converges if the initial vector is $\mathbf{0}_k$ or $\mathbf{1}_k$.

4.1.1. LR-distance. For the MSC based on the LR-distance, the following equation holds:

$$\frac{\partial}{\partial \mathbf{A}} g_{\text{LR}}(\mathbf{A}) = \frac{\partial}{\partial \mathbf{A}} \log |\mathbf{I}_p + \mathbf{A}| = (\mathbf{I}_p + \mathbf{A})^{-1}.$$

Therefore, we have

$$\mathbf{W}^{-1/2} \dot{\mathbf{G}}(\mathbf{B}_\delta^*) \mathbf{W}^{-1/2} = (\mathbf{W} + \mathbf{B}_\delta)^{-1} = \frac{1}{n} \hat{\boldsymbol{\Sigma}}(\boldsymbol{\delta})^{-1}.$$

Hence, the iterative function for solving the MSC minimization method based on the LR-distance is given by

$$\zeta_j(\boldsymbol{\delta}) = 1 - \text{soft} \left(1, n\tau(\boldsymbol{\delta})/z'_j \hat{\boldsymbol{\Sigma}}(\boldsymbol{\delta})^{-1} z_j \right). \quad (17)$$

Furthermore, from Lemma 1, for $\boldsymbol{\epsilon}$ in Proposition 5 and for any p -dimensional vector \mathbf{a} , the following equation holds:

$$\mathbf{a}' \hat{\boldsymbol{\Sigma}}(\boldsymbol{\delta}) \mathbf{a} \leq \mathbf{a}' \hat{\boldsymbol{\Sigma}}(\boldsymbol{\delta} + \boldsymbol{\epsilon}) \mathbf{a} \iff \mathbf{a}' \hat{\boldsymbol{\Sigma}}(\boldsymbol{\delta})^{-1} \mathbf{a} \geq \mathbf{a}' \hat{\boldsymbol{\Sigma}}(\boldsymbol{\delta} + \boldsymbol{\epsilon})^{-1} \mathbf{a}.$$

Let $\hat{\boldsymbol{\delta}}^{\text{LR}}$ be a solution obtained by the iterative method with iterative function (17). Then,

$$\hat{\boldsymbol{\delta}}^{\text{LR}} = \zeta(\hat{\boldsymbol{\delta}}^{\text{LR}}).$$

The ridge parameters optimized by the MSC minimization method based on the LR-distance are given by

$$\hat{\delta}_j^{\text{LR}} = 1 - \text{soft} \left(1, n\tau(\hat{\boldsymbol{\delta}}^{\text{LR}})/z'_j \hat{\boldsymbol{\Sigma}}(\hat{\boldsymbol{\delta}}^{\text{LR}})^{-1} z_j \right) \quad (j = 1, \dots, k).$$

On the other hand, the ridge parameters optimized by the MSC minimization method based on the LH-distance are given by the following form:

$$\hat{\delta}_j^{\text{LH}} = 1 - \text{soft} \left(1, \hat{h}/z'_j \mathbf{S}^{-1} z_j \right) \quad (j = 1, \dots, k).$$

The $\hat{\delta}_j^{\text{LH}}$ includes \mathbf{S}^{-1} and \mathbf{S} is an estimator of the covariance matrix for the full model. Thus, $\hat{\delta}_j^{\text{LH}}$ has a disadvantage because \mathbf{S}^{-1} is unstable when k is large. Whereas, $\hat{\delta}_j^{\text{LR}}$ does not include \mathbf{S}^{-1} , but rather $\hat{\boldsymbol{\Sigma}}(\hat{\boldsymbol{\delta}}^{\text{LR}})^{-1}$ and $\hat{\boldsymbol{\Sigma}}(\hat{\boldsymbol{\delta}}^{\text{LR}})$ is an estimator of the covariance matrix adjusted by $\hat{\boldsymbol{\delta}}^{\text{LR}}$. Thus, $\hat{\delta}_j^{\text{LR}}$ has an advantage because $\hat{\boldsymbol{\Sigma}}(\hat{\boldsymbol{\delta}}^{\text{LR}})^{-1}$ is stable even when k is large.

Example 1. We derive an iterative function for solving the GIC minimization method. From $f(r, u) = f_{\text{GIC}}(r, u)$, we have

$$\dot{f}_r(r, u) = n, \quad \dot{f}_u(r, u) = \alpha,$$

and therefore, $\tau(\boldsymbol{\delta}) = \alpha p / 2n$. Hence, the iterative function for the GIC minimization method is given by

$$\zeta_j(\boldsymbol{\delta}) = 1 - \text{soft} \left(1, \alpha p / 2z'_j \hat{\boldsymbol{\Sigma}}(\boldsymbol{\delta})^{-1} z_j \right) \quad (j = 1, \dots, k). \quad (18)$$

Moreover, since $\tau(\boldsymbol{\delta})$ does not depend on $\boldsymbol{\delta}$, from Proposition 5, the iterative method for solving the GIC minimization method converges under an appropriate initial vector.

Example 2. We derive an iterative function for solving the AIC_C minimization method. From $f(r, u) = f_{\text{AIC}_C}(r, u)$, we have

$$\dot{f}_r(r, u) = n, \quad \dot{f}_u(r, u) = \frac{np(2n - p - 1)}{(n - p - 1 - u)^2},$$

and therefore, we have

$$\tau(\boldsymbol{\delta}) = \frac{p^2(2n - p - 1)}{2\{n - p - 1 - \text{df}(\boldsymbol{\delta})\}^2}.$$

Hence, the iterative function for the AIC_C minimization method is given by

$$\zeta_j(\boldsymbol{\delta}) = 1 - \text{soft} \left(1, \frac{np^2(2n - p - 1)}{2\{n - p - 1 - \text{df}(\boldsymbol{\delta})\}^2 \mathbf{z}'_j \hat{\boldsymbol{\Sigma}}(\boldsymbol{\delta})^{-1} \mathbf{z}_j} \right) \quad (j = 1, \dots, k).$$

Moreover, for $\boldsymbol{\epsilon}$ in Proposition 5, the following equation holds:

$$\text{df}(\boldsymbol{\delta}) \geq \text{df}(\boldsymbol{\delta} + \boldsymbol{\epsilon}).$$

Therefore

$$\tau(\boldsymbol{\delta}) \geq \tau(\boldsymbol{\delta} + \boldsymbol{\epsilon}),$$

and thus, the iterative method for solving the AIC_C minimization method does not satisfy Proposition 5.

4.1.2. BNP-distance. For the MSC based on the BNP-distance, the following equation holds:

$$\begin{aligned} \frac{\partial}{\partial \mathbf{A}} g_{\text{BNP}}(\mathbf{A}) &= \frac{\partial}{\partial \mathbf{A}} \text{tr} \{ \mathbf{A}(\mathbf{I}_p + \mathbf{A})^{-1} \} \\ &= -\frac{\partial}{\partial \mathbf{A}} \text{tr} \{ (\mathbf{I}_p + \mathbf{A})^{-1} \} = (\mathbf{I}_p + \mathbf{A})^{-2}. \end{aligned}$$

Therefore, we have

$$\mathbf{W}^{-1/2} \hat{\mathbf{G}}(\mathbf{B}_\delta^*) \mathbf{W}^{-1/2} = (\mathbf{W} + \mathbf{B}_\delta)^{-1} \mathbf{W} (\mathbf{W} + \mathbf{B}_\delta)^{-1} = \frac{1}{n} \hat{\boldsymbol{\Sigma}}(\boldsymbol{\delta})^{-1} \hat{\boldsymbol{\Sigma}}_0 \hat{\boldsymbol{\Sigma}}(\boldsymbol{\delta})^{-1}.$$

Hence, the iterative function for solving the MSC minimization method based on the BNP-distance is given by

$$\zeta_j(\boldsymbol{\delta}) = 1 - \text{soft} \left(1, \frac{n\tau(\boldsymbol{\delta})}{\mathbf{z}'_j \hat{\boldsymbol{\Sigma}}(\boldsymbol{\delta})^{-1} \hat{\boldsymbol{\Sigma}}_0 \hat{\boldsymbol{\Sigma}}(\boldsymbol{\delta})^{-1} \mathbf{z}_j} \right).$$

Accordingly, using the BNP-distance, the optimal ridge parameters are stable even when k is large.

Example. As an example of MSC based on the BNP-distance, we consider the following criterion:

$$\text{BNPC}(\boldsymbol{\delta}) = ng_{\text{BNP}}(\mathbf{B}_{\boldsymbol{\delta}}^*) + \alpha \text{df}(\boldsymbol{\delta}).$$

Then, since

$$\dot{f}_r(r, u) = n, \quad \dot{f}_u(r, u) = \alpha,$$

we have $\tau(\boldsymbol{\delta}) = \alpha p/2n$. Hence, the iterative function for solving the BNPC minimization method is given by

$$\zeta_j(\boldsymbol{\delta}) = 1 - \text{soft} \left(1, \frac{\alpha p}{2\mathbf{z}'_j \hat{\boldsymbol{\Sigma}}(\boldsymbol{\delta})^{-1} \hat{\boldsymbol{\Sigma}}_0 \hat{\boldsymbol{\Sigma}}(\boldsymbol{\delta})^{-1} \mathbf{z}_j} \right). \quad (19)$$

4.1.3. *ML-distance.* For the MSC based on the ML-distance, the following equation holds:

$$\begin{aligned} \frac{\partial}{\partial \mathbf{A}} g_{\text{ML}}(\mathbf{A}) &= \frac{\partial}{\partial \mathbf{A}} [\text{tr} \{(\mathbf{I}_p + \mathbf{A})^{-1}\} + \log |\mathbf{I}_p + \mathbf{A}|] \\ &= -(\mathbf{I}_p + \mathbf{A})^{-2} + (\mathbf{I}_p + \mathbf{A})^{-1}. \end{aligned}$$

Therefore, we have

$$\begin{aligned} \mathbf{W}^{-1/2} \dot{\mathbf{G}}(\mathbf{B}_{\boldsymbol{\delta}}^*) \mathbf{W}^{-1/2} &= (\mathbf{W} + \mathbf{B}_{\boldsymbol{\delta}})^{-1} - (\mathbf{W} + \mathbf{B}_{\boldsymbol{\delta}})^{-1} \mathbf{W} (\mathbf{W} + \mathbf{B}_{\boldsymbol{\delta}})^{-1} \\ &= \frac{1}{n} \hat{\boldsymbol{\Sigma}}(\boldsymbol{\delta})^{-1} - \frac{1}{n} \hat{\boldsymbol{\Sigma}}(\boldsymbol{\delta})^{-1} \hat{\boldsymbol{\Sigma}}_0 \hat{\boldsymbol{\Sigma}}(\boldsymbol{\delta})^{-1}. \end{aligned}$$

Hence, the iterative function for solving the MSC minimization method based on the ML-distance is given by

$$\zeta_j(\boldsymbol{\delta}) = 1 - \text{soft} \left(1, \frac{n\tau(\boldsymbol{\delta})}{\mathbf{z}'_j \{ \hat{\boldsymbol{\Sigma}}(\boldsymbol{\delta})^{-1} - \hat{\boldsymbol{\Sigma}}(\boldsymbol{\delta})^{-1} \hat{\boldsymbol{\Sigma}}_0 \hat{\boldsymbol{\Sigma}}(\boldsymbol{\delta})^{-1} \} \mathbf{z}_j} \right).$$

Accordingly, using the ML-distance, the optimal ridge parameters are stable even when k is large.

4.1.4. *GLS-distance.* For the MSC based on the GLS-distance, the following equation holds:

$$\frac{\partial}{\partial \mathbf{A}} g_{\text{GLS}}(\mathbf{A}) = \frac{1}{2} \cdot \frac{\partial}{\partial \mathbf{A}} \text{tr}(\mathbf{A}^2) = \mathbf{A}.$$

Therefore, we have

$$\mathbf{W}^{-1/2} \dot{\mathbf{G}}(\mathbf{B}_{\boldsymbol{\delta}}^*) \mathbf{W}^{-1/2} = \mathbf{W}^{-1} \mathbf{B}_{\boldsymbol{\delta}} \mathbf{W}^{-1} = \frac{1}{n} \hat{\boldsymbol{\Sigma}}_0^{-1} \{ \hat{\boldsymbol{\Sigma}}(\boldsymbol{\delta}) - \hat{\boldsymbol{\Sigma}}_0 \} \hat{\boldsymbol{\Sigma}}_0^{-1}.$$

Hence, the iterative function for solving the MSC minimization method based on the GLS-distance is given by

$$\zeta_j(\boldsymbol{\delta}) = 1 - \text{soft} \left(1, \frac{n\tau(\boldsymbol{\delta})}{\mathbf{z}'_j \hat{\boldsymbol{\Sigma}}_0^{-1} \{ \hat{\boldsymbol{\Sigma}}(\boldsymbol{\delta}) - \hat{\boldsymbol{\Sigma}}_0 \} \hat{\boldsymbol{\Sigma}}_0^{-1} \mathbf{z}_j} \right).$$

Since $\hat{\Sigma}_0$ is an estimator of the covariance matrix for the full model, the optimal ridge parameters are unstable when k is large.

4.2. Minimizing MSC via Coordinate Descent. In the previous subsection, we described an algorithm to minimize the MSC via the iterative method with an iterative function obtained by solving $\partial \text{MSC}(\boldsymbol{\delta} \mid g) / \partial \boldsymbol{\delta} = \mathbf{0}_k$. In this subsection, we update $\delta_1, \dots, \delta_k$ individually, not simultaneously. That is, we minimize the MSC via a coordinate descent algorithm.

4.2.1. LR-distance. We partition $\mathbf{W} + \mathbf{B}_\boldsymbol{\delta}$ and $\text{df}(\boldsymbol{\delta})$ into

$$\begin{aligned} \mathbf{W} + \mathbf{B}_\boldsymbol{\delta} &= \mathbf{W}_j + \mathbf{z}_j \mathbf{z}'_j \delta_j^2, & \mathbf{W}_j &= \mathbf{W} + \sum_{\ell \neq j}^k \mathbf{z}_\ell \mathbf{z}'_\ell \delta_\ell^2, \\ \text{df}(\boldsymbol{\delta}) &= q_{1,j} - p\delta_j, & q_{1,j} &= p \left\{ (1+k) - \sum_{\ell \neq j}^k \delta_\ell \right\}. \end{aligned}$$

Then, the following equations hold:

$$\begin{aligned} \mathbf{I}_p + \mathbf{B}_\boldsymbol{\delta}^* &= \mathbf{W}^{-1/2} (\mathbf{W} + \mathbf{B}_\boldsymbol{\delta}) \mathbf{W}^{-1/2} \\ &= \mathbf{W}^{-1/2} (\mathbf{W}_j + \mathbf{z}_j \mathbf{z}'_j \delta_j^2) \mathbf{W}^{-1/2}, \\ |\mathbf{W}_j + \mathbf{z}_j \mathbf{z}'_j \delta_j^2| &= |\mathbf{W}_j| (1 + \mathbf{z}'_j \mathbf{W}_j^{-1} \mathbf{z}_j \delta_j^2). \end{aligned} \tag{20}$$

Therefore, we have

$$\begin{aligned} g_{\text{LR}}(\mathbf{B}_\boldsymbol{\delta}^*) &= \log |\mathbf{I}_p + \mathbf{B}_\boldsymbol{\delta}^*| \\ &= \log (1 + \mathbf{z}'_j \mathbf{W}_j^{-1} \mathbf{z}_j \delta_j^2) + \log |\mathbf{W}_j \mathbf{W}^{-1}| \\ &= \log(1 + q_{2,j} \delta_j^2) + q_{3,j}, \end{aligned}$$

where $q_{2,j}$ and $q_{3,j}$ are constants which do not depend on δ_j given by

$$q_{2,j} = \mathbf{z}'_j \mathbf{W}_j^{-1} \mathbf{z}_j, \quad q_{3,j} = \log |\mathbf{W}_j \mathbf{W}^{-1}|.$$

Hence, the following partial derivative is obtained:

$$\frac{\partial}{\partial \delta_j} g_{\text{LR}}(\mathbf{B}_\boldsymbol{\delta}^*) = \frac{2q_{2,j} \delta_j}{1 + q_{2,j} \delta_j^2}.$$

Example 1. The partial derivative of the GIC is given by

$$\dot{f}_j(\delta_j) = \frac{\partial}{\partial \delta_j} \text{GIC}(\boldsymbol{\delta}) = \frac{1}{1 + q_{2,j} \delta_j^2} (-\alpha p q_{2,j} \delta_j^2 + 2n q_{2,j} \delta_j - \alpha p).$$

An update equation of the coordinate descent algorithm for solving the GIC minimization method is given by the following theorem (the proof is given in Appendix A.5).

THEOREM 3. Let $f_j(\delta)$ be a function for $\delta \in [0, 1]$ and suppose that the derivative of $f_j(\delta)$ is given by the following form:

$$\begin{aligned}\dot{f}_j(\delta) &= \frac{1}{\dot{f}_{j,1}(\delta)} \dot{f}_{j,2}(\delta) \quad (\dot{f}_{j,1}(\delta) > 0), \\ \dot{f}_{j,2}(\delta) &= -c_{j,2}\delta^2 + 2c_{j,1}\delta - c_{j,0} \quad (c_{j,0}, c_{j,1}, c_{j,2} > 0),\end{aligned}$$

and we define $\tilde{\delta}_j$ as

$$\tilde{\delta}_j = \frac{1 - \sqrt{1 - c_{j,2}c_{j,0}/c_{j,1}^2}}{c_{j,2}/c_{j,1}}.$$

Then, $\hat{\delta}_j = \arg \min_{\delta \in [0,1]} f_j(\delta)$ is given by

(1) Case of $1 - c_{j,2}c_{j,0}/c_{j,1}^2 \geq 0$:

$$\hat{\delta}_j = \begin{cases} \tilde{\delta}_j & \left(c_{j,2} > c_{j,1} \text{ or } (c_{j,2} \leq c_{j,1} \text{ and } \tilde{\delta}_j < 1) \right) \\ 1 & \left(c_{j,2} \leq c_{j,1} \text{ and } \tilde{\delta}_j \geq 1 \right) \end{cases}.$$

(2) Case of $1 - c_{j,2}c_{j,0}/c_{j,1}^2 < 0$:

$$\hat{\delta}_j = 1.$$

Example 2. The partial derivative of the AIC_C is given by

$$\begin{aligned}\dot{f}_j(\delta_j) &= \frac{\partial}{\partial \delta_j} \text{AIC}_C(\boldsymbol{\delta}) \\ &= \frac{1}{(1 + q_{2,j}\delta_j^2)(n - p - 1 - q_{1,j} + p\delta_j)^2/n} \dot{f}_{j,2}(\delta_j), \\ \dot{f}_{j,2}(\delta) &= 2p^2q_{2,j}\delta^3 + pq_{2,j}\{4(n - p - 1 - q_{1,j}) - p(2n - p - 1)\}\delta^2 \\ &\quad + 2q_{2,j}(n - p - 1 - q_{1,j})^2\delta - p^2(2n - p - 1).\end{aligned}$$

An update equation of the coordinate descent algorithm for solving the AIC_C minimization method is given by the following theorem (the proof is given in Appendix A.6).

THEOREM 4. Let $f_j(\delta)$ be a function for $\delta \in [0, 1]$ and suppose that the derivative of $f_j(\delta)$ is given by the following form:

$$\begin{aligned}\dot{f}_j(\delta) &= \frac{1}{\dot{f}_{j,1}(\delta)} \dot{f}_{j,2}(\delta) \quad (\dot{f}_{j,1}(\delta) > 0), \\ \dot{f}_{j,2}(\delta) &= c_{j,3}\delta^3 + c_{j,2}\delta^2 + c_{j,1}\delta - c_{j,0} \quad (c_{j,0} > 0),\end{aligned}$$

and let m ($0 \leq m \leq 3$) be the number of stationary points of $\dot{f}_{j,2}(\delta)$ which is included in $(0, 1)$ and $\tilde{\delta}_{j,1}, \dots, \tilde{\delta}_{j,m}$ ($m \geq 1$) be the stationary points satisfying $\tilde{\delta}_{j,1} < \dots < \tilde{\delta}_{j,m}$. Moreover, we define a set \mathcal{S}_j as

$$\mathcal{S}_j = \{1\} \ (m = 0); \ \{\tilde{\delta}_{j,1}\} \ (m = 1); \ \{\tilde{\delta}_{j,1}, 1\} \ (m = 2); \ \{\tilde{\delta}_{j,1}, \tilde{\delta}_{j,3}\} \ (m = 3).$$

Then, $\hat{\delta}_j = \arg \min_{\delta \in [0,1]} f_j(\delta)$ is given by

$$\hat{\delta}_j = \arg \min_{\delta \in \mathcal{S}_j} f_j(\delta).$$

4.2.2. *BNP-distance.* Equation (20) leads to

$$(\mathbf{I}_p + \mathbf{B}_\delta^*)^{-1} = \mathbf{W}^{1/2}(\mathbf{W}_j + \mathbf{z}_j \mathbf{z}'_j \delta_j^2)^{-1} \mathbf{W}^{1/2},$$

and the following holds:

$$(\mathbf{W}_j + \mathbf{z}_j \mathbf{z}'_j \delta_j^2)^{-1} = \mathbf{W}_j^{-1} - \frac{\mathbf{W}_j^{-1} \mathbf{z}_j \mathbf{z}'_j \mathbf{W}_j^{-1} \delta_j^2}{1 + \mathbf{z}'_j \mathbf{W}_j^{-1} \mathbf{z}_j \delta_j^2} = \mathbf{W}_j^{-1} - \frac{\mathbf{W}_j^{-1} \mathbf{z}_j \mathbf{z}'_j \mathbf{W}_j^{-1} \delta_j^2}{1 + q_{2,j} \delta_j^2}.$$

Therefore, we have

$$\begin{aligned} g_{\text{BNP}}(\mathbf{B}_\delta^*) &= p - \text{tr} \{ (\mathbf{I}_p + \mathbf{B}_\delta^*)^{-1} \} \\ &= p - \text{tr} \left\{ \left(\mathbf{W}_j^{-1} - \frac{\mathbf{W}_j^{-1} \mathbf{z}_j \mathbf{z}'_j \mathbf{W}_j^{-1} \delta_j^2}{1 + q_{2,j} \delta_j^2} \right) \mathbf{W} \right\} \\ &= \frac{q_{4,j} \delta_j^2}{1 + q_{2,j} \delta_j^2} + q_{5,j}, \end{aligned}$$

where $q_{4,j}$ and $q_{5,j}$ are constants which do not depend on δ_j given by

$$q_{4,j} = \mathbf{z}'_j \mathbf{W}_j^{-1} \mathbf{W} \mathbf{W}_j^{-1} \mathbf{z}_j, \quad q_{5,j} = p - \text{tr}(\mathbf{W}_j^{-1} \mathbf{W}).$$

Hence, the following partial derivative is obtained:

$$\frac{\partial}{\partial \delta_j} g_{\text{BNP}}(\mathbf{B}_\delta) = \frac{2q_{4,j} \delta_j}{(1 + q_{2,j} \delta_j^2)^2}.$$

Example. The partial derivative of the BNPC is given by

$$\begin{aligned} \dot{f}_j(\delta_j) &= \frac{\partial}{\partial \delta_j} \text{BNPC}(\delta) \\ &= \frac{1}{(1 + q_{2,j} \delta_j^2)^2} (-\alpha p q_{2,j}^2 \delta_j^4 - 2\alpha p q_{2,j} \delta_j^2 + 2n q_{4,j} \delta_j - \alpha p). \end{aligned}$$

An update equation of the coordinate descent algorithm for solving the BNPC minimization method is given by the following theorem obtained which is similar to Theorem 4.

THEOREM 5. *Let $f_j(\delta)$ be a function for $\delta \in [0, 1]$ and suppose that the derivative of $f_j(\delta)$ is given by the following form:*

$$\begin{aligned} \dot{f}_j(\delta) &= \frac{1}{\dot{f}_{j,1}(\delta)} \dot{f}_{j,2}(\delta) \quad (\dot{f}_{j,1}(\delta) > 0), \\ \dot{f}_{j,2}(\delta) &= c_{j,4} \delta^4 + c_{j,3} \delta^3 + c_{j,2} \delta^2 + c_{j,1} \delta - c_{j,0} \quad (c_{j,0} > 0), \end{aligned}$$

and let m ($0 \leq m \leq 4$) be the number of stationary points of $f_{j,2}(\delta)$ which is included in $(0, 1)$ and $\tilde{\delta}_{j,1}, \dots, \tilde{\delta}_{j,m}$ ($m \geq 1$) be the stationary points satisfying $\tilde{\delta}_{j,1} < \dots < \tilde{\delta}_{j,m}$. Moreover, we define a set \mathcal{S}_j as

$$\mathcal{S}_j = \begin{cases} \{1\} & (m = 0); & \{\tilde{\delta}_{j,1}\} & (m = 1) \\ \{\tilde{\delta}_{j,1}, 1\} & (m = 2); & \{\tilde{\delta}_{j,1}, \tilde{\delta}_{j,3}\} & (m = 3) \\ \{\tilde{\delta}_{j,1}, \tilde{\delta}_{j,3}, 1\} & (m = 4) \end{cases}.$$

Then, $\hat{\delta}_j = \arg \min_{\delta \in [0,1]} f_j(\delta)$ is given by

$$\hat{\delta}_j = \arg \min_{\delta \in \mathcal{S}_j} f_j(\delta).$$

4.2.3. ML-distance. Notice that

$$g_{\text{ML}}(\mathbf{A}) = g_{\text{LR}}(\mathbf{A}) - g_{\text{BNP}}(\mathbf{A}).$$

Hence, we have

$$g_{\text{ML}}(\mathbf{B}_{\delta}^*) = \log(1 + q_{2,j}\delta_j^2) - \frac{q_{4,j}\delta_j^2}{1 + q_{2,j}\delta_j^2} + q_{3,j} - q_{5,j},$$

and the following partial derivative is obtained:

$$\frac{\partial}{\partial \delta_j} g_{\text{ML}}(\mathbf{B}_{\delta}^*) = \frac{2q_{2,j}\delta}{1 + q_{2,j}\delta^2} - \frac{2q_{4,j}\delta}{(1 + q_{2,j}\delta^2)^2}.$$

4.2.4. GLS-distance. We have

$$\mathbf{B}_{\delta} \mathbf{W}^{-1} = \mathbf{z}_j \mathbf{z}'_j \mathbf{W}^{-1} \delta_j^2 + \mathbf{W}_j \mathbf{W}^{-1} - \mathbf{I}_p,$$

and therefore

$$g_{\text{GLS}}(\mathbf{B}_{\delta}^*) = \frac{1}{2} \text{tr} \{ (\mathbf{B}_{\delta} \mathbf{W}^{-1})^2 \} = \frac{1}{2} (q_{6,j}\delta_j^4 + 2q_{7,j}\delta_j^2 + q_{8,j}),$$

where $q_{\ell,j}$ ($\ell = 6, 7, 8$) are constants which do not depend on δ_j given by

$$q_{6,j} = (\mathbf{z}'_j \mathbf{W}^{-1} \mathbf{z}_j)^2, \quad q_{7,j} = \mathbf{z}'_j \mathbf{W}^{-1} (\mathbf{W}_j - \mathbf{W}) \mathbf{W}^{-1} \mathbf{z}_j, \\ q_{8,j} = \text{tr} \{ (\mathbf{W}_j \mathbf{W}^{-1} - \mathbf{I}_p)^2 \}.$$

Hence, the following partial derivative is obtained:

$$\frac{\partial}{\partial \delta_j} g_{\text{GLS}}(\mathbf{B}_{\delta}^*) = 2q_{6,j}\delta_j^3 + 2q_{7,j}\delta_j.$$

5. Plug-in Iteration

In the previous section, we described the minimization of MSC extended to general distance. For MSC based on the LH-distance, the class of the optimal ridge parameters is obtained and since the minimizer is given as closed form and is unique, or minimizer candidates are given as closed forms and finite points, MSC can be minimized quickly. In contrast, since the optimal ridge parameters include the inverse of the estimator of the covariance matrix for the full model, those parameters are unstable when k is large. On the other hand, for MSC

based on general distance, in particular the LR-distance, the BNP-distance, and the ML-distance, since the estimator of the covariance matrix which is included in the optimal ridge parameters is an adjusted estimator, the optimal ridge parameters are stable even when k is large. In contrast, such MSC cannot be minimized quickly. As above, MSC based on the LH-distance and MSC based on another distance have contrasting properties. We propose a new approach, called the Plug-in Iteration Method (PIM) which is a hybrid method drawing on the merits of the various MSCs. The PIM optimizes ridge parameters by repeating the following procedure: first, the ridge parameters are optimized by the MSC minimization method based on the LH-distance; next, the ridge parameters are optimized again by using the ridge parameters optimized in the previous step.

The ridge parameters optimized by the MSC minimization method based on the LH-distance include \mathbf{S} , and this derives from the fact that the original distance $\text{tr}\{\hat{\Sigma}(\boldsymbol{\delta})\mathbf{S}^{-1}\}$ includes \mathbf{S} . Although the MSC was hitherto defined by using $\text{tr}(\mathbf{B}_{\boldsymbol{\delta}}^*)$ obtained from the original distance, we now redefine it using the original distance. For any $p \times p$ positive definite matrix \mathbf{A} , we define

$$r_+(\mathbf{A}) = \text{tr}\left(\hat{\Sigma}_0\mathbf{A}^{-1}\right) + \frac{1}{n}\text{tr}\left(\mathbf{Z}\mathbf{A}^{-1}\mathbf{Z}'\right),$$

and let f^\dagger be a bivariate function defined by the following class.

DEFINITION 4 (Class of the bivariate function f^\dagger). The f^\dagger satisfies the following conditions:

- (A1') For any $(r, u) \in (0, r_+(\mathbf{A})) \times [p, np)$, $f^\dagger(r, u)$ is continuous.
- (A2') For any $(r, u) \in (0, r_+(\mathbf{A})) \times [p, np)$, $f^\dagger(r, u)$ is positive.
- (A3') For any $(r, u) \in (0, r_+(\mathbf{A})) \times [p, np)$, $f^\dagger(r, u)$ is first order partially differentiable and its partial derivatives are positive.

Using the bivariate function f^\dagger , we redefine the MSC based on the LH-distance as

$$\text{MSC}^\dagger(\boldsymbol{\delta} \mid \mathbf{A}) = f^\dagger\left(\text{tr}\left\{\hat{\Sigma}(\boldsymbol{\delta})\mathbf{A}^{-1}\right\}, \text{df}(\boldsymbol{\delta})\right). \quad (21)$$

This MSC covers a wider class than the MSC in (10) and is equal to the MSC in (10) when $\mathbf{A} = \mathbf{S}$. For the GC_p and EGCV criteria, f^\dagger is given by

$$f^\dagger(r, u) = \begin{cases} nr + \alpha u & (GC_p \text{ criterion}) \\ r/(1 - u/np)^\alpha & (EGCV \text{ criterion}) \end{cases}.$$

Similar to Theorem 1, the optimal $\boldsymbol{\delta}$ minimizing the MSC in (21) is given by the following corollary.

COROLLARY 1. We define a function $\phi(h \mid \mathbf{A})$ ($h \in \mathbb{R}_+ \setminus \{0\}$) as

$$\phi(h \mid \mathbf{A}) = \text{MSC}^\dagger(\hat{\boldsymbol{\delta}}(h \mid \mathbf{A}) \mid \mathbf{A}),$$

and suppose that $\exists \nu > 0$ s.t. $\phi(\nu | \mathbf{A}) < \lim_{h \rightarrow 0} \phi(h | \mathbf{A})$, where $\hat{\boldsymbol{\delta}}(h | \mathbf{A}) = (\hat{\delta}_1(h | \mathbf{A}), \dots, \hat{\delta}_k(h | \mathbf{A}))'$ is a class of ridge parameters given by

$$\hat{\delta}_j(h | \mathbf{A}) = 1 - \text{soft} \left(1, h / \mathbf{z}'_j \mathbf{A}^{-1} \mathbf{z}_j \right).$$

Then, we have the following:

- (1) The optimal ridge parameters based on minimizing $\text{MSC}^\dagger(\boldsymbol{\delta} | \mathbf{A})$ are given by $\hat{\boldsymbol{\delta}}(\hat{h}_{\mathbf{A}} | \mathbf{A})$ and $\hat{h}_{\mathbf{A}}$ is given by

$$\hat{h}_{\mathbf{A}} = \arg \min_{h \in \mathbb{R}_+ \setminus \{0\}} \phi(h | \mathbf{A}).$$

- (2) The $\phi(h | \mathbf{A})$ has the following properties:

(P1) For all $h \in \mathbb{R}_+ \setminus \{0\}$, $\phi(h | \mathbf{A})$ is continuous.

(P2) For all $h \geq t_k$, $\phi(h | \mathbf{A}) = f^\dagger(r_+(\mathbf{A}), p)$.

(P3) The $\phi(h | \mathbf{A})$ can be expressed as the following piecewise function:

$$\begin{aligned} \phi(h | \mathbf{A}) &= \phi_a(h | \mathbf{A}) \quad (h \in R_a; a = 0, 1, \dots, k) \\ &= f^\dagger \left(\text{tr}(\hat{\boldsymbol{\Sigma}}_0 \mathbf{A}^{-1}) + (c_{1,a} + c_{2,a} h^2) / n, p(1 + k - a - c_{2,a} h) \right), \end{aligned}$$

where R_a , $c_{1,a}$ and $c_{2,a}$ are range and nonnegative constants similar to (11) and Proposition 1, respectively. However, t_j ($j = 1, \dots, k$) is the j th order statistic of $\mathbf{z}'_j \mathbf{A}^{-1} \mathbf{z}_j$ ($j = 1, \dots, k$).

Corollary 1 is an extension of Theorem 1 and Proposition 1 and they are equivalent when $\mathbf{A} = \mathbf{S}$. Furthermore, $\hat{h}_{\mathbf{A}}$ can be obtained by applying Theorem 2.

Using Corollary 1, we describe the PIM algorithm. Let $\mathbf{S}^{(0)} = \mathbf{S}$ and we define $\hat{\boldsymbol{\delta}}^{(0)}(h) = (\hat{\delta}_1^{(0)}(h), \dots, \hat{\delta}_k^{(0)}(h))'$ as $\hat{\boldsymbol{\delta}}^{(0)}(h) = \hat{\boldsymbol{\delta}}(h | \mathbf{S}^{(0)})$ and define the optimal ridge parameters based on minimizing $\text{MSC}^\dagger(\boldsymbol{\delta} | \mathbf{S}^{(0)})$ as

$$\begin{aligned} \hat{\boldsymbol{\delta}}^{(0)} &= \left(\hat{\delta}_1^{(0)}, \dots, \hat{\delta}_k^{(0)} \right)', \quad \hat{\delta}_j^{(0)} = \hat{\delta}_j^{(0)}(\hat{h}^{(0)}), \\ \hat{h}^{(0)} &= \arg \min_{h \in \mathbb{R}_+ \setminus \{0\}} \phi^{(0)}(h), \quad \phi^{(0)}(h) = \text{MSC}^\dagger(\hat{\boldsymbol{\delta}}^{(0)}(h) | \mathbf{S}^{(0)}). \end{aligned}$$

Therefore $\hat{\delta}_j^{(0)}$ is given by

$$\hat{\delta}_j^{(0)} = 1 - \text{soft} \left(1, \hat{h}^{(0)} / \mathbf{z}'_j \{\mathbf{S}^{(0)}\}^{-1} \mathbf{z}_j \right). \quad (22)$$

Furthermore, by substituting $\hat{\boldsymbol{\delta}}^{(0)}$, we define $\mathbf{S}^{(1)}$ as

$$\mathbf{S}^{(1)} = \mathbf{W}^{1/2} \dot{\mathbf{G}}(\mathbf{B}_{\hat{\boldsymbol{\delta}}^{(0)}}^*)^{-1} \mathbf{W}^{1/2},$$

where \mathbf{W} is given in Lemma 1 and $\dot{\mathbf{G}}(\cdot)$ is given by (14), and let $\hat{\boldsymbol{\delta}}^{(1)}(h)$ be a class of ridge parameters wherein the j th element ($j = 1, \dots, k$) is given by

$$\hat{\delta}_j^{(1)}(h) = 1 - \text{soft} \left(1, h / \mathbf{z}'_j \{\mathbf{S}^{(1)}\}^{-1} \mathbf{z}_j \right).$$

Then, we optimize the ridge parameters again as

$$\begin{aligned}\hat{\boldsymbol{\delta}}^{(1)} &= \left(\hat{\delta}_1^{(1)}, \dots, \hat{\delta}_k^{(1)} \right)', \quad \hat{\delta}_j^{(1)} = \hat{\delta}_j^{(1)}(\hat{h}^{(1)}), \\ \hat{h}^{(1)} &= \arg \min_{h \in \mathbb{R}_+ \setminus \{0\}} \phi^{(1)}(h), \quad \phi^{(1)}(h) = \text{MSC}^\dagger(\hat{\boldsymbol{\delta}}^{(1)}(h) \mid \mathbf{S}^{(1)}).\end{aligned}$$

The $\hat{h}^{(1)}$ can be obtained quickly by applying Theorem 2. Since the optimal ridge parameter $\hat{\boldsymbol{\delta}}^{(0)}$ includes \mathbf{S} , it is unstable when k is large. However, since $\mathbf{S}^{(1)}$ is adjusted by substituting $\hat{\boldsymbol{\delta}}^{(0)}$, $\hat{\boldsymbol{\delta}}^{(1)}$ is stable even when k is large. The PIM algorithm is summarized as follows.

PIM Algorithm

Step 1. Let the initial vector $\hat{\boldsymbol{\delta}}^{(0)}$ be the ridge parameters optimized by the MSC minimization method based on the LH-distance and $i \leftarrow 0$.

Step 2. Define $\mathbf{S}^{(i+1)}$ and $\phi^{(i+1)}(h)$ as

$$\begin{aligned}\mathbf{S}^{(i+1)} &= \mathbf{W}^{1/2} \dot{\mathbf{G}}(\mathbf{B}_{\hat{\boldsymbol{\delta}}^{(i)}}^*)^{-1} \mathbf{W}^{1/2}, \\ \phi^{(i+1)}(h) &= \text{MSC}^\dagger \left(\hat{\boldsymbol{\delta}}^{(i+1)}(h) \mid \mathbf{S}^{(i+1)} \right),\end{aligned}$$

where the class of ridge parameters is given by

$$\begin{aligned}\hat{\boldsymbol{\delta}}^{(i+1)}(h) &= \left(\hat{\delta}_1^{(i+1)}(h), \dots, \hat{\delta}_k^{(i+1)}(h) \right)', \\ \hat{\delta}_j^{(i+1)}(h) &= 1 - \text{soft} \left(1, h / \mathbf{z}_j' \{ \mathbf{S}^{(i+1)} \}^{-1} \mathbf{z}_j \right).\end{aligned}$$

Step 3. By using Theorem 2, update the ridge parameters as

$$\hat{\boldsymbol{\delta}}^{(i+1)} = \hat{\boldsymbol{\delta}}^{(i+1)}(\hat{h}^{(i+1)}), \quad \hat{h}^{(i+1)} = \arg \min_{h \in \mathbb{R}_+ \setminus \{0\}} \phi^{(i+1)}(h).$$

Step 4. If $\hat{\boldsymbol{\delta}}^{(i+1)}$ converges, the algorithm is complete. If not, let $i \leftarrow i + 1$ and return to Step 2.

Since the MSC minimized at each iteration is based on the LH-distance, the minimization is fast. Furthermore, an estimator of the covariance matrix which is included in $\hat{\boldsymbol{\delta}}^{(i)}$ is stable by substituting the ridge parameters optimized in the previous step. Thus, the PIM is a hybrid method which leverages the merits of the various MSCs.

The PIM algorithm is similar to the iterative method. In particular, when using the GC_p criterion, for all i ($= 0, 1, \dots$), we have

$$\hat{h}^{(i)} = \frac{\alpha p}{2}.$$

Therefore, the PIM is the following iterative method:

$$\hat{\delta}_j^{(i+1)} = 1 - \text{soft} \left(1, \alpha p / 2 \mathbf{z}_j^* \dot{\mathbf{G}}(\mathbf{B}_{\hat{\boldsymbol{\delta}}^{(i)}}^*) \mathbf{z}_j^* \right),$$

and this is equal to the iterative method wherein the initial vector is the ridge parameters optimized by the GC_p criterion minimization method, the iterative

function is equation (15), and $\tau(\boldsymbol{\delta}) = \alpha p/2$. That is, when using the GC_p criterion, the PIM with the GIC is equal to the GIC minimization method and the PIM with the BNPC is equal to the BNPC minimization method.

6. Relationship with Multivariate Adaptive Group Lasso Regression

In this section, we describe a relationship between the MGR and MAGL estimators after the regularization parameters are optimized by the MSC minimization method based on the LH-distance. The MAGL estimator cannot usually be obtained as closed form. However, it can be obtained as closed form under orthogonal explanatory variables. Although we use general \mathbf{X} until the previous section, this section deals with orthogonal explanatory variables. Furthermore, instead of using the transformed ridge parameters $\delta_1, \dots, \delta_k$, we approach this via the original ridge parameters $\theta_1, \dots, \theta_k$.

6.1. Estimators with Optimal Regularization Parameters under Orthogonality. The orthogonality of \mathbf{X} means $\mathbf{Q} = \mathbf{I}_k$ in (3). Therefore, the LS and the MGR estimators of $\boldsymbol{\Xi}$ in (2) and (4), respectively, are rewritten as

$$\begin{aligned}\hat{\boldsymbol{\Xi}} &= \left(\hat{\boldsymbol{\xi}}_1, \dots, \hat{\boldsymbol{\xi}}_k\right)' = \mathbf{D}^{-1/2} \mathbf{Z}, \quad \hat{\boldsymbol{\xi}}_j = \frac{1}{\sqrt{d_j}} z_j, \\ \hat{\boldsymbol{\Xi}}_{\theta}^{\text{R}} &= \left(\hat{\boldsymbol{\xi}}_{\theta_1,1}^{\text{R}}, \dots, \hat{\boldsymbol{\xi}}_{\theta_k,k}^{\text{R}}\right)' = \mathbf{D}^{1/2} (\mathbf{D} + \boldsymbol{\Theta})^{-1} \mathbf{Z}, \quad \hat{\boldsymbol{\xi}}_{\theta_j,j}^{\text{R}} = \frac{\sqrt{d_j}}{d_j + \theta_j} z_j,\end{aligned}\quad (23)$$

where $\mathbf{D} = \text{diag}(d_1, \dots, d_k)$ and $\mathbf{Z} = (z_1, \dots, z_k)'$ are the $k \times k$ diagonal matrix and the $k \times p$ matrix given by (3) and (6), respectively. The ridge parameters are optimized using the following MSC based on the LH-distance:

$$\text{MSC}_{\text{R}}(\boldsymbol{\theta} \mid \mathbf{A}) = f^{\dagger} \left(\text{tr}\{\hat{\boldsymbol{\Sigma}}_{\text{R}}(\boldsymbol{\theta}) \mathbf{A}^{-1}\}, \text{df}_{\text{R}}(\boldsymbol{\theta}) \right), \quad (24)$$

where $\hat{\boldsymbol{\Sigma}}_{\text{R}}(\boldsymbol{\theta})$ and $\text{df}_{\text{R}}(\boldsymbol{\theta})$ are given by transforming the parameter from $\boldsymbol{\delta}$ to $\boldsymbol{\theta}$ in $\hat{\boldsymbol{\Sigma}}(\boldsymbol{\delta})$ and $\text{df}(\boldsymbol{\delta})$, which are given by (8) and (9), respectively, as

$$\hat{\boldsymbol{\Sigma}}_{\text{R}}(\boldsymbol{\theta}) = \hat{\boldsymbol{\Sigma}}_0 + \frac{1}{n} \sum_{j=1}^k z_j z_j' \left(\frac{\theta_j}{d_j + \theta_j} \right)^2, \quad \text{df}_{\text{R}}(\boldsymbol{\theta}) = p(1+k) - p \sum_{j=1}^k \frac{\theta_j}{d_j + \theta_j}.$$

Thus, the MSC in (24) is the parameter-transformed version of the MSC in (21). Furthermore, since the transformation is a one-to-one correspondence, Corollary 1 gives the following class of ridge parameters optimized by minimizing $\text{MSC}_{\text{R}}(\boldsymbol{\theta} \mid \mathbf{A})$:

$$\begin{aligned}\hat{\boldsymbol{\theta}}(h \mid \mathbf{A}) &= \left(\hat{\theta}_1(h \mid \mathbf{A}), \dots, \hat{\theta}_k(h \mid \mathbf{A}) \right)', \\ \hat{\theta}_j(h \mid \mathbf{A}) &= \begin{cases} \frac{d_j h}{z_j' \mathbf{A}^{-1} z_j - h} & (h < z_j' \mathbf{A}^{-1} z_j) \\ \infty & (h \geq z_j' \mathbf{A}^{-1} z_j) \end{cases}.\end{aligned}$$

Notice that for all $x \in \mathbb{R}_+$,

$$\text{MSC}_R(\hat{\boldsymbol{\theta}}(x | \mathbf{A}) | \mathbf{A}) = \text{MSC}^\dagger(\hat{\boldsymbol{\delta}}(x | \mathbf{A}) | \mathbf{A}).$$

Then, from Corollary 1, the optimal ridge parameters based on minimizing the MSC in (24) are given by

$$\begin{aligned} \hat{\theta}_j &= \hat{\theta}_j(\hat{h}_{\mathbf{A}} | \mathbf{A}) \quad (j = 1, \dots, k), \\ \hat{h}_{\mathbf{A}} &= \arg \min_{h \in \mathbb{R}_+ \setminus \{0\}} \phi(h | \mathbf{A}), \quad \phi(h | \mathbf{A}) = \text{MSC}^\dagger(\hat{\boldsymbol{\delta}}(h | \mathbf{A}) | \mathbf{A}), \end{aligned} \quad (25)$$

and using these optimal ridge parameters, the optimal MGR estimator based on minimizing the MSC in (24) is given by

$$\hat{\boldsymbol{\xi}}_{\theta_j, j}^R = \frac{1}{\sqrt{d_j}} \text{soft} \left(1, \hat{h}_{\mathbf{A}} / \mathbf{z}'_j \mathbf{A}^{-1} \mathbf{z}_j \right) \mathbf{z}_j. \quad (26)$$

Since $\hat{\boldsymbol{\xi}}_{\theta_j, j}^R = \mathbf{0}_p$ when $\hat{h}_{\mathbf{A}} \geq \mathbf{z}'_j \mathbf{A}^{-1} \mathbf{z}_j$, we found that the non-sparse MGR estimator is sparse after the ridge parameters are optimized.

Next, we describe the MAGL estimator of $\boldsymbol{\Xi}$. Ohishi *et al.* [17] derived the AL estimator as closed form under orthogonality of \mathbf{X} . As a natural extension of this result, the MAGL estimator can be obtained as closed form. Let \mathbf{L}_λ be a $k \times k$ diagonal matrix of which the j th diagonal element is given by

$$\ell_{\lambda, j} = \frac{1}{d_j} \text{soft} \left(1, \lambda w_j / \sqrt{d_j} \|\mathbf{z}_j\| \right) \quad (j = 1, \dots, k),$$

where $\lambda \in \mathbb{R}_+$ is a regularization parameter called a tuning parameter, w_j is a weight, and $\|\cdot\|$ is L_2 norm of a vector. Then, the MAGL estimator of $\boldsymbol{\Xi}$ is given by

$$\begin{aligned} \hat{\boldsymbol{\Xi}}_\lambda^L &= \left(\hat{\boldsymbol{\xi}}_{\lambda, 1}^L, \dots, \hat{\boldsymbol{\xi}}_{\lambda, k}^L \right)' = \mathbf{L}_\lambda \mathbf{X}' \mathbf{Y} = \mathbf{L}_\lambda \mathbf{D}^{1/2} \mathbf{Z}, \\ \hat{\boldsymbol{\xi}}_{\lambda, j}^L &= \sqrt{d_j} \ell_{\lambda, j} \mathbf{z}_j = \frac{1}{\sqrt{d_j}} \text{soft} \left(1, \lambda w_j / \sqrt{d_j} \|\mathbf{z}_j\| \right) \mathbf{z}_j. \end{aligned} \quad (27)$$

Since $\mathbf{L}_\lambda = \mathbf{D}^{-1}$ when $\lambda = 0$, the MAGL estimator coincides with the LS estimator when $\lambda = 0$, and the MAGL estimator coincides with the AL estimator given in Ohishi *et al.* [17] when $p = 1$. The MAGL estimator is sparse in the sense that $\hat{\boldsymbol{\xi}}_{\lambda, j}^L = \mathbf{0}_p$ when $\lambda w_j \geq \sqrt{d_j} \|\mathbf{z}_j\|$. The $\hat{\boldsymbol{\Xi}}_\lambda^L$ in (27) denotes the minimizer of the following PRSS:

$$\text{tr} \{ (\mathbf{Y} - \mathbf{1}_n \boldsymbol{\mu}' - \mathbf{X} \boldsymbol{\Xi})' (\mathbf{Y} - \mathbf{1}_n \boldsymbol{\mu}' - \mathbf{X} \boldsymbol{\Xi}) \} + 2\lambda \sum_{j=1}^k w_j \|\boldsymbol{\xi}_j\|. \quad (28)$$

The MGR estimator in (23) depends on k regularization parameters. Whereas, the MAGL estimator in (27) depends on only one regularization parameter. Furthermore, although the MGR estimator is not sparse, the MAGL estimator is characterized by sparsity. Hence, it can be stated that the MGR and MAGL estimators have different properties.

The MAGL estimator in (27) gives a predictive matrix of \mathbf{Y} for the MAGL regression as follows:

$$\hat{\mathbf{Y}}_\lambda^L = \mathbf{1}_n \hat{\boldsymbol{\mu}}' + \mathbf{X} \hat{\boldsymbol{\Xi}}_\lambda^L = \mathbf{H}_\lambda^L \mathbf{Y}, \quad \mathbf{H}_\lambda^L = \mathbf{J}_n + \mathbf{X} \mathbf{L}_\lambda \mathbf{X}'.$$

Using $\hat{\mathbf{Y}}_\lambda^L$ and \mathbf{H}_λ^L , we define an estimator of $\boldsymbol{\Sigma}$ and a GDF as

$$\hat{\boldsymbol{\Sigma}}_L(\lambda) = \frac{(\mathbf{Y} - \hat{\mathbf{Y}}_\lambda^L)'(\mathbf{Y} - \hat{\mathbf{Y}}_\lambda^L)}{n} = \frac{\mathbf{Y}'(\mathbf{I}_n - \mathbf{J}_n - \mathbf{X} \mathbf{L}_\lambda \mathbf{X}')^2 \mathbf{Y}}{n},$$

$$\text{df}_L(\lambda) = p \text{tr}(\mathbf{H}_\lambda^L).$$

Similar to Ohishi *et al.* [17], we have the following lemma concerning $\hat{\boldsymbol{\Sigma}}_L(\lambda)$ and $\text{df}_L(\lambda)$.

LEMMA 5. *The $\hat{\boldsymbol{\Sigma}}_L(\lambda)$ and $\text{df}_L(\lambda)$ are expressed as*

$$\begin{aligned} \hat{\boldsymbol{\Sigma}}_L(\lambda) &= \hat{\boldsymbol{\Sigma}}_0 + \frac{1}{n} \mathbf{Z}'(\mathbf{I}_k - \mathbf{D} \mathbf{L}_\lambda)^2 \mathbf{Z} \\ &= \hat{\boldsymbol{\Sigma}}_0 + \frac{1}{n} \sum_{j=1}^k \left\{ 1 - \text{soft} \left(1, \lambda w_j / \sqrt{d_j} \| \mathbf{z}_j \| \right) \right\}^2 \mathbf{z}_j \mathbf{z}_j', \\ \text{df}_L(\lambda) &= p + p \sum_{j=1}^k \text{soft} \left(1, \lambda w_j / \sqrt{d_j} \| \mathbf{z}_j \| \right). \end{aligned}$$

Then, the MSC for optimizing the tuning parameter in the MAGL regression is given by

$$\text{MSC}_L(\lambda \mid \mathbf{A}) = f^\dagger \left(\text{tr}(\hat{\boldsymbol{\Sigma}}_L(\lambda) \mathbf{A}^{-1}), \text{df}_L(\lambda) \right), \quad (29)$$

and the tuning parameter optimized by the MSC minimization method is given by

$$\hat{\lambda}_{\mathbf{A}} = \arg \min_{\lambda \in \mathbb{R}_+} \text{MSC}_L(\lambda \mid \mathbf{A}).$$

Regarding the weight w_j , in general, an inverse of a norm of an estimator of $\boldsymbol{\xi}_j$ is used. When using the weight $w_j = 1/\|\hat{\boldsymbol{\xi}}_j\|$ based on the LS estimator, the optimal MAGL estimator based on minimizing the MSC in (29) is given by

$$\hat{\boldsymbol{\xi}}_{\hat{\lambda}_{\mathbf{A}}, j}^L = \frac{1}{\sqrt{d_j}} \text{soft} \left(1, \hat{\lambda}_{\mathbf{A}} / \| \mathbf{z}_j \|^2 \right) \mathbf{z}_j. \quad (30)$$

6.2. Equivalence between MGR and MAGL estimators. This subsection investigates a relationship between the MGR and MAGL estimators under the regularization parameters optimized by the MSC minimization method. Although the optimal MGR estimator in (26) and the optimal MAGL estimator in (30) have similar forms, the optimal MGR estimator does not include $\| \mathbf{z}_j \|^2$, but rather $\mathbf{z}_j \mathbf{A}^{-1} \mathbf{z}_j$ normalized by \mathbf{A} . We focus on the difference.

Let \mathbf{T} be an $n \times p$ matrix defined by $\mathbf{T} = \mathbf{Y} \mathbf{A}^{-1/2}$, \mathbf{U} and $\boldsymbol{\Gamma}$ be $k \times p$ matrices defined by $\mathbf{U} = (\mathbf{u}_1, \dots, \mathbf{u}_k)' = \mathbf{Z} \mathbf{A}^{-1/2} = \mathbf{P}'_1 \mathbf{T}$ and $\boldsymbol{\Gamma} = (\boldsymbol{\gamma}_1, \dots, \boldsymbol{\gamma}_k)' =$

$\Xi \mathbf{A}^{-1/2}$, respectively, and \mathbf{v} be a p -dimensional vector defined by $\mathbf{v} = \mathbf{A}^{-1/2} \boldsymbol{\mu}$. Then, we normalize the PRSS for the MGR regression as

$$\begin{aligned} & \text{tr} \{ (\mathbf{Y} - \mathbf{1}_n \boldsymbol{\mu}' - \mathbf{X} \Xi)' (\mathbf{Y} - \mathbf{1}_n \boldsymbol{\mu}' - \mathbf{X} \Xi) \mathbf{A}^{-1} + \Xi' \mathbf{Q} \Theta \mathbf{Q}' \Xi \mathbf{A}^{-1} \} \\ & = \text{tr} \{ (\mathbf{T} - \mathbf{1}_n \mathbf{v}' - \mathbf{X} \Gamma)' (\mathbf{T} - \mathbf{1}_n \mathbf{v}' - \mathbf{X} \Gamma) + \Gamma' \mathbf{Q} \Theta \mathbf{Q}' \Gamma \}. \end{aligned}$$

This normalized PRSS provides the MGR estimator of γ_j as

$$\hat{\gamma}_{\theta_j, j}^{\text{R}} = \frac{\sqrt{d_j}}{d_j + \theta_j} \mathbf{u}_j.$$

Therefore, the MGR normalized estimator of $\boldsymbol{\xi}_j$ is given by

$$\hat{\boldsymbol{\xi}}_{\theta_j, j}^{\text{R}\dagger} = \mathbf{A}^{1/2} \hat{\gamma}_{\theta_j, j}^{\text{R}} = \frac{\sqrt{d_j}}{d_j + \theta_j} \mathbf{z}_j,$$

and this is equal to the MGR estimator in (23). That is, the MGR estimator in (23) is a normalized estimator in spite of the fact that it is obtained from non-normalized PRSS in (5). Thus, the optimal MGR normalized estimator is given by (26). On the other hand, based on Xin *et al.* [23], we normalize the PRSS for the MAGL regression as

$$\begin{aligned} & \text{tr} \{ (\mathbf{Y} - \mathbf{1}_n \boldsymbol{\mu}' - \mathbf{X} \Xi)' (\mathbf{Y} - \mathbf{1}_n \boldsymbol{\mu}' - \mathbf{X} \Xi) \mathbf{A}^{-1} \} + 2\lambda \sum_{j=1}^k w_j \|\mathbf{A}^{-1/2} \boldsymbol{\xi}_j\| \\ & = \text{tr} \{ (\mathbf{T} - \mathbf{1}_n \mathbf{v}' - \mathbf{X} \Gamma)' (\mathbf{T} - \mathbf{1}_n \mathbf{v}' - \mathbf{X} \Gamma) \} + 2\lambda \sum_{j=1}^k w_j \|\boldsymbol{\gamma}_j\|. \end{aligned}$$

When using the general weight $w_j = 1/\|\hat{\boldsymbol{\gamma}}_j\|$ ($\hat{\boldsymbol{\gamma}}_j$ is the LS estimator of $\boldsymbol{\gamma}_j$), this normalized PRSS provides the MAGL estimator of $\boldsymbol{\gamma}_j$ as

$$\hat{\boldsymbol{\gamma}}_{\lambda, j}^{\text{L}} = \frac{1}{\sqrt{d_j}} \text{soft}(1, \lambda/\|\mathbf{u}_j\|^2) \mathbf{u}_j.$$

Therefore, the MAGL normalized estimator of $\boldsymbol{\xi}_j$ is given by

$$\hat{\boldsymbol{\xi}}_{\lambda, j}^{\text{L}\dagger} = \mathbf{A}^{1/2} \hat{\boldsymbol{\gamma}}_{\lambda, j}^{\text{L}} = \frac{1}{\sqrt{d_j}} \text{soft}(1, \lambda/\mathbf{z}_j' \mathbf{A}^{-1} \mathbf{z}_j) \mathbf{z}_j,$$

and this is different from the MAGL estimator in (27) obtained as the minimizer of the PRSS in (28) with the weight $w_j = 1/\|\hat{\boldsymbol{\xi}}_j\|$. Hence, the difference between the two optimal estimators (26) and (30) is whether the estimator is normalized or not. If $\hat{h}_{\mathbf{A}} = \hat{\lambda}_{\mathbf{A}}$, the two optimal normalized estimators are equivalent. The equivalence is given by the following theorem (the proof is given in Appendix A.7).

THEOREM 6. *Suppose that $w_j = 1/\|\hat{\boldsymbol{\gamma}}_j\|$ and let $\hat{\theta}_j$ ($j = 1, \dots, k$) and $\hat{\lambda}$ be the regularization parameters optimized by the MSC minimization method based*

on the LH-distance defined by

$$\hat{\theta}_j = \hat{\theta}_j(\hat{h}_{\mathbf{A}} | \mathbf{A}), \quad \hat{h}_{\mathbf{A}} = \arg \min_{h \in \mathbb{R}_+} \text{MSC}_{\text{R}}(\hat{\theta}(h | \mathbf{A}) | \mathbf{A}),$$

$$\hat{\lambda} = \hat{\lambda}_{\mathbf{A}} = \arg \min_{\lambda \in \mathbb{R}_+} \text{MSC}_{\text{L}}(\lambda | \mathbf{A}).$$

Then, the following equation holds:

$$\hat{\xi}_{\hat{\theta}_j, j}^{\text{R}\dagger} = \hat{\xi}_{\hat{\lambda}, j}^{\text{L}\dagger} \quad (j = 1, \dots, k).$$

7. Numerical Studies

In this section, we explore the performance of the MSC minimization methods for optimizing ridge parameters by evaluating prediction accuracies of predictive matrices via simulation. This simulation is executed using R (ver. 3.6.0) on a computer with a Windows 10 Pro operating system, Intel (R) Core i7-7700 processor, and 16 GB of RAM. Let $\mathbf{R}_k = \text{diag}(1, \dots, k)$ and let $\Omega_k(\rho)$ be a $k \times k$ matrix of which the (i, j) element is given by $\rho^{|i-j|}$. Then, the simulation data are generated from the following model:

$$\mathbf{Y} \sim N_{n \times p}(\mathbf{X}\Xi, \Sigma \otimes \mathbf{I}_n),$$

$$\mathbf{X} = (\mathbf{I}_n - \mathbf{J}_n)\mathbf{X}_0\Psi(0.99)^{1/2}, \quad \Sigma = \mathbf{R}_p^{1/2}\Omega_p(\rho_y)\mathbf{R}_p^{1/2},$$

where Ξ and \mathbf{X}_0 are $k \times p$ and $n \times k$ matrices wherein all the elements are identically and independently distributed according to $U(-1, 1)$ and $\Psi(\rho)$ is a correlation matrix of \mathbf{X} defined by $\Psi(\rho) = \mathbf{R}_k^{1/2}\Omega_k(\rho)\mathbf{R}_k^{1/2}$. Furthermore, $\rho = 0.99$ and thus this simulation is a highly correlated setting. Finally, Ξ and \mathbf{X}_0 are fixed throughout the simulation iterations.

Let $\hat{\mathbf{Y}}_{\hat{\delta}}$ be the predictive matrix of \mathbf{Y} obtained from the optimal MGR estimator based on minimizing the MSC and $\hat{\mathbf{Y}}$ be the predictive matrix of \mathbf{Y} obtained from the LS estimator, i.e., $\hat{\mathbf{Y}} = \hat{\mathbf{Y}}_{\mathbf{0}_k}$. Then, we evaluate the prediction accuracy of $\hat{\mathbf{Y}}_{\hat{\delta}}$ by the following relative mean square error (RMSE):

$$\text{RMSE}[\hat{\mathbf{Y}}_{\hat{\delta}}] = \frac{\text{MSE}[\hat{\mathbf{Y}}_{\hat{\delta}}]}{p(k+1)} \times 100(\%),$$

$$\text{MSE}[\hat{\mathbf{Y}}_{\hat{\delta}}] = \text{E} \left[\text{tr} \left\{ (\mathbf{X}\Xi - \hat{\mathbf{Y}}_{\hat{\delta}})'(\mathbf{X}\Xi - \hat{\mathbf{Y}}_{\hat{\delta}})\Sigma^{-1} \right\} \right].$$

In this setting, $\text{MSE}[\hat{\mathbf{Y}}] = p(k+1)$. This means that the prediction accuracies are evaluated in terms of the amount of improvement of the prediction accuracy of $\hat{\mathbf{Y}}$. Specifically, $\text{RMSE} < 100$ means the prediction accuracy of $\hat{\mathbf{Y}}_{\hat{\delta}}$ is superior to that of $\hat{\mathbf{Y}}$ and $\text{RMSE} > 100$ means the prediction accuracy of $\hat{\mathbf{Y}}_{\hat{\delta}}$ is inferior to that of $\hat{\mathbf{Y}}$. The smaller the RMSE value, the better the prediction accuracy. The expectation of the MSE is evaluated by Monte Carlo simulation with 10,000 iterations. Furthermore, it can be considered that the MSE value strongly relates to the amount of shrinkage of the MGR estimator, in particular, more shrinkage is required when there are highly correlated variables in \mathbf{X} . When $\hat{\delta}_j = 1$,

the amount of shrinkage of the MGR estimator is maximized and this means that the j th eigenvalue (and corresponding eigenvector) is removed from the model. From this, we measure the amount of shrinkage of the MGR estimator by calculating the following relative number of removed eigenvalues (RNRE):

$$\text{RNRE}(\hat{\delta}) = \frac{\#\{j \in \{1, \dots, k\} \mid \hat{\delta}_j = 1\}}{k} \times 100 (\%).$$

The RNRE expresses the ratio of the number of removed eigenvalues. If the RNRE value is small (large), then the amount of shrinkage is also small (large).

In this simulation, we estimate the mean structure of model. Thus, we use the LH-, LR-, and BNP-distances as the distance in the MSC. RMSE comparison 1 explores the prediction accuracies of predictive matrices where ridge parameters are optimized by the following methods:

- GC_p : GC_p criterion minimization method.
- EGCV: EGCV criterion minimization method.
- GIC: GIC minimization method via the iterative method with the initial vector $\mathbf{0}_k$.
- BNPC: BNPC minimization method via the iterative method with the initial vector $\mathbf{0}_k$.
- PIM1: PIM with EGCV criterion and GIC.
- PIM2: PIM with EGCV criterion and BNPC.

For all MSCs, we use $\alpha = 2, 2 \log \log n, \log n$, and they are labeled as 1, 2, and 3, respectively. Furthermore, the quartic equation in the BNPC minimization method is solved by the R function “polyroot”.

Table 1 summarizes the RMSE and RNRE values for $\rho_y = 0.2, 0.5, 0.9$ and $k = 0.1n, 0.3n, 0.5n$ when $p = 5$ and $n = 50$. From this table, it can be discerned that the prediction accuracy of $\hat{\mathbf{Y}}_{\hat{\delta}}$ is greater than that of $\hat{\mathbf{Y}}$ in most cases. We also found that the RNRE values increase as α increases, i.e., as the amount of shrinkage increases. Moreover, the RMSE values tend to increase with increasing ρ_y or k , and this may be caused by decreasing shrinkage. However, we could not find relationships between prediction accuracies and the amount of shrinkage. Table 2 summarizes the results when $p = 5$ and $n = 200$. Overall, trends are similar to those in Table 1. However, when $n = 200$ the amount of shrinkage substantially decreases. Table 3 summarizes the results when $p = 5$ and $n = 500$. In this case, the optimal ridge parameters often do not lead to improvements in prediction accuracies. This is because the amount of shrinkage is too large for the BNPC and too small for the methods. Tables 4 – 6 show the results when $p = 10$, and we can see that trends are similar compared to the case where $p = 5$.

In RMSE comparison 1, the iteration method was used to optimize the ridge parameters using the GIC and BNPC minimization methods. However, these optimal ridge parameters can also be calculated by using the coordinate descent algorithm or the PIM with the GC_p criterion. RMSE comparison 2 confirms whether the three algorithms minimize the MSC or not by comparing the results obtained from these algorithms. Although the initial vector used

TABLE 1 RMSE comparison 1 when $p = 5$ and $n = 50$

		ρ_y k	0.2			0.5			0.9		
			5	15	25	5	15	25	5	15	25
GC_p	1	RMSE	49.40	47.52	48.94	51.43	50.44	52.18	65.17	66.74	71.24
		RNRE	36.22	31.99	28.52	34.80	29.95	26.40	24.02	19.03	14.69
	2	RMSE	45.30	44.06	45.24	47.60	47.54	49.20	63.96	66.23	71.53
		RNRE	50.10	45.39	41.63	48.26	42.82	38.77	34.74	28.39	22.35
	3	RMSE	43.19	43.69	44.27	45.65	48.03	49.31	66.18	69.91	76.91
		RNRE	64.97	60.97	58.01	63.32	57.91	54.72	47.83	40.62	33.31
EGCV	1	RMSE	49.59	48.21	50.37	51.60	51.01	53.33	65.21	66.88	71.40
		RNRE	35.72	30.11	25.18	34.27	28.31	23.49	23.80	18.35	13.68
	2	RMSE	45.29	44.00	44.99	47.57	47.47	48.97	63.95	66.28	72.02
		RNRE	50.34	45.71	43.00	48.50	43.29	40.47	35.08	29.25	24.39
	3	RMSE	43.12	44.08	45.02	45.60	48.70	51.01	66.52	71.98	86.35
		RNRE	66.00	64.46	66.79	64.36	61.71	64.07	49.11	44.60	43.72
GIC	1	RMSE	50.02	50.80	57.71	52.08	53.59	60.48	65.78	69.00	76.44
		RNRE	37.11	27.27	16.50	35.57	25.32	14.83	24.39	14.91	6.82
	2	RMSE	45.21	44.74	47.42	47.60	48.28	51.31	64.38	66.66	72.30
		RNRE	53.65	46.23	38.21	51.77	43.42	34.76	37.31	27.15	15.71
	3	RMSE	42.88	44.54	45.42	45.51	49.67	51.73	67.94	72.38	83.41
		RNRE	70.30	68.78	70.97	69.00	66.07	68.09	53.81	46.64	42.07
BNPC	1	RMSE	48.17	45.60	50.84	50.42	49.82	61.45	65.45	68.53	145.94
		RNRE	46.53	54.76	85.12	44.61	51.70	84.47	31.15	29.32	57.38
	2	RMSE	43.73	48.99	59.09	46.39	57.16	75.94	66.25	102.30	***
		RNRE	66.23	79.62	91.35	64.70	78.68	91.40	49.52	62.36	91.06
	3	RMSE	42.80	66.04	89.59	45.60	79.82	118.06	76.70	***	***
		RNRE	78.76	88.69	92.94	78.34	88.57	93.34	69.75	85.94	94.79
PIM1	1	RMSE	48.84	47.34	48.91	50.94	50.30	52.11	65.06	66.78	71.46
		RNRE	39.93	34.80	30.91	38.20	32.79	29.01	26.84	21.69	17.51
	2	RMSE	44.59	43.59	44.28	47.01	47.43	48.84	64.38	67.36	75.72
		RNRE	56.00	52.84	52.60	54.31	50.29	50.00	39.97	35.09	32.80
	3	RMSE	42.80	45.39	46.58	45.43	51.05	53.90	68.89	77.46	103.47
		RNRE	71.71	72.21	76.51	70.49	69.92	74.35	55.97	52.99	56.55
PIM2	1	RMSE	48.02	46.50	47.93	50.22	49.78	51.97	65.02	67.28	78.35
		RNRE	45.41	42.84	44.79	43.57	40.82	42.86	31.25	27.98	30.58
	2	RMSE	43.90	44.34	46.32	46.49	49.30	52.71	65.52	72.54	103.73
		RNRE	63.43	63.54	67.10	61.89	61.48	65.29	47.13	46.20	52.24
	3	RMSE	42.64	49.52	50.09	45.46	57.84	59.65	73.92	113.98	***
		RNRE	77.31	80.65	84.19	76.70	79.66	82.99	65.95	67.35	73.37

Note: Emboldened entries represent the minimum of the RMSE values in each column; *** denotes values greater than 150.

in the iterative method is $\mathbf{0}_k$, the PIM with the GC_p criterion is the iterative method by changing the initial vector from $\mathbf{0}_k$ to the ridge parameters optimized by the GC_p criterion minimization method. Hence, by comparing the results obtained from the two methods, we can confirm whether the iterative method depends on the initial vector or not.

Table 7 compares the three algorithms for solving the GIC minimization method in terms of the RMSE, i.e., from the iterative method (GIC_IM), the coordinate descent algorithm (GIC_CD), and the PIM with the GC_p criterion

TABLE 2 RMSE comparison 1 when $p = 5$ and $n = 200$

		$\rho_{y,k}$	0.2			0.5			0.9		
			20	60	100	20	60	100	20	60	100
GC_p	1	RMSE	68.57	75.92	81.39	73.54	80.32	84.51	89.94	92.30	94.24
		RNRE	19.37	13.63	9.84	15.86	10.71	7.94	5.17	3.77	2.56
	2	RMSE	73.88	82.70	89.17	80.85	88.41	92.48	97.35	98.59	99.68
		RNRE	37.35	26.81	20.12	31.19	21.43	16.44	12.30	7.62	5.63
	3	RMSE	89.84	101.56	110.81	100.53	110.59	114.55	117.17	116.58	115.30
		RNRE	55.56	41.74	32.45	48.38	34.31	27.06	21.37	12.67	9.73
EGCV	1	RMSE	68.56	75.88	81.27	73.52	80.25	84.39	89.93	92.27	94.20
		RNRE	19.16	13.23	9.34	15.73	10.45	7.62	5.16	3.74	2.52
	2	RMSE	74.32	84.71	95.15	81.45	90.81	98.34	97.81	99.70	101.84
		RNRE	38.18	29.04	24.51	31.99	23.36	20.07	12.62	8.08	6.43
	3	RMSE	92.62	115.19	***	104.26	129.24	***	120.70	128.37	148.20
		RNRE	57.77	48.88	48.61	50.87	42.05	42.90	22.52	15.02	14.72
GIC	1	RMSE	68.76	76.45	83.20	73.60	80.41	85.79	89.66	91.89	94.34
		RNRE	18.08	9.19	3.57	14.79	7.07	2.79	4.57	2.29	0.75
	2	RMSE	74.21	80.60	83.03	81.29	85.67	85.72	96.60	94.62	94.11
		RNRE	38.07	24.25	12.93	31.78	18.89	9.73	11.91	5.69	2.35
	3	RMSE	94.07	110.50	136.68	106.84	123.88	141.34	120.95	114.66	103.06
		RNRE	59.24	47.18	42.03	52.74	40.57	35.23	22.57	12.38	6.67
BNPC	1	RMSE	68.88	76.24	82.13	73.89	80.43	85.06	89.88	91.92	94.25
		RNRE	20.41	12.45	5.71	16.51	9.24	4.03	5.01	2.61	0.87
	2	RMSE	78.89	119.39	***	87.55	147.83	***	99.54	99.52	***
		RNRE	45.23	51.28	94.39	38.44	49.56	95.01	14.02	8.05	5.59
	3	RMSE	110.61	***	***	130.66	***	***	***	***	***
		RNRE	68.76	83.54	97.03	65.39	85.89	97.02	30.66	94.16	97.15
PIM1	1	RMSE	68.67	76.04	81.62	73.68	80.49	84.80	90.06	92.44	94.41
		RNRE	19.83	13.84	10.06	16.28	10.99	8.23	5.39	3.93	2.72
	2	RMSE	75.29	86.59	101.64	82.87	93.71	106.56	98.83	101.75	105.66
		RNRE	39.87	30.98	27.91	33.62	25.36	23.46	13.40	8.83	7.37
	3	RMSE	95.74	121.86	***	109.29	141.32	***	126.93	147.81	***
		RNRE	60.36	52.26	53.79	54.14	47.10	50.12	24.11	18.30	25.30
PIM2	1	RMSE	68.80	76.32	82.79	73.88	80.89	86.11	90.21	92.67	94.78
		RNRE	20.62	14.67	11.50	16.91	11.76	9.49	5.64	4.17	3.01
	2	RMSE	76.58	90.23	122.20	84.85	100.37	140.97	100.18	106.38	***
		RNRE	41.95	34.07	35.28	35.68	29.12	32.68	14.36	10.24	15.98
	3	RMSE	100.12	133.69	***	116.54	***	***	139.98	***	***
		RNRE	63.44	56.30	59.23	58.54	53.67	58.34	26.59	34.39	67.16

Note: Emboldened entries represent the minimum of the RMSE values in each column; *** denotes values greater than 150.

(PIM- GC_p). Settings are as per RMSE comparison 1, where α is only $\alpha = 2$. From these results, it can be discerned that there is equivalent performance among the three algorithms. Although there is a bit of error, it can be considered that the error is made when convergence judgment. Thus, the three algorithms all converge and achieve minimization of the GIC. Furthermore, we found that the iterative method does not depend on the initial vector.

Table 8 shows a runtime comparison of the three algorithms for the GIC minimization method in terms of time (s) per repeat, where the reported values

TABLE 3 RMSE comparison 1 when $p = 5$ and $n = 500$

		ρ_y	0.2			0.5			0.9		
		k	50	150	250	50	150	250	50	150	250
GC_p	1	RMSE	87.42	94.49	94.98	89.93	95.84	96.11	97.21	99.19	98.84
		RNRE	6.75	2.27	2.14	5.23	1.58	1.56	1.11	0.11	0.40
	2	RMSE	98.40	104.50	103.78	100.49	105.05	104.03	103.37	103.50	102.07
		RNRE	15.90	6.95	5.71	12.45	5.07	4.13	3.62	0.60	1.07
	3	RMSE	130.30	131.79	129.36	130.42	130.22	127.53	120.89	116.88	112.61
		RNRE	26.47	14.69	11.48	21.95	11.59	8.63	7.77	1.97	2.09
EGCV	1	RMSE	87.40	94.47	94.95	89.92	95.83	96.08	97.20	99.19	98.84
		RNRE	6.72	2.26	2.11	5.21	1.57	1.55	1.11	0.11	0.40
	2	RMSE	99.18	106.79	108.56	101.17	106.90	107.68	103.57	103.84	102.57
		RNRE	16.33	7.79	7.05	12.76	5.66	4.97	3.70	0.63	1.14
	3	RMSE	137.37	***	***	136.36	149.81	***	122.82	121.41	121.10
		RNRE	27.93	19.22	23.61	23.31	15.38	18.15	8.11	2.38	2.66
GIC	1	RMSE	87.10	93.58	94.63	89.59	94.93	95.70	96.92	98.70	98.62
		RNRE	5.94	1.17	0.60	4.59	0.78	0.44	0.92	0.04	0.10
	2	RMSE	97.08	98.71	95.24	99.03	99.46	96.23	102.06	100.40	98.76
		RNRE	15.27	4.64	2.32	11.84	3.19	1.62	3.24	0.25	0.37
	3	RMSE	134.11	129.57	115.52	132.87	126.22	111.41	119.31	108.82	101.60
		RNRE	27.37	14.24	8.70	22.65	10.65	5.69	7.62	1.20	1.00
BNPC	1	RMSE	87.22	93.62	94.55	89.70	94.98	95.65	96.96	98.71	98.62
		RNRE	6.28	1.30	0.67	4.82	0.86	0.48	0.95	0.04	0.11
	2	RMSE	100.13	103.66	***	101.56	102.85	97.88	102.71	100.70	98.84
		RNRE	17.00	6.77	4.32	13.10	4.41	2.34	3.51	0.29	0.41
	3	RMSE	***	***	***	***	***	***	124.39	114.88	***
		RNRE	32.58	98.69	100*	27.14	99.36	100*	8.64	1.85	1.43
PIM1	1	RMSE	87.44	94.54	95.05	89.97	95.89	96.19	97.23	99.21	98.86
		RNRE	6.83	2.31	2.19	5.29	1.61	1.60	1.13	0.12	0.41
	2	RMSE	99.66	107.65	111.08	101.67	107.89	110.47	103.83	104.11	103.06
		RNRE	16.66	8.14	7.67	13.06	5.96	5.49	3.82	0.68	1.21
	3	RMSE	140.71	***	***	139.49	***	***	124.29	124.55	132.50
		RNRE	28.64	20.44	28.29	24.08	16.84	24.37	8.46	2.66	3.24
PIM2	1	RMSE	87.49	94.61	95.20	90.01	95.97	96.34	97.26	99.23	98.89
		RNRE	6.94	2.38	2.28	5.38	1.66	1.67	1.16	0.12	0.43
	2	RMSE	100.23	108.98	119.90	102.25	109.44	121.35	104.11	104.46	103.86
		RNRE	17.03	8.64	9.25	13.37	6.39	7.14	3.97	0.73	1.31
	3	RMSE	145.66	***	***	143.99	***	***	126.11	131.62	***
		RNRE	29.54	23.09	35.64	25.03	20.81	38.87	8.88	3.19	59.05

Note: Emboldened entries represent the minimum of the RMSE values in each column; *** denotes values greater than 150; * denotes an exact value.

are 10,000 times the actual runtime values. The PIM is the fastest algorithm in most cases. Sometimes the iterative method is faster than the PIM. In such cases, the RNRE value is very small, that is, the optimal ridge parameters are close to 0. The difference between the iterative method and the PIM is the initial vector, and the initial vector of the iterative method is $\mathbf{0}_k$. Hence, it can be considered that the iterative method is faster than the PIM when the optimal ridge parameters are close to 0. On the other hand, the coordinate

TABLE 4 RMSE comparison 1 when $p = 10$ and $n = 50$

		ρ_y k	0.2			0.5			0.9		
			5	15	25	5	15	25	5	15	25
GC_p	1	RMSE	47.25	43.85	48.37	49.21	46.54	51.65	61.62	62.52	68.72
		RNRE	24.78	19.56	12.83	23.24	18.20	11.48	14.52	10.18	5.72
	2	RMSE	42.71	38.81	42.17	44.95	41.82	46.06	59.32	60.16	65.84
		RNRE	42.72	35.07	24.76	40.42	32.90	22.35	27.17	19.76	11.70
	3	RMSE	40.61	37.16	38.43	43.07	40.62	43.24	60.65	61.73	66.31
		RNRE	63.05	55.28	43.57	60.99	52.36	39.81	43.92	34.58	22.26
EGCV	1	RMSE	47.36	44.27	49.05	49.30	46.90	52.21	61.65	62.63	68.87
		RNRE	24.49	18.68	11.86	22.94	17.40	10.71	14.44	9.94	5.53
	2	RMSE	42.60	38.34	40.39	44.85	41.40	44.55	59.30	60.09	65.50
		RNRE	43.50	37.75	31.19	41.25	35.54	28.33	27.92	21.71	14.89
	3	RMSE	40.55	37.68	38.50	43.03	41.41	44.97	61.08	63.98	73.76
		RNRE	64.88	62.77	63.23	62.91	59.69	59.41	46.10	41.77	37.62
GIC	1	RMSE	46.61	46.01	58.58	48.71	48.85	61.80	61.73	65.21	76.91
		RNRE	32.68	21.04	6.86	30.68	19.21	5.69	19.44	9.33	1.96
	2	RMSE	41.50	37.85	40.88	44.02	41.08	45.91	59.74	61.02	69.04
		RNRE	58.42	50.97	40.64	56.12	47.47	34.49	39.19	27.44	10.41
	3	RMSE	40.14	40.11	40.84	42.93	44.33	49.30	65.66	68.82	81.02
		RNRE	77.07	78.59	82.57	76.24	75.06	78.72	60.97	57.74	51.56
BNPC	1	RMSE	42.71	41.12	51.02	45.39	45.33	68.20	61.06	68.18	***
		RNRE	64.79	79.62	91.59	62.73	75.73	91.38	44.03	54.31	78.60
	2	RMSE	39.83	52.83	72.67	42.90	62.50	97.41	68.37	91.06	***
		RNRE	79.04	89.89	93.29	78.69	87.98	93.56	67.08	73.92	92.37
	3	RMSE	41.21	61.96	111.45	43.32	79.90	138.42	82.95	***	***
		RNRE	80.03	93.12	95.21	79.99	92.84	95.38	78.97	83.33	94.81
PIM1	1	RMSE	45.28	41.15	43.12	47.42	44.00	46.92	60.75	61.49	66.64
		RNRE	36.96	32.41	27.88	34.84	30.58	25.62	23.15	18.80	14.55
	2	RMSE	41.01	37.02	37.71	43.57	40.32	43.17	59.85	62.00	70.16
		RNRE	61.72	59.62	61.03	59.60	56.79	57.30	43.06	39.97	37.79
	3	RMSE	40.20	41.42	42.05	42.99	46.22	52.06	67.20	73.52	102.37
		RNRE	77.91	81.02	85.84	77.25	77.75	83.09	63.07	64.21	65.64
PIM2	1	RMSE	42.94	39.22	41.39	45.49	42.37	47.38	60.56	64.27	80.77
		RNRE	61.28	61.61	64.39	59.20	59.00	61.61	43.11	44.50	46.22
	2	RMSE	39.90	41.28	42.22	42.97	46.11	52.02	66.13	73.27	111.18
		RNRE	78.32	81.28	83.70	77.78	78.42	81.46	64.66	65.39	67.05
	3	RMSE	40.45	53.45	48.40	43.14	63.21	63.07	81.84	95.55	***
		RNRE	79.99	90.19	91.41	79.97	88.32	90.86	78.23	75.29	80.91

Note: Emboldened entries represent the minimum of the RMSE values in each column; *** denotes values greater than 150.

descent algorithm is overwhelmingly slowest of all. Hence, the best option for solving the GIC minimization method is to use the PIM with the GC_p criterion.

Table 9 compares the three algorithms for solving the BNPC minimization method, in terms of RMSE as similar to Table 7. It can be discerned that the three algorithms converge and achieve minimization of the BNPC, and the iterative method does not depend on the initial vector.

Table 10 shows a runtime comparison of the three algorithms for the BNPC minimization method in terms of time (s) as per Table 8. Similar to what

TABLE 5 RMSE comparison 1 when $p = 10$ and $n = 200$

		ρ_y k	0.2			0.5			0.9		
			20	60	100	20	60	100	20	60	100
GC_p	1	RMSE	57.38	67.35	72.47	62.60	72.10	76.68	83.29	88.44	90.86
		RNRE	16.40	10.25	7.31	13.21	8.01	5.61	3.39	2.11	1.34
	2	RMSE	61.78	73.26	78.22	68.67	78.80	82.61	90.30	93.85	95.20
		RNRE	40.86	27.18	20.49	33.81	21.51	16.27	11.54	6.27	4.25
	3	RMSE	76.54	92.15	98.54	87.31	100.35	103.24	111.52	111.78	110.88
		RNRE	63.02	45.94	36.23	55.35	38.00	29.93	23.44	13.10	8.85
EGCV	1	RMSE	57.39	67.37	72.49	62.61	72.10	76.68	83.29	88.44	90.86
		RNRE	16.24	10.00	6.98	13.11	7.86	5.43	3.38	2.10	1.33
	2	RMSE	62.35	76.04	85.74	69.43	81.88	89.90	90.91	95.27	97.88
		RNRE	42.32	31.11	27.98	35.16	24.74	22.49	12.08	6.98	5.29
	3	RMSE	79.09	107.06	149.90	90.91	119.41	***	115.99	126.01	***
		RNRE	65.46	54.49	55.01	58.39	47.89	48.46	25.20	17.08	17.74
GIC	1	RMSE	57.70	68.76	77.17	62.88	73.26	80.71	83.29	88.83	92.49
		RNRE	16.01	6.06	1.54	12.75	4.55	1.09	3.03	0.98	0.18
	2	RMSE	63.40	73.48	74.93	70.68	78.53	78.36	90.51	90.67	90.93
		RNRE	45.18	27.99	15.35	37.62	21.40	10.36	12.30	4.58	1.37
	3	RMSE	83.94	110.14	144.62	97.64	122.96	***	121.44	118.06	105.65
		RNRE	69.91	56.63	54.61	64.36	51.24	47.69	27.74	15.57	7.72
BNPC	1	RMSE	57.66	68.03	87.36	63.06	72.70	78.78	83.50	88.68	92.23
		RNRE	21.14	12.84	13.01	16.82	8.71	3.12	3.88	1.36	0.26
	2	RMSE	71.91	***	***	82.22	***	***	96.74	***	***
		RNRE	59.59	71.87	92.54	52.43	69.76	93.04	17.72	15.24	77.56
	3	RMSE	105.05	***	***	126.17	***	***	***	***	***
		RNRE	82.08	87.21	95.67	80.44	87.80	96.30	50.65	91.06	99.96
PIM1	1	RMSE	57.43	67.50	72.73	62.73	72.31	77.03	83.47	88.61	91.07
		RNRE	18.23	11.71	8.94	14.77	9.26	7.03	3.91	2.48	1.72
	2	RMSE	64.42	80.28	96.26	72.19	87.80	103.49	93.06	99.03	107.22
		RNRE	47.27	36.70	35.58	39.83	30.17	30.40	14.29	8.93	8.09
	3	RMSE	85.21	120.90	***	99.44	136.43	***	128.16	***	***
		RNRE	70.72	60.15	63.32	65.58	56.05	59.38	29.66	23.98	33.96
PIM2	1	RMSE	57.59	68.00	74.63	62.99	72.97	79.46	83.71	88.93	91.81
		RNRE	20.71	14.32	13.19	16.93	11.42	10.84	4.60	3.02	2.49
	2	RMSE	67.66	88.23	118.06	76.73	100.31	140.80	96.47	110.29	***
		RNRE	53.40	44.46	45.76	46.19	39.71	42.74	17.45	13.81	23.59
	3	RMSE	95.20	***	***	112.90	***	***	***	***	***
		RNRE	77.46	68.98	71.41	74.39	65.19	70.63	38.43	49.73	67.58

Note: Emboldened entries represent the minimum of the RMSE values in each column; *** denotes values greater than 150.

was noted above regarding the GIC minimization method, to solve the BNPC minimization method, using the PIM with the GC_p criterion is the best option.

Acknowledgement

I would like to express my deepest gratitude to my supervisor, Prof. Hirokazu Yanagihara of Hiroshima University, for stimulating my interest in statistics and for his invaluable support and guidance. I also thank Prof. Yasunori Fujikoshi, Prof. Hirofumi Wakaki, Dr. Keisuke Fukui, Dr. Shinpei Imori, Dr. Shintaro

TABLE 6 RMSE comparison 1 when $p = 10$ and $n = 500$

		ρ_y	0.2			0.5			0.9		
		k	50	150	250	50	150	250	50	150	250
GC_p	1	RMSE	81.19	88.94	90.00	85.14	91.51	92.14	95.37	97.82	97.67
		RNRE	4.32	1.56	1.66	2.93	0.89	1.12	0.38	0.03	0.20
	2	RMSE	93.28	98.48	97.82	96.38	100.05	99.03	100.55	101.06	100.33
		RNRE	15.09	7.32	6.46	10.64	4.69	4.47	2.03	0.27	0.77
	3	RMSE	131.47	128.21	124.25	132.40	127.22	122.64	119.74	113.93	111.01
		RNRE	28.73	17.16	13.68	22.07	12.58	10.26	4.66	1.06	1.63
EGCV	1	RMSE	81.19	88.93	89.99	85.14	91.51	92.14	95.36	97.82	97.67
		RNRE	4.31	1.55	1.64	2.93	0.89	1.11	0.38	0.03	0.20
	2	RMSE	94.53	101.57	103.97	97.40	102.38	103.49	100.75	101.35	100.85
		RNRE	15.80	8.72	8.72	11.11	5.54	5.89	2.07	0.29	0.83
	3	RMSE	142.15	***	***	141.78	***	***	122.24	118.84	122.10
		RNRE	31.45	23.46	30.84	24.22	17.55	22.35	4.93	1.35	2.43
GIC	1	RMSE	81.06	88.96	91.45	84.98	91.45	93.26	95.24	97.75	97.99
		RNRE	3.66	0.59	0.22	2.44	0.32	0.14	0.28	0.01	0.02
	2	RMSE	92.92	93.59	90.43	95.60	95.22	92.38	99.36	98.62	97.64
		RNRE	15.10	4.93	2.18	10.48	2.82	1.33	1.83	0.10	0.19
	3	RMSE	145.39	144.66	131.69	143.38	137.46	119.42	118.85	106.79	100.25
		RNRE	32.53	20.22	14.90	24.88	14.36	9.51	4.75	0.68	0.77
BNPC	1	RMSE	81.18	88.85	91.17	85.07	91.39	93.08	95.26	97.75	97.98
		RNRE	4.18	0.76	0.29	2.75	0.39	0.18	0.31	0.01	0.02
	2	RMSE	99.64	***	***	100.71	106.53	***	100.06	98.92	97.71
		RNRE	18.76	16.06	99.59	13.00	6.64	69.70	2.09	0.14	0.23
	3	RMSE	***	***	***	***	***	***	127.85	***	***
		RNRE	50.52	96.67	99.90	40.25	96.76	100.00	5.99	37.40	90.00
PIM1	1	RMSE	81.27	89.05	90.15	85.21	91.62	92.29	95.39	97.84	97.70
		RNRE	4.55	1.70	1.85	3.09	0.98	1.25	0.41	0.04	0.22
	2	RMSE	96.06	104.80	111.93	98.81	105.66	111.48	101.12	101.89	101.90
		RNRE	16.79	10.01	10.84	11.94	6.62	7.83	2.24	0.35	0.95
	3	RMSE	***	***	***	***	***	***	125.33	127.66	***
		RNRE	34.26	28.55	38.11	26.62	22.41	34.55	5.45	1.94	5.85
PIM2	1	RMSE	81.36	89.21	90.51	85.30	91.78	92.62	95.41	97.86	97.73
		RNRE	4.82	1.90	2.18	3.27	1.09	1.46	0.44	0.04	0.25
	2	RMSE	98.10	111.58	137.23	100.68	113.05	147.94	101.54	102.68	104.57
		RNRE	18.01	12.06	15.23	12.95	8.61	12.78	2.43	0.45	1.18
	3	RMSE	***	***	***	***	***	***	129.75	***	***
		RNRE	38.48	38.87	46.20	30.39	36.78	48.39	6.17	7.24	65.25

Note: Emboldened entries represent the minimum of the RMSE values in each column; *** denotes values greater than 150.

Hashimoto, and Dr. Ryoya Oda of Hiroshima University and Dr. Mariko Yamamura of Radiation Effects Research Foundation for their helpful comments and encouragement. Moreover, I thank the associate editor and the reviewer for their valuable comments.

TABLE 7 RMSE comparison 2 (GIC; $\alpha = 2$)

n	ρ_y	k	$p = 5$			$p = 10$		
			GIC.IM	GIC.CD	PIM.GC _p	GIC.IM	GIC.CD	PIM.GC _p
50	0.2	5	50.02	50.02	50.01	46.61	46.60	46.61
		15	50.80	50.79	50.78	46.01	45.99	45.97
		25	57.71	57.69	57.66	58.58	58.54	58.47
	0.5	5	52.08	52.08	52.07	48.71	48.70	48.70
		15	53.59	53.58	53.57	48.85	48.83	48.81
		25	60.48	60.46	60.43	61.80	61.77	61.71
	0.9	5	65.78	65.78	65.77	61.73	61.73	61.73
		15	69.00	68.99	68.99	65.21	65.20	65.18
		25	76.44	76.44	76.43	76.91	76.90	76.89
200	0.2	20	68.76	68.76	68.76	57.70	57.70	57.70
		60	76.45	76.45	76.45	68.76	68.76	68.76
		100	83.20	83.19	83.19	77.17	77.16	77.16
	0.5	20	73.60	73.60	73.60	62.88	62.88	62.88
		60	80.41	80.41	80.41	73.26	73.26	73.25
		100	85.79	85.79	85.79	80.71	80.71	80.70
	0.9	20	89.66	89.66	89.67	83.29	83.29	83.29
		60	91.89	91.89	91.89	88.83	88.83	88.83
		100	94.34	94.34	94.34	92.49	92.49	92.49
500	0.2	50	87.10	87.10	87.10	81.06	81.06	81.06
		150	93.58	93.58	93.58	88.96	88.96	88.96
		250	94.63	94.63	94.63	91.45	91.45	91.45
	0.5	50	89.59	89.59	89.59	84.98	84.98	84.98
		150	94.93	94.93	94.93	91.45	91.45	91.45
		250	95.70	95.70	95.70	93.26	93.26	93.26
	0.9	50	96.92	96.92	96.92	95.24	95.24	95.24
		150	98.70	98.70	98.70	97.75	97.75	97.75
		250	98.62	98.62	98.62	97.99	97.99	97.99

Appendix

A.1. Proof of Theorem 1. Let $r(\boldsymbol{\delta}) = \text{tr}(\mathbf{B}_{\boldsymbol{\delta}}^*)$ and $u(\boldsymbol{\delta}) = \text{df}(\boldsymbol{\delta})$. From Lemma 2, the domain of f is included in $[0, r_+] \times [p, np]$. We define $\tau(\boldsymbol{\delta})$ as

$$\tau(\boldsymbol{\delta}) = \frac{nbpf_u(r(\boldsymbol{\delta}), u(\boldsymbol{\delta}))}{2\dot{f}_r(r(\boldsymbol{\delta}), u(\boldsymbol{\delta}))}.$$

It is straightforward that $\tau(\boldsymbol{\delta}) > 0$ from f satisfies Definition 1. Then, we have

$$\begin{aligned} \frac{\partial}{\partial \delta_j} \text{MSC}(\boldsymbol{\delta}) &= \frac{\partial}{\partial \delta_j} r(\boldsymbol{\delta}) \cdot \frac{\partial}{\partial r} f(r, u) \Big|_{(r, u) = (r(\boldsymbol{\delta}), u(\boldsymbol{\delta}))} \\ &\quad + \frac{\partial}{\partial \delta_j} u(\boldsymbol{\delta}) \cdot \frac{\partial}{\partial u} f(r, u) \Big|_{(r, u) = (r(\boldsymbol{\delta}), u(\boldsymbol{\delta}))} \\ &= \frac{2}{nb} \mathbf{z}'_j \mathbf{S}^{-1} \mathbf{z}_j \delta_j \dot{f}_r(r(\boldsymbol{\delta}), u(\boldsymbol{\delta})) - p \dot{f}_u(r(\boldsymbol{\delta}), u(\boldsymbol{\delta})) \\ &= \frac{2}{nb} \mathbf{z}'_j \mathbf{S}^{-1} \mathbf{z}_j \dot{f}_r(r(\boldsymbol{\delta}), u(\boldsymbol{\delta})) \left(\delta_j - \frac{\tau(\boldsymbol{\delta})}{\mathbf{z}'_j \mathbf{S}^{-1} \mathbf{z}_j} \right), \end{aligned}$$

TABLE 8 Runtime comparison (GIC; $\times 1/10,000$ (s))

n	ρ_y	k	$p = 5$			$p = 10$		
			GIC.IM	GIC.CD	PIM.GC _p	GIC.IM	GIC.CD	PIM.GC _p
50	0.2	5	3.04	10.81	2.30	4.57	12.46	3.48
		15	5.15	43.41	3.76	8.78	52.13	6.10
		25	8.06	95.43	5.87	13.65	122.30	10.74
	0.5	5	3.40	10.97	2.09	4.65	12.41	3.53
		15	6.04	46.88	3.66	8.86	52.57	6.45
		25	8.48	104.61	5.87	13.57	124.52	11.11
	0.9	5	3.52	12.28	2.23	5.70	12.36	4.05
		15	6.04	46.51	3.80	8.97	52.90	6.70
		25	8.74	104.99	6.15	13.61	121.29	11.77
200	0.2	20	4.00	52.16	2.76	4.99	57.31	3.62
		60	7.51	205.03	5.26	9.66	228.49	6.95
		100	15.20	443.87	12.82	20.92	489.25	18.33
	0.5	20	3.96	52.96	2.75	4.92	58.35	3.43
		60	7.64	207.20	5.77	10.24	227.12	7.16
		100	15.74	450.99	13.95	22.31	504.68	20.00
	0.9	20	3.74	49.59	2.53	4.69	54.66	3.40
		60	6.12	174.76	4.48	9.25	197.91	7.01
		100	9.30	303.43	9.91	16.13	379.86	13.15
500	0.2	50	4.70	128.49	3.16	5.86	137.78	4.05
		150	13.66	456.75	10.85	23.53	528.58	20.47
		250	41.80	851.61	38.38	81.79	1051.05	54.42
	0.5	50	4.50	126.82	3.24	5.87	134.42	3.99
		150	13.52	440.54	11.35	22.19	521.07	18.68
		250	34.72	798.15	37.24	67.56	986.76	47.96
	0.9	50	3.91	109.08	2.66	5.13	112.50	3.37
		150	10.16	348.91	10.19	15.00	360.90	14.80
		250	21.28	559.92	26.01	32.38	607.38	37.82

Note: Emboldened entries represent the fastest time in each row.

$$\left. \frac{\partial}{\partial \delta_j} \text{MSC}(\boldsymbol{\delta}) \right|_{\boldsymbol{\delta}=\mathbf{0}_k} < 0.$$

Let $\boldsymbol{\delta}^* = (\delta_1^*, \dots, \delta_k^*)'$ be the minimizer of $\text{MSC}(\boldsymbol{\delta})$. Then, $\delta_j^* \neq 0$ ($j = 1, \dots, k$), and the necessary condition of δ_j^* is given by

$$\delta_j^* = \begin{cases} \frac{\tau(\boldsymbol{\delta}^*)}{\mathbf{z}'_j \mathbf{S}^{-1} \mathbf{z}_j} & (\tau(\boldsymbol{\delta}^*) < \mathbf{z}'_j \mathbf{S}^{-1} \mathbf{z}_j) \\ 1 & (\tau(\boldsymbol{\delta}^*) \geq \mathbf{z}'_j \mathbf{S}^{-1} \mathbf{z}_j) \end{cases} \quad (j = 1, \dots, k).$$

Let \mathcal{G} be a set defined by

$$\mathcal{G} = \left\{ \boldsymbol{\delta} \in [0, 1]^k \mid \boldsymbol{\delta} = \hat{\boldsymbol{\delta}}(h), \forall h \in \mathbb{R}_+ \setminus \{0\} \right\},$$

where $\hat{\boldsymbol{\delta}}(h)$ is a k -dimensional vector of which the j th element is given by

$$\hat{\delta}_j(h) = \begin{cases} \frac{h}{\mathbf{z}'_j \mathbf{S}^{-1} \mathbf{z}_j} & (h < \mathbf{z}'_j \mathbf{S}^{-1} \mathbf{z}_j) \\ 1 & (h \geq \mathbf{z}'_j \mathbf{S}^{-1} \mathbf{z}_j) \end{cases} \quad (j = 1, \dots, k).$$

TABLE 9 RMSE comparison 2 (BNPC; $\alpha = 2$)

n	ρ_y	k	$p = 5$			$p = 10$		
			BNPC_IM	BNPC_CD	PIM_GC $_p$	BNPC_IM	BNPC_CD	PIM_GC $_p$
50	0.2	5	48.17	48.15	48.16	42.71	42.67	42.71
		15	45.60	45.58	45.60	41.12	47.36	41.12
		25	50.84	60.14	50.83	51.02	602.41	51.02
	0.5	5	50.42	50.41	50.42	45.39	45.36	45.39
		15	49.82	49.83	49.82	45.33	55.91	45.33
		25	61.45	87.35	61.45	68.20	1127.10	68.20
	0.9	5	65.45	65.45	65.45	61.06	61.12	61.06
		15	68.53	68.56	68.52	68.18	138.36	68.18
		25	145.94	510.55	146.63	193.23	9745.56	193.62
200	0.2	20	68.88	68.88	68.88	57.66	57.67	57.66
		60	76.24	76.24	76.24	68.03	68.04	68.03
		100	82.13	82.13	82.12	87.36	89.21	87.41
	0.5	20	73.89	73.89	73.89	63.06	63.06	63.06
		60	80.43	80.44	80.44	72.70	72.70	72.69
		100	85.06	85.06	85.05	78.78	78.77	78.74
	0.9	20	89.88	89.88	89.88	83.50	83.51	83.50
		60	91.92	91.92	91.92	88.68	88.68	88.68
		100	94.25	94.24	94.24	92.23	92.23	92.22
500	0.2	50	87.22	87.22	87.22	81.18	81.18	81.18
		150	93.62	93.62	93.62	88.85	88.85	88.85
		250	94.55	94.55	94.55	91.17	91.17	91.17
	0.5	50	89.70	89.70	89.70	85.07	85.08	85.08
		150	94.98	94.98	94.98	91.39	91.39	91.39
		250	95.65	95.65	95.65	93.08	93.08	93.08
	0.9	50	96.96	96.96	96.96	95.26	95.26	95.27
		150	98.71	98.71	98.71	97.75	97.75	97.75
		250	98.62	98.62	98.62	97.98	97.98	97.98

Then, from δ^* is the minimizer of $MSC(\delta)$, the following equation holds:

$$MSC(\delta^*) = \min_{\delta \in [0,1]^k \setminus \{\mathbf{0}_k\}} MSC(\delta) \leq \min_{\delta \in \mathcal{G}} MSC(\delta) = \min_{h \in \mathbb{R}_+ \setminus \{0\}} MSC(\hat{\delta}(h)).$$

However, because $\delta^* \in \mathcal{G}$ the following equation holds:

$$MSC(\delta^*) \geq \min_{\delta \in \mathcal{G}} MSC(\delta) = \min_{h \in \mathbb{R}_+ \setminus \{0\}} MSC(\hat{\delta}(h)).$$

These results lead to

$$MSC(\delta^*) = \min_{h \in \mathbb{R}_+ \setminus \{0\}} MSC(\hat{\delta}(h)),$$

and hence, we have

$$\delta^* = \hat{\delta}(\hat{h}), \quad \hat{h} = \arg \min_{h \in \mathbb{R}_+ \setminus \{0\}} MSC(\hat{\delta}(h)).$$

Consequently, Theorem 1 is proved.

TABLE 10 Runtime comparison (BNPC; $\times 1/10,000$ (s))

n	ρ_y	k	$p = 5$			$p = 10$		
			BNPC_IM	BNPC_CD	PIM_GC $_p$	BNPC_IM	BNPC_CD	PIM_GC $_p$
50	0.2	5	4.39	68.80	3.43	5.92	76.06	4.95
		15	9.73	337.69	8.10	11.66	281.44	10.57
		25	11.36	621.02	9.32	12.21	472.33	9.96
	0.5	5	4.34	71.33	3.45	6.07	73.57	5.23
		15	10.17	364.55	8.64	12.77	283.79	11.63
		25	12.20	650.78	10.31	13.28	482.18	10.76
	0.9	5	5.29	72.58	4.26	7.73	70.38	6.87
		15	12.20	418.82	9.86	16.47	320.11	14.59
		25	18.36	886.39	15.46	23.12	604.47	20.86
200	0.2	20	5.27	376.38	4.04	7.73	390.91	6.68
		60	17.64	2295.43	16.25	26.98	2116.54	24.41
		100	27.54	3635.11	26.46	53.83	4494.30	48.62
	0.5	20	5.47	388.16	4.37	8.05	414.10	7.12
		60	20.60	2500.19	19.32	28.91	2341.24	26.35
		100	28.81	3581.45	26.60	41.19	3368.07	39.14
	0.9	20	6.28	418.43	4.93	11.00	492.37	9.63
		60	15.07	1676.25	12.75	25.66	2106.34	25.16
		100	19.40	2759.43	17.84	45.79	3869.82	39.98
500	0.2	50	6.28	958.66	5.03	10.85	1175.67	9.16
		150	35.73	4509.76	29.75	68.50	6669.25	64.67
		250	59.00	6650.66	53.98	142.62	8408.83	127.37
	0.5	50	6.19	928.01	4.70	11.59	1274.18	10.46
		150	40.44	4488.88	34.88	53.09	4748.88	48.16
		250	52.95	5933.89	47.70	176.77	12455.95	148.31
	0.9	50	4.80	774.13	3.45	6.37	788.19	4.89
		150	14.96	2655.99	11.09	57.38	5380.94	55.70
		250	34.27	4518.84	39.54	118.36	8531.93	97.23

Note: Emboldened entries represent the fastest time in each row.

A.2. Proof of Lemma 4. To prove Lemma 4, it is sufficient to prove $\text{df}(h_1) \geq \text{df}(h_2)$. From Lemma 1, $\text{df}(h)$ is expressed as

$$\text{df}(h) = p + p \sum_{j=1}^k \text{soft}(1, h/\mathbf{z}'_j \mathbf{S}^{-1} \mathbf{z}_j).$$

Therefore, we have

$$\text{df}(h_1) - \text{df}(h_2) = p \sum_{j=1}^k \{ \text{soft}(1, h_1/\mathbf{z}'_j \mathbf{S}^{-1} \mathbf{z}_j) - \text{soft}(1, h_2/\mathbf{z}'_j \mathbf{S}^{-1} \mathbf{z}_j) \},$$

and regarding the RHS of the above equation, the following equation holds:

$$\text{soft}(1, h_1/\mathbf{z}'_j \mathbf{S}^{-1} \mathbf{z}_j) - \text{soft}(1, h_2/\mathbf{z}'_j \mathbf{S}^{-1} \mathbf{z}_j)$$

$$= \begin{cases} 0 & (\mathbf{z}'_j \mathbf{S}^{-1} \mathbf{z}_j \leq h_1) \\ 1 - \frac{h_1}{\mathbf{z}'_j \mathbf{S}^{-1} \mathbf{z}_j} > 0 & (h_1 < \mathbf{z}'_j \mathbf{S}^{-1} \mathbf{z}_j \leq h_2) \\ \frac{h_2 - h_1}{\mathbf{z}'_j \mathbf{S}^{-1} \mathbf{z}_j} > 0 & (h_2 < \mathbf{z}'_j \mathbf{S}^{-1} \mathbf{z}_j) \end{cases}.$$

Hence, $\text{df}(h_1) \geq \text{df}(h_2)$ holds with equality only when $t_k \leq h_1$. Consequently, Lemma 4 is proved.

A.3. Proof of Proposition 4. First, we prove (1) by reduction to absurdity. Let $\hat{h}_{\alpha_1} = t_k$ and suppose that $\hat{h}_{\alpha_2} \neq t_k$. Then, the definition \hat{h}_α gives

$$\phi(\hat{h}_{\alpha_2} | \alpha_1) \geq \phi(t_k | \alpha_1), \quad \phi(t_k | \alpha_2) \geq \phi(\hat{h}_{\alpha_2} | \alpha_2),$$

and we have $\hat{h}_{\alpha_2} \neq t_k \Rightarrow \hat{h}_{\alpha_2} < t_k$ from (P2) in Proposition 1. Furthermore, $\phi(h | \alpha) = \eta(h | \alpha - \alpha_0) \phi(h | \alpha_0)$ holds from the definition of $\phi(h | \alpha)$. Therefore, from Lemma 4, we have

$$\begin{aligned} \phi(t_k | \alpha_2) &= \eta(t_k | \alpha_2 - \alpha_1) \phi(t_k | \alpha_1) \\ &< \eta(\hat{h}_{\alpha_2} | \alpha_2 - \alpha_1) \phi(\hat{h}_{\alpha_2} | \alpha_1) = \phi(\hat{h}_{\alpha_2} | \alpha_2). \end{aligned}$$

However, this contradicts $\phi(t_k | \alpha_2) \geq \phi(\hat{h}_{\alpha_2} | \alpha_2)$. Hence, (1) is proved.

Next, regarding (2), it is sufficient to prove $\hat{h}_{\alpha_1} < \hat{h}_{\alpha_2}$. We approach this via reduction to absurdity again. Let $\alpha_1 < \alpha_2$ and suppose that $\hat{h}_{\alpha_2} \leq \hat{h}_{\alpha_1}$. Now, we have $\hat{h}_{\alpha_2} < t_k$ from $\hat{h}_{\alpha_2} \neq t_k$. Therefore,

$$\begin{aligned} \phi(\hat{h}_{\alpha_1} | \alpha_2) &= \eta(\hat{h}_{\alpha_1} | \alpha_2 - \alpha_1) \phi(\hat{h}_{\alpha_1} | \alpha_1) \\ &< \eta(\hat{h}_{\alpha_2} | \alpha_2 - \alpha_1) \phi(\hat{h}_{\alpha_2} | \alpha_1) = \phi(\hat{h}_{\alpha_2} | \alpha_2). \end{aligned}$$

However, this contradicts the definition of \hat{h}_{α_2} . Hence, (2) is proved.

Consequently, Proposition 4 is proved.

A.4. Proof of Proposition 5. First, we consider when $\forall j \in \{1, \dots, k\}, \delta_j^{(1)} \geq \delta_j^{(0)}$. Suppose that $\delta_j^{(i)} \geq \delta_j^{(i-1)}$ for all $j \in \{1, \dots, k\}$. Then, $\delta_j^{(i)}$ is updated as

$$\delta_j^{(i+1)} = \zeta_j(\boldsymbol{\delta}^{(i)}) = 1 - \text{soft} \left(1, \tau(\boldsymbol{\delta}^{(i)}) / \mathbf{z}_j^{*'} \dot{\mathbf{G}}(\mathbf{B}_{\boldsymbol{\delta}^{(i)}}^*) \mathbf{z}_j^* \right),$$

and we have

$$\tau(\boldsymbol{\delta}^{(i)}) \geq \tau(\boldsymbol{\delta}^{(i-1)}), \quad \mathbf{z}_j^{*'} \dot{\mathbf{G}}(\mathbf{B}_{\boldsymbol{\delta}^{(i)}}^*) \mathbf{z}_j^* \leq \mathbf{z}_j^{*'} \dot{\mathbf{G}}(\mathbf{B}_{\boldsymbol{\delta}^{(i-1)}}^*) \mathbf{z}_j^*.$$

This gives $\delta_j^{(i+1)} \geq \delta_j^{(i)}$ for all $j \in \{1, \dots, k\}$, and hence the sequence $\{\delta_j^{(i)}\}$ is a monotonically increasing sequence. Moreover, the sequence is bounded. Hence, the iterative method converges.

Next, we consider when $\forall j \in \{1, \dots, k\}, \delta_j^{(1)} \leq \delta_j^{(0)}$. Suppose that $\delta_j^{(i)} \leq \delta_j^{(i-1)}$ for all $j \in \{1, \dots, k\}$. Then, we have

$$\tau(\boldsymbol{\delta}^{(i)}) \leq \tau(\boldsymbol{\delta}^{(i-1)}), \quad \mathbf{z}_j^{*'} \dot{\mathbf{G}}(\mathbf{B}_{\boldsymbol{\delta}^{(i)}}^*) \mathbf{z}_j^* \geq \mathbf{z}_j^{*'} \dot{\mathbf{G}}(\mathbf{B}_{\boldsymbol{\delta}^{(i-1)}}^*) \mathbf{z}_j^*.$$

This gives $\delta_j^{(i+1)} \leq \delta_j^{(i)}$ for all $j \in \{1, \dots, k\}$, and hence the sequence $\{\delta_j^{(i)}\}$ is a monotonically decreasing sequence. Moreover, the sequence is bounded. Hence, the iterative method converges.

Consequently, Proposition 5 is proved.

A.5. Proof of Theorem 3. Now, we have

$$\dot{f}_j(0) = -\frac{c_{j,0}}{f_{j,1}(0)} < 0, \quad \dot{f}_j(\delta) = 0 \iff \delta = \frac{1 \pm \sqrt{1 - c_{j,2}c_{j,0}/c_{j,1}^2}}{c_{j,2}/c_{j,1}}.$$

Therefore, $\hat{\delta}_j \neq 0$ and the smaller of the two real distinct roots or the double root of the quadratic equation $\dot{f}_{j,2}(\delta) = 0$ is the local minimizer. Notice that $\delta \in [0, 1]$. Then, to obtain the minimizer of $f_j(\delta)$, it is sufficient to confirm whether the local minimizer is included in $[0, 1]$ or not.

When $1 - c_{j,2}c_{j,0}/c_{j,1}^2 \geq 0$, there is one local minimizer, and let this be $\tilde{\delta}_j$, i.e.,

$$\tilde{\delta}_j = \frac{1 - \sqrt{1 - c_{j,2}c_{j,0}/c_{j,1}^2}}{c_{j,2}/c_{j,1}}.$$

This is positive and the following equation holds when $c_{j,2} > c_{j,1}$:

$$\tilde{\delta}_j < 1 - \sqrt{1 - c_{j,2}c_{j,0}/c_{j,1}^2} < 1.$$

Hence, we can obtain (1) in Theorem 3.

When $1 - c_{j,2}c_{j,0}/c_{j,1}^2 < 0$, there are no stationary points, and therefore $f_j(\delta)$ is a monotonically decreasing function. Hence, we can obtain (2) in Theorem 3.

Consequently, Theorem 3 is proved.

A.6. Proof of Theorem 4. Now, we have

$$\dot{f}_j(0) = -\frac{c_{j,0}}{f_{j,1}(0)} < 0, \quad \dot{f}_j(\delta) = 0 \iff \dot{f}_{j,2}(\delta) = 0.$$

Thus $\hat{\delta}_j \neq 0$. Moreover, from $\delta \in [0, 1]$, minimizer candidates are local minimizers of $f_{j,2}(\delta)$ included in $(0, 1)$ and the right end point of the range. Hence, we can obtain the set of minimizer candidates \mathcal{S}_j by calculating stationary points of the cubic function $\dot{f}_{j,2}(\delta)$ and by confirming whether each stationary point is included in $(0, 1)$ or not. Consequently, Theorem 4 is proved.

A.7. Proof of Theorem 6. To prove the equivalence between the two estimators, it is sufficient to prove $\hat{h}_A = \hat{\lambda}_A$. The two terms which constitute the

MSC for optimizing ridge parameters are

$$\begin{aligned} \text{tr} \left\{ \hat{\Sigma}_R(\hat{\theta}(h | \mathbf{A})) \mathbf{A}^{-1} \right\} &= b \text{tr} \left(\hat{\Sigma}_0 \mathbf{A}^{-1} \right) \\ &\quad + \frac{1}{n} \sum_{j=1}^k \left\{ 1 - \text{soft} \left(1, h / \mathbf{z}'_j \mathbf{A}^{-1} \mathbf{z}_j \right) \right\}^2 \mathbf{z}'_j \mathbf{A}^{-1} \mathbf{z}_j, \\ \text{df}_R(\hat{\theta}(h | \mathbf{A})) &= p + p \sum_{j=1}^k \text{soft} \left(1, h / \mathbf{z}'_j \mathbf{A}^{-1} \mathbf{z}_j \right). \end{aligned}$$

On the other hand, when $w_j = 1/\|\hat{\gamma}_j\|$, from Lemma 5, the two terms which constitute the MSC for optimizing the tuning parameter are given by

$$\begin{aligned} \text{tr} \left\{ \hat{\Sigma}_L(\lambda) \mathbf{A}^{-1} \right\} &= b \text{tr} \left(\hat{\Sigma}_0 \mathbf{A}^{-1} \right) \\ &\quad + \frac{1}{n} \sum_{j=1}^k \left\{ 1 - \text{soft} \left(1, \lambda / \mathbf{z}'_j \mathbf{A}^{-1} \mathbf{z}_j \right) \right\}^2 \mathbf{z}'_j \mathbf{A}^{-1} \mathbf{z}_j, \\ \text{df}_L(\lambda) &= p + p \sum_{j=1}^k \text{soft} \left(1, \lambda / \mathbf{z}'_j \mathbf{A}^{-1} \mathbf{z}_j \right). \end{aligned}$$

Hence, for all $x \in \mathbb{R}_+$, the following equation holds:

$$\text{MSC}_R(\hat{\theta}(x | \mathbf{A}) | \mathbf{A}) = \text{MSC}_L(x | \mathbf{A}).$$

Thus $\hat{h}_\mathbf{A} = \hat{\lambda}_\mathbf{A}$ and consequently, Theorem 6 is proved.

References

- [1] H. Akaike, Information theory and an extension of the maximum likelihood principle, In: B. N. Pretrov, F. Csáki (Eds.), 2nd International Symposium on Information Theory, Akadémiai Kiadó, Budapest, **26** (1973), 267–281.
- [2] T. W. Anderson, An Introduction to Multivariate Statistical Analysis, John Wiley & Sons, Inc., New Jersey, 2003.
- [3] A. C. Atkinson, A note on the generalized information criterion for choice of a model, *Biometrika*, **67** (1980), 413–418.
- [4] K. A. Bollen, Structural Equations with Latent Variables, John Wiley & Sons, Inc., New York, 1989.
- [5] P. Craven & G. Wahba, Smoothing noisy data with spline functions: estimating the correct degree of smoothing by the method of generalized cross-validation, *Numer. Math.*, **31** (1979), 377–403.
- [6] A. C. David, Galois Theory, John Wiley & Sons, Inc., New Jersey, 2004.
- [7] D. L. Donoho & I. M. Johnstone, Ideal spatial adaptation by wavelet shrinkage, *Biometrika*, **81** (1994), 425–455.
- [8] Y. Fujikoshi & K. Satoh, Modified AIC and C_p in multivariate linear regression, *Biometrika*, **84** (1997), 707–716.
- [9] E. J. Hannan & B. G. Quinn, The determination of the order of an autoregression, *J. R. Stat. Soc. Ser. B. Stat. Methodl.*, **41** (1979), 190–195.
- [10] A. E. Hoerl & R. W. Kennard, Ridge regression: biased estimation for nonorthogonal problems, *Technometrics*, **12** (1970), 55–67.

- [11] C. M. Hurvich & C.-L. Tsai, Regression and time series model selection in small samples, *Biometrika*, **76** (1989), 297–307.
- [12] C. L. Mallows, Some comments on C_p , *Technometrics*, **15** (1973), 661–675.
- [13] Y. Mori & T. Suzuki, Generalized ridge estimator and model selection criteria in multivariate linear regression, *J. Multivariate Anal.*, **165** (2018), 243–261.
- [14] I. Nagai, H. Yanagihara & K. Satoh, Optimization of ridge parameters in multivariate generalized ridge regression by plug-in methods, *Hiroshima Math. J.*, **42** (2012), 301–324.
- [15] R. Nishii, Asymptotic properties of criteria for selection of variables in multiple regression, *Ann. Statist.*, **12** (1984), 758–765.
- [16] M. Ohishi, H. Yanagihara & Y. Fujikoshi, A fast algorithm for optimizing ridge parameters in a generalized ridge regression by minimizing a model selection criterion, *J. Statist. Plann. Inference*, **204** (2020), 187–205.
- [17] M. Ohishi, H. Yanagihara & S. Kawano, Equivalence between adaptive-Lasso and generalized ridge estimators in linear regression with orthogonal explanatory variables after optimizing regularization parameters, *Ann. Inst. Statist. Math.*, **72** (2020), 1501–1516.
- [18] G. Schwarz, Estimating the dimension of a model, *Ann. statist.*, **6** (1978), 461–464.
- [19] M. S. Srivastava, *Methods of Multivariate Statistics*, John Wiley & Sons, Inc., New York, 2002.
- [20] R. Tibshirani, Regression shrinkage and selection via the Lasso, *J. R. Stat. Soc. Ser. B. Stat. Methodl.*, **58** (1996), 267–288.
- [21] J.-P. Tignol, *Galois’ Theory of Algebraic Equations*, World Scientific Publishing, Singapore, 2001.
- [22] N. H. Timm, *Applied Multivariate Analysis*, Springer-Verlag, New York, 2002.
- [23] X. Xin, J. Hu & L. Liu, On the oracle property of a generalized adaptive elastic-net for multivariate linear regression with a diverging number of parameters, *J. Multivariate Anal.*, **162** (2017), 16–31.
- [24] H. Yanagihara, Explicit solution to the minimization problem of generalized cross-validation criterion for selecting ridge parameters in generalized ridge regression, *Hiroshima Math. J.*, **48** (2018), 203–222.
- [25] H. Yanagihara, I. Nagai & K. Satoh, A bias-corrected C_p criterion for optimizing ridge parameters in multivariate generalized ridge regression, *J. Appl. Statist.*, **38** (2009), 151–172 (in Japanese).
- [26] H. Zou, The adaptive Lasso and its oracle properties, *J. Amer. Statist. Assoc.*, **101** (2006), 1418–1429.

Mineaki Ohishi
Department of Mathematics
Graduate School of Science
Hiroshima University
Higashi-Hiroshima 739-8526 JAPAN
E-mail: mineaki-ohishi@hiroshima-u.ac.jp