

# 論文内容要旨

Impact of Prosthesis-Patient Mismatch After Transcatheter Aortic Valve Replacement on Changes in Cardiac Sympathetic Nervous Function Assessment by <sup>123</sup>I-Metaiodobenzylguanidine Myocardial Scintigraphy  
(経カテーテル的大動脈弁置換術後の患者人工弁ミスマッチが心臓交感神経機能の変化に及ぼす影響：<sup>123</sup>I-MIBG 心筋シンチグラフィでの評価)  
International Heart Journal, 2020, in press.

主指導教員：中野 由紀子教授

(医系科学研究科 循環器内科学)

副指導教員：東 幸仁教授

(原爆放射線医科学研究所 ゲノム障害病理)

副指導教員：栗栖 智准教授

(医系科学研究科 循環器内科学)

新田 和宏

(医歯薬保健学研究科 医歯薬学専攻)

## **Introduction**

Aortic stenosis (AS) is a cardiovascular disease which is common among elderly patients. Transcatheter aortic valve replacement (TAVR) has recently emerged as a minimally invasive therapeutic option. However, prosthesis-patient mismatch (PPM)—which occurs when the effective orifice area of the implanted prosthetic valve is too small in relation to the patient's body surface area—is an important potential complication of TAVR. <sup>123</sup>I-metaiodobenzylguanidine (MIBG) myocardial scintigraphy has been widely used for the assessment of CSN activity and represents a useful prognostic marker in patients with heart failure. However, the impact of PPM after TAVR on changes in CSN function remains unclear. The present study aimed to investigate the impact of PPM after TAVR on CSN activity using <sup>123</sup>I-MIBG scintigraphy.

## **Methods**

### **Study population**

We recruited a total of 117 patients with symptomatic severe AS who underwent TAVR between February 2016 and May 2019 at the Hiroshima University Hospital. Of these, 58 underwent <sup>123</sup>I-MIBG scintigraphy at baseline and at about 9 months after TAVR. Exclusion criteria were as follows: (1) prior cardiac surgery (n = 2), (2) presence of unstable pre-procedural conditions (n = 2), (3) undergoing TAVR with the trans-apical or trans-aortic approach (n = 3), and (4) permanent pacemaker implantation (n = 6) or hemodialysis (n = 1) after TAVR. Finally, 44 patients were enrolled in this study.

### **<sup>123</sup>I-metaiodobenzylguanidine scintigraphy**

<sup>123</sup>I-MIBG was administered intravenously at a dose of 111 MBq. Anterior planar images were obtained at 15 min (early image) and 210 min (delayed image) after injection using a dual-detector 90° γ-camera with a medium-energy general collimator. Early and delayed heart-to-mediastinum (H/M) ratios were calculated by measuring the average counts in each region. The washout rate (WR) was calculated by:  $[(H-M)_{\text{early}} - (H-M)_{\text{delayed}}/k] \times 100/(H-M)_{\text{early}}$ , with background subtraction and time-decay correction (k = time-decay coefficient). Changes between pre- and post-procedural <sup>123</sup>I-MIBG parameters were calculated using the following formulae: (1)  $\Delta H/M \text{ ratio} = (\text{post-procedural H/M ratio}) - (\text{baseline H/M ratio})$  and (2)  $\Delta WR = (\text{baseline WR}) - (\text{post-procedural WR})$ .

### **Transthoracic echocardiography**

We obtained echocardiographic data at baseline and at 7 days and 9 months after TAVR. The severity of PPM was classified using an indexed effective orifice area, with moderate defined as  $\geq 0.65$  and  $\leq 0.85$  cm<sup>2</sup>/m<sup>2</sup> and severe defined as  $< 0.65$  cm<sup>2</sup>/m<sup>2</sup>, and this was assessed by echocardiogram at 7 days after TAVR.

## **Results**

### **Patient characteristics prior to transcatheter aortic valve replacement in relation to patient-prosthesis mismatch**

There were 9 male and 35 female patients with a mean age of  $84 \pm 4$  years. There were no significant differences in terms of age, sex, body surface area, body mass index, past medical history, serum creatinine, medications, and transthoracic echocardiographic data between patients with PPM and those without PPM before TAVR. As for 123I-MIBG parameters before TAVR, patients without PPM had higher early H/M ratio and WR than those with PPM.

### **Patient characteristics prior to and after transcatheter aortic valve replacement in relation to patient-prosthesis mismatch**

The rate of overall and severe PPM among the study population was 39% ( $n = 17$ ) and 11% ( $n = 5$ ), respectively. Significant improvements in AVA, peak pressure gradient, and mean pressure gradient after TAVR were observed in patients with and without PPM. Those without PPM exhibited significant regression in LVMI after TAVR, whereas those with PPM did not. Patients without PPM exhibited significant improvements in delayed H/M ratio and WR after TAVR, while those with PPM did not.

### **Factors associated with improvements in 123I-metaiodobenzylguanidine parameters**

Multivariable linear regression analysis revealed PPM was a negative predictor of improvement in delayed H/M ratio. Furthermore, PPM and baseline WR were independent predictors of improvement in WR.

### **Discussion**

This investigation of the impact of PPM after TAVR on changes in CSN function using 123I-MIBG scintigraphy demonstrates the following: (1) delayed H/M ratio and WR improve significantly after TAVR for patients without PPM, whereas patients with PPM do not experience these improvements, and (2) PPM is a negative predictor of improvements in delayed H/M ratio and WR in patients undergoing TAVR.

Our findings of the impact of PPM after TAVR on changes in CSN function may be explained by the association between PPM and reduced hemodynamic improvement after TAVR. We found that patients with PPM experienced less favorable changes after TAVR compared to those without PPM including lower AVA, higher peak and mean pressure gradients, and limited LVMI regression. The small indexed effective orifice area and its residual high-pressure gradient may suppress improvements in 123I-MIBG parameters.

In conclusion, our findings suggest that delayed H/M ratio and WR improve significantly after TAVR for patients who do not experience PPM. Because of the negative effects in terms of improvement in 123I-MIBG parameters, strategies to prevent or minimize PPM are important, especially for patients with CSN overactivation.