Check for updates

OPEN ACCESS

1

2

3

4

5

6

7

8

9

10

11

12

13

14

15

16

17

18

19

20

21

22

23

24

25

26

27

28

29

30

31

32

33

34

35

36

37

38

39

40

41

42

43

44

45

46

47

48 49 Q5

EDITED BY Nina M. Van Sorge, Amsterdam University Medical Center, Netherlands

REVIEWED BY María Guembe, Gregorio Marañón Hospital, Spain Timothy J. Foster, Trinity College Dublin, Ireland Chia Y. Lee, University of Arkansas for Medical Sciences, United States *CORRESPONDENCE

Motoyuki Sugai Sugai@niid.go.jp

SPECIALTY SECTION

This article was submitted to Infectious Agents and Disease, a section of the journal Frontiers in Microbiology

RECEIVED 18 November 2022 ACCEPTED 14 December 2022 PUBLISHED XX XX 2022

CITATION

Kutsuno S, Hayashi I, Yu L, Yamada S, Hisatsune J and <mark>Sugai M</mark> (2022)

Non-deacetylated poly-*N*acetylglucosamine-hyperproducing *Staphylococcus aureus* undergoes immediate autoaggregation upon vortexing. *Front. Microbiol.* 13:1101545. doi: 10.3389/fmicb.2022.1101545

COPYRIGHT

© 2022 Kutsuno, Hayashi, Yu, Yamada, Hisatsune and Sugai. This is an openaccess article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.

Non-deacetylated poly-*N*-acetylglucosaminehyperproducing *Staphylococcus aureus* undergoes immediate autoaggregation upon vortexing

Shoko Kutsuno^{1,2}, Ikue Hayashi³, Liansheng Yu^{1,2}, Sakuo Yamada⁴, Junzo Hisatsune^{1,2} and Motoyuki Sugai^{1,2}*

¹Antimicrobial Resistance Research Center, National Institute of Infectious Diseases, Tokyo, Japan, ²Department of Antimicrobial Resistance, Hiroshima University Graduate School of Biomedical & Health Sciences, Hiroshima, Japan, ³Research Facility, Hiroshima University Faculty of Dentistry, Hiroshima, Japan, ⁴Department of Medical Technology, Faculty of Health Sciences & Technology, Kawasaki University of Medical Welfare, Kurashiki, Japan

Biofilms are microbial communities of cells embedded in a matrix of extracellular polymeric substances generated and adhering to each other or to a surface. Cell aggregates formed in the absence of a surface and floating pellicles that form biofilms at the air-liquid interface are also considered to be a type of biofilm. Staphylococcus aureus is a well-known cause of biofilm infections and high-molecular-weight polysaccharides, poly-N-acetylglucosamine (PNAG) is a main constituent of the biofilm. An *icaADBC* operon comprises major machinery to synthesize and extracellularly secrete PNAG. Extracellular PNAG is partially deacetylated by IcaB deacetylase, and the positively charged PNAG hence interacts with negatively charged cell surface to form the major component of biofilm. We previously reported a new regulator of biofilm (Rob) and demonstrated that Rob binds to a unique 5-bp motif, TATTT, present in intergenic region between *icaADBC* operon and its repressor gene *icaR* (Yu et al., 2017). The deletion of the 5-bp motif induces excessive adherent biofilm formation. The real function of the 5-bp motif is still unknown. In an attempt to isolate the 5-bp motif deletion mutant, we isolated several non-adherent mutants. They grew normally in turbid broth shaking culture but immediately auto-aggregated upon weak vortexing and sedimented as a lump resulting in a clear supernatant. Whole genome sequencing of the mutants identified they all carried mutations in *icaB* in addition to deletion of the 5-bp motif. Purification and molecular characterization of auto-aggregating factor in the culture supernatant of the mutant identified that the factor was a massively produced non-deacetylated PNAG. Therefore, we created a double deficient strain of biofilm inhibitory factors (5-bp motif, icaR, rob) and icaB to confirm the aggregation phenomenon. This peculiar phenomenon was only observed in $\Delta 5 bp \Delta i caB$ double mutant but not in $\Delta i caB$ or $\Delta rob \Delta i caB$ mutant. This study explains large amount of extracellularly produced non-deacetylated PNAG by $\Delta 5 bp \Delta i caB$ double mutation induced rapid auto-aggregation of S. aureus cells by vortexing. This phenomenon indicated that Staphylococcus

96

97

98

100

101

102

103

104

105

106

107

108

109

110

111

112

113

114

115

116

117

118

119

120

121

122

123

124

125

126

127

128

129

130

131

132

133

134

135

136

137

138

139

140

141

142

143

144

145

146

147

148

149

150

151

152

Q10

153

155

156

157

158

159

160

161

162

163

164

165

166

167

168

169

170

171

172

173

175

176

177

178

179

180

181

182

183

184

185

186

187

188

189

190

191

192

193

194

195

196

197

198

199

200

201

202

203

204

205

206

Q11 174

Q7 154

Q8

aureus may form biofilms that do not adhere to solid surfaces and we propose this as a new mechanism of non-adherent biofilm formation of *S. aureus*.

KEYWORDS

biofilm, *Staphylococcus aureus*, *icaB*, aggregation, poly-*N*-acetylglucosamine, PNAG

Introduction

Staphylococcus aureus is a facultative anaerobic grampositive coccus indigenous to human skin, pharynx, fecal, and nasal mucosa (Lowy, 1998; Claassen-Weitz et al., 2016; Mehraj et al., 2016; Byrd et al., 2018). S. aureus, the most common cause of nosocomial infections, accounts for a high percentage of isolates from hospitalized patients. S. aureus causes a variety of infections, including chronic biofilm infections, such as catheter bloodstream infections, osteomyelitis, and endocarditis. Such biofilm infections are caused by the persistent attachment of S. aureus to host tissues, such as bone and heart valves, and to implanted materials, such as catheters and prostheses (Parsek and Singh, 2003; Otto, 2008; Kiedrowski and Horswill, 2011; Barrett and Atkins, 2014; Di Domenico et al., 2022). Biofilmforming bacteria are more resistant to host defense mechanisms and drugs than non-biofilm-forming bacteria, and the removal of biofilms from indwelling catheters and artificial organs using drugs and immune cells tends to be difficult, resulting in a strong tendency for infections to become refractory (Chatterjee et al., 2014). Biofilms are defined as adherent microbial communities in which cells adhere to surfaces and other cells and are encased in a protective extracellular polymeric matrix (Costerton et al., 1978; Vaccari et al., 2017; Trunk et al., 2018). This mode of growth exhibits altered physiology with respect to gene expression and protein production (Archer et al., 2011; Schilcher and Horswill, 2020). The extracellular matrix is the basis for the attachment of bacteria to the surface of objects and is also involved in the binding of bacteria to each other (Flemming and Wingender, 2010). Its matrix is composed of various components such as nucleic acids, polysaccharides, proteins, and lipids, and the ratio of these components varies with environmental factors among strains as well as species (Mayer et al., 1999; Li et al., 2016). Therefore, the amounts of biofilms as well as their physicochemical and biochemical properties vary among different strains of the same bacterial species.

In general, the major components of staphylococcal biofilms are recognized as high-molecular-weight polysaccharides (poly-*N*-acetylglucosamine, PNAG or polysaccharide intercellular adhesion, PIA in *S. epidermidis*; Mack et al., 1996; Cue et al., 2012), surface protein (Foster et al., 2014), and eDNA (Montanaro et al., 2011; Ibáñez de Aldecoa et al., 2017). These facilitate the attachment of bacterial cells to the surface of objects, followed by bacterial colonization (Lister and Horswill, 2014). The process of PNAG/PIA formation on the cell surface has been well studied using the *S. epidermidis* model (Mack et al., 1996; Vuong et al., 2004; Le et al., 2018). As an extracellular polysaccharide, partially deacetylated PNAG has been obtained from a variety of bacterial sources such as *Escherichia coli* (Wang et al., 2004), *Klebsiella pneumoniae* (Chen et al., 2014), and *Acinetobacter baumannii* (Choi et al., 2009).

PNAG of Staphylococcus sp. is achieved via a combination of four gene products, icaA, icaD, icaB, and icaC, which are tandemly encoded on the ica operon of the chromosome (Heilmann et al., 1996a; Cramton et al., 1999). During ica operon-induced synthesis of PNAG, N-acetylglucosamine undergoes polymerization via the combined function of IcaA and IcaD, and then, the resulting N-acetylglucosamine polymers are exported via the IcaC transporter (Gerke et al., 1998). Exported N-acetylglucosamine polymers are partially deacetylated in situ by IcaB, and the products accumulate on the cell surface as PNAG (Vuong et al., 2004; Pokrovskaya et al., 2013). In the S. epidermidis model, extracellular IcaB partially deacetylated PNAG; subsequently, the positively charged deacetylated forms interacted with the negatively charged cell surface via electrostatic interactions and accumulated on the cell surface (Pokrovskaya et al., 2013). Several transcription factors are involved in the regulation of PNAG (Conlon et al., 2002; Jefferson et al., 2004; You et al., 2014; Yu et al., 2017). IcaR, located just upstream of the icaADBC operon, is a well-studied negative regulator that suppresses the *icaADBC* operon (Conlon et al., 2002; Jefferson et al., 2004). The product of rob (Yu et al., 2017), which is a newly discovered negative regulator (also known as GbaA (You et al., 2014)), binds to a 5-bp motif (TATTT) in the *icaR-icaA* intergenic region and suppresses the expression of *icaADBC*. The 5-base motif and rob are not present in S. epidermidis; thus, rob is a unique negative regulator of *icaADBC* in S. aureus. Jefferson et al. (2003) demonstrated that a clinically identified S. aureus strain, which produced a large amount of biofilm, lacked the 5-base motif, suggesting its importance in the regulation of biofilm production (Jefferson et al., 2003). It remains unclear whether the product of rob is the only regulator capable of recognizing the 5-base motif.

In order to further analyze the physiological function of the 5-base motif, a 5-base motif deletion mutant of *S. aureus* FK300, poor biofilm forming strain, was generated. Although most mutants showed normal colony morphology, a few revealed

262

263

264

265

266

267

268

269

270

271

272

273

274

275

276

277

278

279

280

281

282

283

284

285

286

287

288

289

290

291

292

293

294

295

296

297

298

299

300

301

302

303

304

305

306

307

308

309

310

311

312

313

314

207

peculiar phenotypes, as indicated by colonies showing shiny flat morphology and fusion with adjacent colonies.

These colonies grew normally in turbid broth shaking culture but immediately autoaggregated upon weak vortexing and sedimented as a lump resulting in a clear supernatant. These mutants completely lacked the ability to form biofilms on the surface.

In this study, we investigated the molecular mechanism underlying this autoaggregation just after the vortexing of mutants with a 5-base deletion.

Materials and methods

Bacterial strains and growth media

In this study, the shaking culture was performed in a water bath using a test tube of q15 mm×150mm with shaking at 140 rpm at an angle of 45°. The bacterial strains and plasmids used in this study are listed in Supplementary Table 1. Standard strain S. aureus FK300, a rsbU-repaired (Giachino et al., 2001) derivative of NCTC8325-4 (Novick, 1967; Herbert et al., 2010), was used in a functional study of the role of icaB. DNA restriction systemdeficient S. aureus RN4220 (Kreiswirth et al., 1983) was used as the initial recipient for manipulation of recombinant plasmids. S. aureus was routinely grown in brain heart infusion (Becton, Dickinson and Company, MD, United States) broth, tryptic soy broth (TSB; Becton, Dickinson and Company), or tryptic soy agar plates. Tetracycline (5 µg/ml) or chloramphenicol (5 µg/ml) was added to retain plasmids. The Escherichia coli strain DH5α was used for plasmid construction and maintenance. E. coli was grown in lysogeny broth (LB; 5 g yeast extract, 10 g polypeptone, and 10 g NaCl per liter; pH 7.2) or LB agar. Ampicillin (100 µg/ml) or tetracycline (12.5µg/ml) was added to the medium when the plasmid was to be retained.

Plasmid and strain construction

Routine DNA manipulation was performed as previously described (Sambrook et al., 1989). FK300 mutants were constructed via allelic replacement using pKFT (Kato and Sugai, 2011). Polymerase chain reaction (PCR) was performed using TaKaRa LA Taq (TaKaRa, Shiga, Japan). The thermal cycling conditions were as follows: 94°C for 2 min, followed by 30 cycles of 94°C for 15 s, 50°C, for 30 s, and 68°C for 2 min, finishing with a final extension step at 72°C for 1 min. The oligonucleotides used in this study are listed in Supplementary Table 2. Fragments were cloned into the plasmid pGEM-T Easy (Promega, Madison, WI, United States) using TA cloning and transformed into E. coli DH5a. A fragment excised from the pKS101 plasmid using a restriction enzyme was cloned into the pKFT plasmid and transformed into E. coli DH5a. Recombinant plasmids were introduced into S. aureus RN4220 via electroporation (Kraemer and Iandolo, 1990; Löfblom et al., 2007; Hisatsune et al., 2016).

The modified plasmids were then electroporated into *S. aureus* FK300 cells for allelic replacement. Marker-less deletion mutants in tetracycline-sensitive colonies were screened using PCR. Fragments were verified using DNA sequencing with a BigDye Terminator v 3.1 Cycle Sequencing Kit (Applied Biosystems, Waltham, MA, United States). In the complementation experiments, genes were amplified using PCR with the corresponding primer pairs and then cloned into the SmaI site of pKAT (Kato, 2004). The plasmids, pKS103 and pKS104, carrying *the icaR* and *icaB* coding regions of FK300, respectively, were constructed and transformed into the *S. aureus* strains listed in Supplementary Table 1 using electroporation. Inserts in all plasmid constructs were verified using PCR and DNA sequencing.

Biofilm assay

A biofilm assay using polystyrene plates was performed as described previously (Heilmann et al., 1996b) with a few modifications. Briefly, overnight cultures were diluted 1:100 in TSB. Ten microliters of this diluted solution was transferred in triplicate into flat-bottom, 96-well polystyrene plates (TrueLine; Nippon Genetics Co., Ltd., Japan) containing 100µl of TSB or TSB plus 1% glucose. Following incubation at 37°C for 20 h, the wells were gently washed thrice with 320µl of sterile phosphate-buffered saline (PBS; 137mM NaCl, 2.7mM KCl, 10mM Na2HPO4·12H2O, and 1.8 mM KH₂PO₄; pH 7.4), and the biofilm was stained with 1% crystal violet for 15 min. Unbound crystal violet was then removed by washing the plate in a container by immersing and agitating it gently 10 times in tap water and then drying it. Biofilm-bound crystal violet was solubilized in 150 µl of 33% glacial acetic acid at 25°C for 15 min. The extracts were diluted 10-fold, and then, absorbance at 590nm was measured using an Immuno-Mini NJ-2300 spectrophotometer (Nalgene Nunc International K. K., Tokyo, Japan). Each assay was performed in triplicate and repeated three times.

Autoaggregation assay

Bacteria from the plate were inoculated into 3 ml of TSB in a test tube (φ 15 × 150 mm), placed in a water bath, and shaken at 140 rpm at an angle of 45° for 6h. The culture was vortexed (Vortex-Genie 2; Scientific Industries, Inc.) for 10 s. Turbidity of the culture was visually monitored using video photography. For swapping experiments, the culture supernatant of interest was passed through a 0.2-µm filter, incubated with the cells of interest, and washed several times with 10 mM PBS; thereafter, the autoaggregation assay was carried out as described above.

RNA isolation and quantitative real-time reverse transcription-PCR (qRT-PCR)

Real-time PCR of *ica* operon in *S. aureus* has been previously described (Yu et al., 2017). Overnight *S. aureus* cultures were

316

317

318

319

320

321

322

323

324

325

326

327

328

329

330

331

332

333

334

335

336

337

338

339

340

341

342

343

344

345

346

347

348

349

350

351

352

353

354

355

356

357

358

359

360

361

362

363

364

365

366

367

368

diluted in TSB containing 1% glucose to an initial optical density (OD) of 0.02 at 660 nm and harvested after 6h of incubation with shaking at 37°C. Total RNA was isolated using a FastRNA Pro Blue kit (MP Biomedicals, Santa Ana, CA, United States) according to the manufacturer's instructions. DNA was extracted by treatment with RQ1 RNase-free DNase (Promega) at 37°C for 30 min. After DNase inactivation, PCR was performed to verify the absence of contaminating DNA. RNA was then reversetranscribed using a Transcriptor First-Strand cDNA Synthesis Kit (Roche, Mannheim, Germany). The resulting cDNA was diluted 10-fold with Tris-EDTA buffer (10mM Tris-HCl and 1mM EDTA; pH 8.0) and used as a template in the real-time PCR. qRT-PCR was performed using the SsoAdvanced Universal SYBR Green SuperMix (Bio-Rad, Hercules, CA, United States) and a CFX96 real-time PCR detection system (Bio-Rad). The thermal cycling conditions were as follows: 95°C for 1 min, followed by 40 cycles of 95°C for 15 s, 60°C (icaA), or 62°C (gyrB) for 15 s, and 72°C for 30 s. All PCR runs were performed in triplicate, and data were analyzed using the CFX Manager software (version 3.0; Bio-Rad) according to the manufacturer's instructions. The housekeeping gene, gyrase subunit B (gyrB), was used as a reference gene to normalize the expression level of the target gene in each reaction. The primers used for real-time PCR are listed in Supplementary Table 2. All samples were inspected 3 times and the data were analyzed using the $2^{-\Delta\Delta Ct}$ method (Livak and Schmittgen, 2001).

Electron microscopy

Electron microscopic observations were performed using TEM, as previously reported (Yamada et al., 1996). Bacterial cells were harvested, washed twice with 0.1 M PBS, and collected using centrifugation $(2,300 \times g, 15 \text{ min})$. For TEM, cells were fixed with 2.5% glutaraldehyde and 1% OsO₄. Samples were dehydrated using an ethanol series and embedded in New Spurr (Agar Scientific Ltd., United Kingdom). Ultrathin sections were cut with an ultramicrotome (ULTRACUTS, Leica, Tokyo, Japan) and examined with a JEOL JEM-2000 EXII electron microscope (JEOL Ltd., Tokyo, Japan) at 80 or 100 kV.

Element analysis

Elemental analysis of autoaggregates from the culture supernatant of FK300 Δ 5bp Δ Bm was performed using a scanning electron microscope connected to an energy dispersive X-ray spectroscope (Miniscope TM-3030, Hitachi High-Technologies Corporation, Tokyo, Japan).

Gel permeation HPLC

The culture supernatant was concentrated *via* centrifugation using an Amicon Ultra 15 ml filter (3 kDa cut off, Merck Millipore 369

370

371

372

373

374

375

376

377

378

379

380

381

382

383

384

385

386

387

388

389

390

391

392

393

394

395

396

397

398

399

400

401

402

403

404

405

406

407

408

409

410

411

412

413

414

415

416

417

418

419

420

421

422

Corporation, Darmstadt, Germany) for 40 min at 4,000 × g, and the concentrate was treated with trichloroacetic acid (final conc. 5%) to remove proteins by centrifugation for 15 min at 6,000 × g. Subsequently, the supernatant was subjected to gel permeation chromatography using Shim-pack Diol-300 (500 × 7.9 mm) with water as the mobile phase at a flow rate 0.5 ml/min. The effluent was fractionated every 2 ml from fraction 1–21, 11 min after sample injection. Samples necessary for hydrolysis were treated at 100°C for 2 h in the presence of 2 N HCl and rehydrated with distilled water after evaporation for the colorimetric assay.

Colorimetric determination of amino sugars

Fractions obtained by HPLC were analyzed for amino sugars using the Morgan-Elson assay (Enghofer and Kress, 1979). Samples (100μ l) were incubated with 20μ l acetone containing 1.5% acetic anhydride and 100μ l boric acid buffer (pH 9.0) at 95°C for 8 min and kept on ice. Then, the samples were mixed with 750 µl of p-(dimethylamino) benzaldehyde containing 12.5% HCl and 50 µl of 2-ethoxyethanol, and left to stand at room temperature for 15 min. OD at 545 nm was measured using a microplate reader (Varioskan LUX multimode reader, Thermo Fisher Scientific, Waltham, MA, United States).

PNAG dot blot

PNAG dot blotting was performed on each fraction obtained by fractionating the supernatant using gel filtration chromatography (Cramton et al., 1999). Alternatively, Bacteria from the plate were inoculated and incubated in 3 ml TSB medium at 37°C for 6h with shaking. The incubated bacteria were centrifuged (8,000 g×2 min), and the supernatant and bacteria were separated. To 1 ml of supernatant passed through a 0.2 µm sterile filter, 50 µl of Proteinase K (50 µl/ml; TaKaRa) was added and allowed to react at 55°C for 30 min, followed by incubation at 85°C for 30 min to inactivate the protease. The bacterial cells were washed once with PBS, resuspended in 100 µl of 0.5 M EDTA, and boiled at 100°C for 5 min. To 40 µl of the supernatant after centrifugation, 10 µl of Proteinase K (50 µl/ml; TaKaRa) was added and reacted at 55°C for 30 min, followed by incubation at 85°C for 30 min to inactivate the protease. The resulting bacterial surface samples and culture supernatant samples were serially diluted. The bacterial cell surface sample, supernatant sample or HPLC fraction was dropped onto a nitrocellulose membrane (Amersham Protran NC 0.45; General Electric Company, Connecticut, United States) and the membrane was immersed in a TBS-T [50 mM Tris-HCl (pH 8.0), 150 mM NaCl, and 0.05% Tween 20 (Sigma-Aldrich, Inc.)] solution containing 5% skim milk and allowed to react for 1 h to block the membranes. The membrane was then washed thrice with TBS-T for 15 min and incubated with rabbit anti- Poly-β-1,6-N-acetyl-D-glucosamine (PNAG) antiserum supplied by Dr. Gerald Pier (Skurnik et al., 2010)

diluted 1:4,000 with 0.5% skimmed milk in PBS-T for 2h at room temperature. Bound antibody was detected with peroxidase-conjugated goat anti-rabbit immunoglobulin G (IgG) antibodies (MP Biomedicals, LLC-Cappel Products, Ohio, United States; 1:6,000) and developed using Pierce enhanced chemiluminescence (ECL) western blotting substrate (Thermo Fisher Scientific, Rockford, IL, United States). Results were detected using a charge-coupled device camera-based imager (GE Healthcare Life Sciences, Piscataway, NJ, United States) or FujiRX-U film (Fujifilm, Tokyo, Japan), and developed using HI-RENDOL (Fujifilm) and HI-RENFIX (Fujifilm).

Electrospray ionization mass spectrometry (ESI-MS) analysis

MS analysis was performed using an LTQ Orbitrap XL mass spectrometer (Thermo Fisher Scientific) on a direct infusion MS using a nanospray capillary.

Samples in 50% methanol containing 0.1% formic acid were ionized using electrospray ionization in positive ion mode, and the following parameters were set: spray voltage, 1.5 kV; tube lens voltage, 250 V(max); source fragmentation, ON (100 Vmax); and mass acquisition range, m/z 200–4,000. Data were processed using the Xcalibur software package provided by Thermo Fisher Scientific.

Statistical analysis

Differences between two means were evaluated by the Mann– Whitney U test. Data were analyzed by Kruskal-Wallis test of variance to compare multiple means. Differences with *p*-values less than 0.05 were considered statistically significant. Statistical analysis was performed using R 4. 1.3 was used, with a significance level of 5%.

Results

Effects of *icaB* on cell autoaggregation and biofilm elaboration of *Staphylococcus aureus* FK300

The deletion of the 5-bp motif mutant from *S. aureus* wild type (WT) strain FK300 *via* allelic replacement resulted in the isolation of three independent mutants (Δ 5bpBm 1, 2, and 3) showing a common but unusual phenotype (Supplementary Figure 1). Most isolates upon 5-bp deletion mutation showed normal colony morphology, but a few revealed unusual phenotypes, as indicated by colonies showing shiny flat morphology, compounded by fusion between adjacent colonies (Figure 1, colony). In broth culture, most possible 5-bp motif-deletion mutants revealed a super-biofilm-producing phenotype, where these mutants

exhibited thick pellicles in test tube broth culture, as observed in a previous study (You et al., 2014; Figure 1, left, Δ 5bp, before vortex). By contrast, the three isolates in our study did not show any pellicles on the wall of the test tube (Figure 1, right, Δ 5bpBm 2, before vortex). Furthermore, weak mixing using a vortex mixer for a few seconds induced instant autoaggregation and subsequent sedimentation of the bacteria, resulting in a clear supernatant (Figure 1, right, Δ 5bpBm 2, after vortex), whereas the genuine mutant Δ 5bp remained turbid (Figure 1, left, Δ 5bp, after vortex; Supplementary Video 1).

Whole-genome sequencing of the three mutants established that their 5-bp motif was defective, indicating that the deletion of 5-bp had been successful. In addition, we identified a point mutation in the coding sequence of *icaB* in each strain (Supplementary Figure 2). For Δ 5bpBm 3, a frameshift mutation



Autoaggregation assay of *S. aureus* FK300 5bp deletion mutant culture. Autoaggregation assay of *S. aureus* FK300 5-bp deletion mutant culture. During the process of isolating the 5-bp (*icaR-icaA* intergenic region) deletion mutant from *S. aureus* FK300, we obtained colonies showing a distinct phenotype and the isolates were designated as Δ 5bpBm 1, 2, and 3. These isolates were cultured in TSB for 6h (before), and vortexed for 10 s (after), following which images of the culture were obtained (Supplementary Video 1). Shown is an image of the shape of the colonies growing on TSA plate. Left, genuine Δ 5bpBm 2. Other Δ 5bpBm mutants revealed a similar phenotype (Supplementary Figure 1).

10.3389/fmicb.2022.1101545

occurred due to the insertion of T, resulting in the creation of a termination codon 67 bases downstream, whereas Δ5bpBm 2 and Δ 5bpBm 1 demonstrated single and double missense mutations, respectively. Thereafter, FK300∆5bpBm 2 was used as the representative strain in the experiment. When FK300∆5bpBm 2 mutant was complemented with *icaB*, it formed adherent biofilms after culture and shown pellicle formation in vitro (Figure 2A). After voltexing, the formed biofilm was not completely dissolved as that of Δ 5bp probably due to overproduction of adherent biofilm and remained detached was suspended as clumps. Biofilm assays using plastic plates substantiated that FK300∆5bpBm 2 had completely lost the ability to form adherent biofilms, whereas complementation with *icaB* restored strong biofilm production (Figure 2B). These results suggested that *icaB* was involved in the unusual phenomenon displayed by the three FK300 Δ 5bpBm 2 mutant. Therefore, we generated FK300 Δ 5bp Δ *icaB* and verified that the mutant showed the same phenotype as FK300 Δ 5bpBm 2. This phenotype was also successfully complemented by *icaB*. To establish that this phenomenon was dependent on the ica operon, we overexpressed *icaR* in FK300 Δ 5bp Δ *icaB* to regulate the



By Marketon Feas and a strain expansion of lease distant production and 5-bp deletion cause autoaggregation. Autoaggregation assay (**A**) and biofilm assay (**B**) of various *Staphylococcus aureus* FK300 and the associated mutants: FK300Δ5bp; FK300Δ5bpBm 2; FK300Δ5bpΔ*icaB*; and FK300Δ5bpBm 2 complemented with pKAT carrying *icaR* (*picaR*), *icaB* (*picaB*), or pKAT, whereas FK300Δ5bpΔ*icaB* complemented with *picaB* or pKAT. Autoaggregation assay was carried out as shown in Figure 1. Biofilm assay was conducted using a microtiter plate. Bacteria were grown in TSB in the presence (Glc) or absence (Glc-) of 1% glucose. Biofilm stained with crystal violet was solubilized and OD 590 nm was measured using the polystyrene microtiter plate as described in the Methods section. Bars indicate mean values, error bars indicate standard error of the mean (n=3). WT, wild type strain FK300. pKAT, empty vector control. production of PNAG. FK300 Δ 5bpBm 2 p*icaR* completely lost the autoaggregating phenotype observed in FK300 Δ 5bp Δ *icaB*, indicating its dependence on the *ica* operon (Figure 2).

Deletion of the 5-bp motif is known to increase biofilm formation (Yu et al., 2017). However, other regulatory repressor regullators (rob and icaR) are also involved in biofilm formation, and the deletion of these factors has been shown to increase biofilm formation (Jefferson et al., 2004; Yu et al., 2017). We therefore created *icaB* deletion mutants and double deletion mutants combining the repressor and $\Delta i caB$ to see if it would develop a self-aggregating phenotype as observed with $\Delta 5 bp \Delta i ca B$. Although all three deletion mutants lost the adherent biofilm formation ability (Figure 3B), only FK300 Δ 5bp Δ icaB showed an autoaggregating phenotype (Figure 3A). This result was reproduced when the culture was shaken even 250 rpm (Unpublished data). Tiny autoaggregates were observed in the broth of FK300 Δ rob Δ icaB and FK300 Δ icaR Δ icaB, but there were no pellicles and the supernatant remained turbid even after mixing using a vortex mixer. This may likely have been due to the differences between the magnitudes of activation of the ica operon in the deletion mutants (Δrob , $\Delta icaR$, and $\Delta 5bp$), and we, therefore, measured mRNA levels of *ica* operon in the respective



Comparison of each biofilm formation inhibitory factor and *icaB* double deficient strains. Autoaggregation and biofilm formation of Δ 5bp Δ *icaB*, Δ *icaR\DeltaicaB*, and Δ *rob\DeltaicaB* double mutants. Autoaggregation assay (A) and biofilm formation (B) of FK300 and FK300 Δ 5bp, FK300 Δ 5bpBm 2, FK300 Δ 5bp Δ *icaB*, FK300 Δ *icaB*. Status before and after mixing for a few seconds on a vortex mixer is shown. Each bacterium was cultured in TSB medium at 37°C for 6 h with shaking. (B) Bacteria were grown in TSB in the presence (Glc) or absence (Glc-) of 1% glucose. Biofilm formation was measured using the polystyrene microtiter plate assay as described in the Methods section. Bars indicate mean values, error bars indicate standard error of the mean (*n*=3). WT; wild type strain FK300.

640

641

642 643

644

645

646

647

648

649

650

651 652

653

654

655

656

657

658

659

660 661

662

663

664

665

666

667

668

669

670

671

672

673

674

675

676

677

678

679

680

681

682

683

684

685

686

687

688

689

690

691

692

mutants. The results showed that expression of *ica* operon by Δ 5bp was much stronger than that by Δ *rob* or Δ *icaR* (Figure 4). The ica operon expression levels of each defective mutant compared to WT were about 31.46-fold for Δrob , 173.90-fold for $\Delta icaR$, and 2486.17-fold for $\Delta 5$ bp. This is consistent with the report of Yu et al. (2017). Based on these observations, we analyzed fractions of the culture supernatant and cell surface of each strain using anti-PNAG antiserum dot blot in order to investigate the actual production of PNAG. No signal was observed on the cell surfaces of FK300 Δ 5bp Δ *icaB*, FK300 Δ *ica*R Δ *icaB*, or FK300 Δ rob Δ *icaB*. However, in the culture supernatant, a clear strong signal demonstrating a reaction with the anti-PNAG antiserum was seen for FK300 Δ 5bp Δ *icaB* and a weak one for FK300 $\Delta icaR\Delta icaB$, but not for FK300 $\Delta rob\Delta icaB$ (Figures 5A,B). On the other hand, strong signals were detected in both the culture supernatant and the cell surface in FK300 Δrob , FK300 $\Delta i caR$, and FK300 $\Delta 5 bp$, respectively as expected. For FK300, a very weak signal was observed only at the cell surface, and no signal was detected in the culture supernatant. For FK300*\DeltaicaB*, no signal was observed in both culture supernatant and cell surface.

Substances that caused autoaggregation of cells

We conducted a swapping experiment to identify the factor(s) involved in autoaggregation on the cell surface or in the culture supernatant of FK300 Δ 5bp Δ *icaB*. The culture



FIGURE 4

Measurements of *icaA* transcription by qPCR. *Ica* operon expression of hyper-biofilm elaborating mutants. Total RNA preparation, cDNA synthesis, and quantitative PCR were performed as described in the Methods section. Relative expression level of *ica* operon; transcript levels in the Δ 5bp, *icaR*, or *rob* deletion mutants as compared to those in WT strain FK300 are shown. The expression of *gyrB* was used for sample normalization. Bars indicate mean values, error bars indicate standard error of the mean (*n*=3). Kruskal-Wallis test results *p*<0.05. There were significant differences between all deficient strains and WT. *: *p*<0.05 (Mann–Whitney U test). WT; wild type strain FK300. medium of each bacteria was separated into cells and culture supernatant, and a vortex experiment was performed by multiplying the selected supernatant with the selected cells to see which sample was important for aggregation. The strains used in the experiment were FK300, FK300∆5bp, or FK300 Δ 5bp Δ *icaB*. The autoaggregating factor(s) was present in the culture supernatant of FK300 Δ 5bp Δ *icaB*, but not in the culture supernatant of FK300∆5bp or on the cell surface of FK300 Δ 5bp Δ *icaB* (Figure 6). The results suggest that the important factor for aggregation is in the culture supernatant of FK300 Δ 5bp Δ *icaB*. We examined the ultrastructure of the autoaggregates using transmission electron microscopy (TEM). The autoaggregate preparation obtained from sedimented FK300∆5bpBm 2 was compared with a cell pellet of WT FK300. There was no significant difference between the ultrastructures of the cell wall peptidoglycan layers of the strains (Figure 7A). However, string-like substances surrounding FK300∆5bpBm 2 were observed, implying the presence of string-like substances in the culture supernatant of FK300∆5bpBm 2. We further analyzed the elements of the autoaggregates obtained from the culture supernatant of FK300Δ5bpΔicaB using a TM-3030 Hitachi microscope connected to energy dispersive X-ray spectroscope. Dried autoaggregate was spotted using a miniscope, and the spotted site was analyzed. The results suggested that the ratios of carbon, oxygen, and nitrogen were similar to those of N-acetylglucosamine (Figure 7B). To gain further insights into the biochemical structure of the string-like substances surrounding the mutant, we prepared a culture supernatant of FK300 Δ 5bp Δ *icaB*, which was then vigorously shaken to determine whether any autoaggregation occurred without bacterial cells and also whether any tiny but distinct whitish autoaggregates would appear after shaking (Supplementary Figure 3). Such substances appeared in FK300 Δ 5bp Δ *icaB* culture supernatants but not in FK300 and FK300Δ5bp culture supernatants. We studied in vitro autoaggregation in the culture supernatant of FK Δ 5bp Δ *icaB*, to analyze the inhibitory effect of several additives, including NaCl, ethylenediaminetetraacetic acid (EDTA), dispersin B, proteinase K, and DNase. Only dispersin B, a β-hexosaminidase that specifically hydrolyzes β-1,6-glycosidic bonds in acetylglucosamine polymers (Kaplan et al., 2003; Ramasubbu et al., 2005; Itoh et al., 2005), inhibited the appearance of aggregates in vitro. This finding strongly suggested that acetylglucosamine polymers were the major factor(s) involved in the autoaggregation of FK300 Δ 5bp Δ *icaB*.

To further analyze the biochemistry of these polymers, the FK300 Δ 5bp Δ *icaB* culture supernatant was fractionated using gel filtration chromatography, and each fraction with or without hydrolysis was analyzed for amino sugar concentration. A broad peak appeared in the void-volume-column (fractions 3–8) of the HCl-hydrolyzed fractions, indicating that the polymers in the supernatant of FK300 Δ 5bp Δ *icaB* contained amino sugars (Figure 8A). Furthermore, dot blot analysis with antisera against

746

693



FIGURE 5

PIA / PNAG production on the culture supernatant and cell surface in each strain. PIA/PNAG production in the culture supernatant and on cell surface of the mutants. PIA/PNAG collected from the culture supernatant and cell surface of FK300 and FK300\[Delta]5bp\]icaB, FK300\[Delta]caB, FK300\[Delta]ca



FIGURE 6

Swapping experiment of culture supernatant and bacterial cells. Swapping of culture supernatant and bacterial cells. Filtered culture supernatant of FK300, FK300 Δ 5bp Δ icaB was incubated with bacterial cells of FK300 or FK300 Δ 5bp Δ icaB (before) and vortexed for 10 s (after), and photographic images of the culture were obtained. Aggregates identified after vortexing were indicated by red arrows. WT; wild type strain FK300.

PNAG clearly demonstrated that the corresponding fractions were positive for PNAG (Figure 8B). Collectively, these results strongly suggested that the factor(s) involved in the autoaggregation of FK300 Δ 5bp Δ *icaB* in the supernatant was an N-acetylglucosamine polymer.

N-acetylglucosamine polymers secreted from the cytosol of *S. aureus* undergo partial deacetylation by IcaB, which functions as a deacetylase (Vuong et al., 2004). The resulting extracellular positively charged deacetylated polymers interact with the

negatively charged cell surface to form a biofilm matrix around the cells. FK300 Δ 5bp Δ *icaB* lacks IcaB deacetylase and is therefore unable to catalyze the deacetylation of *N*-acetylglucosamine polymers, which releases the polymer into the culture supernatant. Therefore, we investigated whether the polymers involved in the autoaggregation of FK300 Δ 5bp Δ *icaB* were deacetylated, by performing mass spectrometry (MS) analysis of the polymers. PNAG on the surface layer of the biofilm-producing strain, FK300 Δ *rob*, was extracted and used as в



INTH NOVE	
a trans	2 VA
NO YA GAL	
A Cit	- Alera
No Right	25-1
	1
Stant -	- Human
Aspping	50 µm

Element	Carbon	Oxygen	Nitrogen	Total
Norm. C [wt.%]	46.6	46.6	6.8	100.0
Atom. C [at.%]	53.3	40.0	6.7	100.0
Δ5bpΔ <i>icaB</i> Aggregates of culture supernatant				
Element	Carbon	Oxygen	Nitrogen	Total
Element Norm. C [wt.%]	Carbon 46.0	Oxygen 43.7	Nitrogen 10.3	Total 100.0
Element Norm. C [wt.%] Atom. C [at.%]	Carbon 46.0 52.6	Oxygen 43.7 37.4	Nitrogen 10.3 10.0	Total 100.0 100.0

N-acetyl-glucosamine [positive control]

FIGURE 7

Searching for autoaggregation factors using electron microscopy and elemental analysis. Morphological and elemental analysis of autoaggregation factor(s) using electron microscopy. (A) Ultrastructural observation of *Staphylococcus aureus* FK300\Delta5bpBm 2 using transmission electron microscope. *S. aureus* FK300 (left) and *S. aureus* FK300\Delta5bpBm 2 (right). (B) Elemental analysis of the autoaggregate from FK300\Delta5bpΔ*icaB* supernatant. Autoaggregates obtained by vortexing the culture supernatant of FK300Δ5bpΔ*icaB* were dried and examined *via* a scanning electron microscope (Mini Scope TM-3030) for element analysis. As a control, the value of *N*-acetylglucosamine is shown. WT; wild type strain FK300. The top left figure is SEM image of a sample irradiated with characteristic X-rays. The bottom left figure is EDS spectrum. The right is a table of elemental ratios of flocculence samples and *N*-acetylglucosamine.

a control. MS analysis of PNAG gathered from the surface of FK300 Δ *rob* cells showed MS signals with a regular interval of m/z=203, which corresponded to the *N*-acetylglucosamine (GlcNAc) molecule. In addition, several signals corresponding to partially deacetylated polymers, such as (GlcN)₁-(GlcNAc)₆, (GlcN)₂-(GlcNAc)₅, and (GlcN)₃-(GlcNAc)₄ between (GlcNAc)₆ and (GlcNAc)₇, were observed (Figure 9A). By contrast, polymers recovered from the culture supernatant of FK300 Δ 5bp Δ *icaB* showed signals corresponding to (GlcNAc)_n, but not those corresponding to deacetylated fragments (Figure 9B). These results clearly showed that the polymers in the supernatant of FK300 Δ 5bp Δ *icaB* were polymers of *N*-acetylglucosamine without deacetylation.

Discussion

IcaB plays a crucial role in the biogenesis of extracellular *N*-acetylglucosamine polymers for adherent biofilm formation, by deacetylating *N*-acetylglucosamine polymers, which makes them electrostatically positive. These polymers then interact with the negatively charged bacterial cell surface *via* electrostatic interactions leading to PIA in *S. epidermidis* (Mack et al., 1996). Deletion of 5-bp in the icaR-icaA intergenic region leads to strong biofilm formation in *S. aureus* (Yu et al., 2017). Defective IcaB functioning leads to the production of fully acetylated *N*-acetylglucosamine polymers, which no longer interact with the cell surface and remain in the culture supernatant. Therefore, we assumed that FK300 Δ 5bp Δ *icaB* produces many fully acetylated poly-*N*-acetylglucosamine molecules and secretes them into the culture supernatant. Our study clearly demonstrated that the secretion of a large amount of poly-*N*-acetylglucosamine molecules without deacetylation played a direct role in the autoaggregation of *S. aureus* after vortexing.

Similar to the vortex-induced autoaggregation of *S. aureus* in this study, there was a report of vortex-induced aggregation in single-walled carbon nanotubes (Fernandes et al., 2017). Although the detailed physical mechanism is not known, the aggregates grew larger with increasing addition up to a certain concentration for single-walled carbon nanotubes, and then leveled off. This is similar to the present experimental results.

Kutsuno et al



FIGURE 8

Gel permeation high performance liquid chromatography analyses of culture supernatants of FK300 Δ 5bp Δ icaB and WT. Gel permeation high performance liquid chromatography analyses of culture supernatants of FK300 Δ 5bp Δ icaB and WT. Concentrated culture supernatants of FK300 Δ 5bp Δ icaB and WT. Concentrated culture supernatants of FK300 Δ 5bp Δ icaB and FK300 were subjected to gel permeation chromatography (GPC) using Shim-pack Diol-300 (500×7.9 mm) under water as the mobile phase with a flow rate 0.5 ml/min. Effluents were fractionated every 2 ml from fraction (Fr.) 1 to 21, 11 min after sample injection. Each fraction, with or without hydrolysis, was analyzed for amino sugar (A) using the Morgan-Elson colorimetric method and the PNAG (B) signal using dot blot with the rabbit anti-PNAG as described in the Methods section. WT; wild type strain FK300. Previous studies of carbon nanotubes suggested that aggregation was caused by the interaction of single-walled carbon nanotubes with each other due to the weakening of the shielding effect that prevented the tubes from aggregating. From this, it was inferred that the aggregation in this study was caused by the loss of positive charge due to deacetylation of PNAG caused by the dysfunction of *icaB*, which facilitated interaction of PNAG with each other.

In this study, Allelic exchange to create a 5-bp deficient strain confirmed the presence of an unselected secondary mutation in *icaB*. Three of the mutants with altered colony morphology each had a different mutation in *icaB*, and these were not clonal. This suggested that the 5-bp deletion mutation alone had a fitness cost under the conditions employed in mutagenesis. In Figure 3, clearly aberrant autoaggregation was observed in FK300 Δ 5bp Δ *icaB*, but not in FK300 Δ *rob\DeltaicaB* and FK300 $\Delta i ca R \Delta i ca B$. We analyzed the mRNA expression of icaA in the three double mutants and found that ica operon expression was considerably high in the FK300 Δ 5bp Δ icaB mutants (Figure 4). In agitation agglutination experiments with FK300 $\Delta i ca R \Delta i ca B$ and FK300 $\Delta rob \Delta i ca B$ strains, the supernatant did not become clear, but small clumps were visible, suggesting that high expression of the *ica* operon is required to induce agglutination like FK300 Δ 5bo Δ *icaB* in this study. Deletion of the 5-bp motif induced maximum expression



FIGURE 9

The typical ESI positive ion mass spectrum of PIA fractionated by GPC, showing the major mass differences to be deacetylation of poly-N-acetyl-D-glucosamine polymer. Typical ESI-positive ion mass spectrum of PNAG from FK300 Δ 5bp Δ *icaB* fractionated using GPC. Samples of cell surface FK300 Δ rob (A) and culture supernatant FK300 Δ 5bp Δ *icaB* (B) from the collected fraction by GPC were analyzed using ESI-MS. MS analyses of authentic PNAG (A) showed fragmentation of polysaccharide homopolymers resulting in a series of ESI-MS of m/z=203Da, which corresponded to GlcNAc. Besides these spectra, several peaks defecting of m/z=42Da, which corresponded to deacetylation of [GlcNAc] polymer were observed in FK300 Δ rob (A). By contrast, the culture supernatant of autoaggregation inducing strain FK300 Δ 5bp Δ *icaB* (B) contained non deacetylated [GlcNAc] polymer. *N*-acetyl-D-glucosamine is expressed as GlcNAC, Glucosamine as GlcN. WT; wild type strain FK300.

of the *ica* operon. A previous study demonstrated that *rob* binds to the 5-bp motif (Yu et al., 2017). However, the magnitude of *icaA* expression in KF300 Δ 5bp Δ *icaB* was far greater than that in FK300 Δ rob Δ *icaB*. This suggested that factor(s) other than *rob* were involved in 5-bp motif dependent *icaA* activation. The identity of these factor(s) remains to be explored.

Considering massive production of PNAG without deacetylation in culture supernatant of FK300 Δ 5bp Δ *icaB*, the signal in the culture supernatant in dot blot assay was weak (Figure 5A). This may be attributed to structural differences between fully acetylated poly-*N*-acetylglucosamine and PNAG used to generate antiserum. It remains unclear as to what extent the fully acetylated poly-*N*-acetylglucosamine reacted with the anti-PNAG serum used in this study. Therefore, it was not possible to ascertain quantitative differences between strains with and without *icaB* deletion *via* this comparison.

Biofilms are defined as adherent communities of microbial origin, represented by cells that adhere to substrates, interfaces, or each other, are embedded in a matrix of extracellular polymeric material, and exhibit an altered phenotype concerning growth, gene expression, and protein production. Cell aggregates formed in the absence of a surface (Trunk et al., 2018) and floating pellicles that form biofilms at the air-liquid interface (Vaccari et al., 2017) are also considered to be a type of biofilm. In this study, we found that strains with mutations in *icaB* caused the self-aggregation and settling as clumps by vortexing. This suggests that deacetylation of PNAG, a major component of staphylococcal biofilms, by *icaB* is necessary for biofilm adherence, but we have demonstrated that accumulation non-deacetylated PNAG formed huge of microbial aggregates as non-adherent biofilm. Non-stick biofilms have also been reported in previous papers with Escherichia coli (Schembri et al., 2003) and Clostridium perfringens (Obana et al., 2020).

Staphylococci are known to be the causative agent of endocarditis, and a high incidence of thromboembolism in mechanical heart valves, where non-physiological flow patterns cause platelet aggregation and free thrombus. It has been previously reported that mechanical valves cause eddy-like turbulence that carries thrombus downstream (Bluestein et al., 2000), and autoaggregation-causing strains such as *S. aureus* in this study may increase the risk of thrombus formation in patients implanted with mechanical valves. Autoaggregating mechanism of *S. aureus* demonstrated in this study may have implication to explain the phenomena of autoaggregation of *S.aureus* observed in chronic wounds (Kirketerp-Møller et al., 2008) and/or synovial fluid (Staats et al., 2021). In such a way, non-adherent biofilms of *S. aureus* could be potentially involved in pathogenicity and need further investigation.

In summary, we discovered abnormal autoaggregation of *S. aureus* mutants lacking both TATTT motif in the intergenic region between *icaR* and *ica* operon, and *icaB* function upon

vortexing. This was illustrated by the production of a large amount of fully acetylated PNAG polymer in the culture supernatant, which induced autoaggregation by tiny physical stimuli and formed clumps. We propose this as an another type of biofilm, non-adherent biofilm.

Genome sequencing

Genomic DNA extraction and whole-genome sequencing were performed as described previously27. The raw data reads of the 5-bp (*icaR-icaA* intergenic region) deletion mutants from S. aureus FK300, FK300 Δ 5bpBm 1, 2, and 3, have been deposited in the DDBJ/Sequence Read Archive under the accession numbers DRA013849.

Data availability statement

The datasets presented in this study can be found in online repositories. The names of the repository/repositories and accession number(s) can be found at: https://www.ddbj.nig.ac.jp/, DRA013849.

Author contributions

LY, JH, and MS conceived and designed the experimental studies. SK performed most experiments described in the paper and wrote the paper. JH analyzed the sequence data and supported experimental design and technique. LY provided assistance with the RNA isolation and qRT-PCR. SY provided assistance with the electron microscopy. IH provided assistance with Element analysis, Gel permeation HPLC and ESI-MS analysis. MS helped with writing and editing. All authors contributed to the article and approved the submitted version.

Funding

The work was supported by Research Program on Emerging and Re-emerging infectious Diseases from the Japan Agency for Medical Research and Development (AMED) under grant number JP21fk0108604j0001, JP20gm1010001j0305, the Health and Labor Sciences Research Grant (21HA2009) and JSPS KAKENHI Grant Number JP20K10268.

Acknowledgments

We thank Gerald Pier, Brigham and Women's Hospital Channing Labs for anti-PNAG serum.

```
1125
       1126
       1127
       1128
       1129
       1130
       1131
       1132
       1133
       1134
       1135
       1136
       1137
       1138
       1139
       1140
       1141
      1142
Q14
       1143
       1144
       1145
       1146
       1147
       1148
       1149
      1150
       1151
       1152
       1153
       1154
       1155
       1156
       1157
       1158
       1159
       1160
       1161
       1162
       1163
      1164
Q16
       1165
       1166
       1167
       1168
       1169
       1170
       1171
```

1179 **Q17**

1180

1181

1182

1183

1184

1185

1186

1187

1188

1189

1190

1191

1192

1193

1194

1195

1196

1197

1198

1199

1200

1201

1202

1203

1204

1205

1206

1207

1208

1209

1210

1211

1212

1213

1214

1215

1216

1217

1218

1219

1220

1221

1222

1223

1224

1225

1226

1227

1228

1229

1230

1231

1232

Conflict of interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Publisher's note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated

References

Archer, N. K., Mazaitis, M. J., Costerton, J. W., Leid, J. G., Powers, M. E., and Shirtliff, M. E. (2011). Staphylococcus aureus biofilms: properties, regulation, and roles in human disease. *Virulence* 2, 445–459. doi: 10.4161/viru.2.5.17724

Barrett, L., and Atkins, B. (2014). The clinical presentation of prosthetic joint infection. J. Antimicrob. Chemother. 69, i25–i27. doi: 10.1093/jac/dku250

Bluestein, D., Rambod, E., and Gharib, M. (2000). Vortex shedding as a mechanism for free emboli formation in mechanical heart valves. *J. Biomech. Eng.* 122, 125–134. doi: 10.1115/1.429634

Byrd, A. L., Belkaid, Y., and Segre, J. A. (2018). The human skin microbiome. *Nat. Rev. Microbiol.* 16, 143–155. doi: 10.1038/nrmicro.2017.157

Chatterjee, S., Maiti, P., Dey, R., Kundu, A., and Dey, R. (2014). Biofilms on indwelling urologic devices: microbes and antimicrobial management prospect. *Ann. Med. Health Sci. Res.* 4, 100–104. doi: 10.4103/2141-9248.126612

Chen, K. M., Chiang, M. K., Wang, M., Ho, H. C., Lu, M. C., and Lai, Y. C. (2014). The role of pgaC in *Klebsiella pneumoniae* virulence and biofilm formation. *Microb. Pathog.* 77, 89–99. doi: 10.1016/j.micpath.2014.11.005

Choi, A. H., Slamti, L., Avci, F. Y., Pier, G. B., and Maira-Litrán, T. (2009). The pgaABCD locus of *Acinetobacter baumannii* encodes the production of polybeta-1-6-*N*-acetylglucosamine, which is critical for biofilm formation. *J. Bacteriol.* 191, 5953–5963. doi: 10.1128/JB.00647-09

Claassen-Weitz, S., Shittu, A. O., Ngwarai, M. R., Thabane, L., Nicol, M. P., and Kaba, M. (2016). Fecal carriage of Staphylococcus aureus in the hospital and community setting: a systematic review. *Front. Microbiol.* 7:449. doi: 10.3389/fmicb.2016.00449

Conlon, K. M., Humphreys, H., and O'Gara, J. P. (2002). *icaR* encodes a transcriptional repressor involved in environmental regulation of *Ica* operon expression and biofilm formation in *Staphylococcus epidermidis*. *J. Bacteriol.* 184, 4400–4408. doi: 10.1128/JB.184.16.4400-4408.2002

Costerton, J. W., Geesey, G. G., and Cheng, K. J. (1978). How bacteria stick. Sci. Am. 238, 86–95. doi: 10.1038/scientificamerican0178-86

Cramton, S. E., Gerke, C., Schnell, N. F., Nichols, W. W., and Götz, F. (1999). The intercellular adhesion (*Ica*) locus is present in *Staphylococcus aureus* and is required for biofilm formation. *Infect. Immun.* 67, 5427–5433. doi: 10.1128/ IAI.67.10.5427-5433.1999

Cue, D., Lei, M. G., and Lee, C. Y. (2012). Genetic regulation of the intercellular adhesion locus in staphylococci. *Front. Cell. Infect. Microbiol.* 2:38. doi: 10.3389/ fcimb.2012.00038

Di Domenico, E. G., Oliva, A., and Guembe, M. (2022). The current knowledge on the pathogenesis of tissue and medical device-related biofilm infections. *Microorganisms* 10:1259. doi: 10.3390/microorganisms10071259

Enghofer, E., and Kress, H. (1979). An evaluation of the Morgan--Elson assay for 2-amino-2-deoxy sugars. *Carbohydr. Res.* 76, 233–238. doi: 10.1016/0008-6215(79)80022-1

Fernandes, R., Buzaglo, M., Regev, O., Furó, I., and Marques, E. F. (2017). Mechanical agitation induces counterintuitive aggregation of pre-dispersed carbon nanotubes. J. Colloid Interface Sci. 493, 398–404. doi: 10.1016/j.jcis.2017.01.040

Flemming, H. C., and Wingender, J. (2010). The biofilm matrix. Nat. Rev. Microbiol. 8, 623-633. doi: 10.1038/nrmicro2415

Foster, T. J., Geoghegan, J. A., Ganesh, V. K., and Höök, M. (2014). Adhesion, invasion and evasion: the many functions of the surface proteins of *Staphylococcus aureus*. *Nat. Rev. Microbiol.* 12, 49–62. doi: 10.1038/nrmicro3161

Gerke, C., Kraft, A., Süssmuth, R., Schweitzer, O., and Götz, F. (1998). Characterization of the N-acetylglucosaminyltransferase activity involved in the organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

Supplementary material

The Supplementary material for this article can be found online at: https://www.frontiersin.org/articles/10.3389/fmicb.2022.1101545/ full#supplementary-material

biosynthesis of the *Staphylococcus epidermidis* polysaccharide intercellular adhesin. J. Biol. Chem. 273, 18586–18593. doi: 10.1074/jbc.273.29.18586

Giachino, P., Engelmann, S., and Bischoff, M. (2001). Sigma(B) activity depends on RsbU in *staphylococcus aureus*. J. Bacteriol. 183, 1843–1852. doi: 10.1128/ JB.183.6.1843-1852.2001

Heilmann, C., Gerke, C., Perdreau-Remington, F., and Götz, F. (1996b). Characterization of Tn917 insertion mutants of *Staphylococcus epidermidis* affected in biofilm formation. *Infect. Immun.* 64, 277–282. doi: 10.1128/iai.64.1.277-282.1996

Heilmann, C., Schweitzer, O., Gerke, C., Vanittanakom, N., Mack, D., and Götz, F. (1996a). Molecular basis of intercellular adhesion in the biofilm-forming *Staphylococcus epidermidis. Mol. Microbiol.* 20, 1083–1091. doi: 10.1111/j.1365-2958.1996.tb02548.x

Herbert, S., Ziebandt, A. K., Ohlsen, K., Schäfer, T., Hecker, M., Albrecht, D., et al. (2010). Repair of global regulators in *Staphylococcus aureus* 8325 and comparative analysis with other clinical isolates. *Infect. Immun.* 78, 2877–2889. doi: 10.1128/IAI.00088-10

Hisatsune, J., Sato'o, Y., Yu, L., Kutsuno, S., Hayakawa, Y., and Sugai, M. (2016). Efficient transformation of *Staphylococcus aureus* using multi-pulse electroporation. *J. Microbiol. Methods* 130, 69–72. doi: 10.1016/j.mimet.2016.08.012

Ibáñez de Aldecoa, A. L., Zafra, O., and González-Pastor, J. E. (2017). Mechanisms and regulation of extracellular DNA release and its biological roles in microbial communities. *Front. Microbiol.* 8:1390. doi: 10.3389/fmicb.2017.01390

Itoh, Y., Wang, X., Hinnebusch, B. J., Preston, J. F. 3rd, and Romeo, T. (2005). Depolymerization of beta-1,6-*N*-acetyl-D-glucosamine disrupts the integrity of diverse bacterial biofilms. *J. Bacteriol.* 187, 382–387. doi: 10.1128/ JB.187.1.382-387.2005

Jefferson, K. K., Cramton, S. E., Götz, F., and Pier, G. B. (2003). Identification of a 5-nucleotide sequence that controls expression of the Ica locus in *Staphylococcus aureus* and characterization of the DNA-binding properties of IcaR. *Mol. Microbiol.* 48, 889–899. doi: 10.1046/j.1365-2958.2003.03482.x

Jefferson, K. K., Pier, D. B., Goldmann, D. A., and Pier, G. B. (2004). The teicoplanin-associated locus regulator (TcaR) and the intercellular adhesin locus regulator (IcaR) are transcriptional inhibitors of the Ica locus in *Staphylococcus aureus*, *J. Bacteriol.* 186, 2449–2456. doi: 10.1128/JB.186.8.2449-2456.2004

Kaplan, J. B., Ragunath, C., Ramasubbu, N., and Fine, D. H. (2003). Detachment of Actinobacillus actinomycetemcomitans biofilm cells by an endogenous betahexosaminidase activity. J. Bacteriol. 185, 4693–4698. doi: 10.1128/ IB.185.16.4693-4698.2003

Kato, F. (2004). Induction of staphylococcal bicomponent toxins; exploration of inducers and their functions. [dissertation/master's thesis]. [Sendai City, Japan]: University of Tohoku.

Kato, F., and Sugai, M. (2011). A simple method of markerless gene deletion in *Staphylococcus aureus*. J. Microbiol. Methods 87, 76–81. doi: 10.1016/j. mimet.2011.07.010

Kiedrowski, M. R., and Horswill, A. R. (2011). New approaches for treating staphylococcal biofilm infections. *Ann. N. Y. Acad. Sci.* 1241, 104–121. doi: 10.1111/j.1749-6632.2011.06281.x

Kirketerp-Møller, K., Jensen, P. Ø., Fazli, M., Madsen, K. G., Pedersen, J., Moser, C., et al. (2008). Distribution, organization, and ecology of bacteria in chronic wounds. *J. Clin. Microbiol.* 46, 2717–2722. doi: 10.1128/JCM.00501-08

Kraemer, G. R., and Iandolo, J. J. (1990). High-frequency transformation of *Staphylococcus aureus* by electroporation. *Curr. Microbiol.* 21, 373–376. doi: 10.1007/BF02199440

Kreiswirth, B. N., Löfdahl, S., Betley, M. J., O'Reilly, M., Schlievert, P. M., Bergdoll, M. S., et al. (1983). The toxic shock syndrome exotoxin structural gene is not detectably transmitted by a prophage. *Nature* 305, 709–712. doi: 10.1038/305709a0

Le, K. Y., Park, M. D., and Otto, M. (2018). Immune evasion mechanisms of *Staphylococcus epidermidis* biofilm infection. *Front. Microbiol.* 9:359. doi: 10.3389/ fmicb.2018.00359

Li, D., Li, J., Zhao, F., Wang, G., Qin, Q., and Hao, Y. (2016). The influence of fermentation condition on production and molecular mass of EPS produced by *Streptococcus thermophilus* 05-34 in milk-based medium. *Food Chem.* 197, 367–372. doi: 10.1016/j.foodchem.2015.10.129

Lister, J. L., and Horswill, A. R. (2014). *Staphylococcus aureus* biofilms: recent developments in biofilm dispersal. *Front. Cell. Infect. Microbiol.* 4:178. doi: 10.3389/fcimb.2014.00178

Livak, K. J., and Schmittgen, T. D. (2001). Analysis of relative gene expression data using real-time quantitative PCR and the 2(-Delta Delta C(T)). *Method.* 25, 402–408. doi: 10.1006/meth.2001.1262

Löfblom, J., Kronqvist, N., Uhlén, M., Ståhl, S., and Wernérus, H. (2007). Optimization of electroporation-mediated transformation: *Staphylococcus carnosus* as model organism. *J. Appl. Microbiol.* 102, 736–747. doi: 10.1111/j.1365-2672.2006.03127.x

Lowy, F. D. (1998). Staphylococcus aureus infections. N. Engl. J. Med. 339, 520–532. doi: 10.1056/NEJM199808203390806

Mack, D., Fischer, W., Krokotsch, A., Leopold, K., Hartmann, R., Egge, H., et al. (1996). The intercellular adhesin involved in biofilm accumulation of *Staphylococcus epidermidis* is a linear beta-1,6-linked glucosaminoglycan: purification and structural analysis. *J. Bacteriol.* 178, 175–183. doi: 10.1128/jb.178.1.175-183.1996

Mayer, C., Moritz, R., Kirschner, C., Borchard, W., Maibaum, R., Wingender, J., et al. (1999). The role of intermolecular interactions: studies on model systems for bacterial biofilms. *Int. J. Biol. Macromol.* 26, 3–16. doi: 10.1016/ s0141-8130(99)00057-4

Mehraj, J., Witte, W., Akmatov, M. K., Layer, F., Werner, G., and Krause, G. (2016). Epidemiology of Staphylococcus aureus nasal carriage patterns in the community. *Curr. Top. Microbiol. Immunol.* 398, 55–87. doi: 10.1007/82_2016_497

Montanaro, L., Poggi, A., Visai, L., Ravaioli, S., Campoccia, D., Speziale, P., et al. (2011). Extracellular DNA in biofilms. *Int. J. Artif. Organs* 34, 824–831. doi: 10.5301/ ijao.5000051

Novick, R. (1967). Properties of a cryptic high-frequency transducing phage in *Staphylococcus aureus*. *Virology* 33, 155–166. doi: 10.1016/0042-6822(67)90105-5

Obana, N., Nakamura, K., and Nomura, N. (2020). Temperature-regulated heterogeneous extracellular matrix gene expression defines biofilm morphology in *Clostridium perfringens. NPJ biofilms and microbiomes* 6:29. doi: 10.1038/ s41522-020-00139-7

Otto, M. (2008). Staphylococcal biofilms. Curr. Top. Microbiol. Immunol. 322, 207–228. doi: 10.1007/978-3-540-75418-3_10

Parsek, M. R., and Singh, P. K. (2003). Bacterial biofilms: an emerging link to disease pathogenesis. *Annu. Rev. Microbiol.* 57, 677–701. doi: 10.1146/annurev. micro.57.030502.090720

Pokrovskaya, V., Poloczek, J., Little, D. J., Griffiths, H., Howell, P. L., and Nitz, M. (2013). Functional characterization of *Staphylococcus epidermidis* IcaB, a de-*N*-acetylase important for biofilm formation. *Biochemistry* 52, 5463–5471. doi: 10.1021/bi400836g

Ramasubbu, N., Thomas, L. M., Ragunath, C., and Kaplan, J. B. (2005). Structural analysis of dispersin B, a biofilm-releasing glycoside hydrolase from the periodontopathogen *Actinobacillus actinomycetemcomitans. J. Mol. Biol.* 349, 475–486. doi: 10.1016/j.jmb.2005.03.082

Sambrook, J., Fritsch, E. F., and Maniatis, T. (1989). *Molecular cloning: A laboratory manual*. Plainview, NY: Cold Spring Harbor Laboratory Press.

Schembri, M. A., Hjerrild, L., Gjermansen, M., and Klemm, P. (2003). Differential expression of the *Escherichia coli* autoaggregation factor antigen 43. *J. Bacteriol.* 185, 2236–2242. doi: 10.1128/JB.185.7.2236-2242.2003

Schilcher, K., and Horswill, A. R. (2020). Staphylococcal biofilm development: structure, regulation, and treatment strategies. *Microbiology and molecular biology reviews: MMBR* 84, e00026–e00019. doi: 10.1128/MMBR.00026-19

Skurnik, D., Merighi, M., Grout, M., Gadjeva, M., Maira-Litran, T., Ericsson, M., et al. (2010). Animal and human antibodies to distinct *Staphylococcus aureus* antigens mutually neutralize opsonic killing and protection in mice. *J. Clin. Invest.* 120, 3220–3233. doi: 10.1172/JCI42748

Staats, A., Burback, P. W., Eltobgy, M., Parker, D. M., Amer, A. O., Wozniak, D. J., et al. (2021). Synovial fluid-induced aggregation occurs across Staphylococcus aureus clinical isolates and is mechanistically independent of attached biofilm formation. *Microbiology spectrum* 9:e0026721. doi: 10.1128/Spectrum.00267-21

Trunk, T., Khalil, H. S., and Leo, J. C. (2018). Bacterial autoaggregation. *AIMS microbiology* 4, 140–164. doi: 10.3934/microbiol.2018.1.140

Vaccari, L., Molaei, M., Niepa, T., Lee, D., Leheny, R. L., and Stebe, K. J. (2017). Films of bacteria at interfaces. *Adv. Colloid Interf. Sci.* 247, 561–572. doi: 10.1016/j. cis.2017.07.016

Vuong, C., Kocianova, S., Voyich, J. M., Yao, Y., Fischer, E. R., DeLeo, F. R., et al. (2004). A crucial role for exopolysaccharide modification in bacterial biofilm formation, immune evasion, and virulence. *J. Biol. Chem.* 279, 54881–54886. doi: 10.1074/jbc.M411374200

Wang, X., Preston, J. F. 3rd, and Romeo, T. (2004). The pgaABCD locus of *Escherichia coli* promotes the synthesis of a polysaccharide adhesin required for biofilm formation. *J. Bacteriol.* 186, 2724–2734. doi: 10.1128/JB.186.9.2724-2734.2004

Yamada, S., Sugai, M., Komatsuzawa, H., Nakashima, S., Oshida, T., Matsumoto, A., et al. (1996). An autolysin ring associated with cell separation of *Staphylococcus aureus*. *J. Bacteriol.* 178, 1565–1571. doi: 10.1128/ jb.178.6.1565-1571.1996

You, Y., Xue, T., Cao, L., Zhao, L., Sun, H., and Sun, B. (2014). *Staphylococcus aureus* glucose-induced biofilm accessory proteins, GbaAB, influence biofilm formation in a PIA-dependent manner. *International journal of medical microbiology*: *IJMM* 304, 603–612. doi: 10.1016/j.ijmm.2014.04.003

Yu, L., Hisatsune, J., Hayashi, I., Tatsukawa, N., Sato'o, Y., Mizumachi, E., et al. (2017). A novel repressor of the Ica locus discovered in clinically isolated superbiofilm-elaborating *Staphylococcus aureus*. *MBio* 8, e02282–e02216. doi: 10.1128/ mBio.02282-16